



Fundamentals of Organic Chemistry

CHEM 109

For Students of Health Colleges

Credit hrs.: (2+1)


King Saud University

College of Science, Chemistry Department

The General Structure of Nucleic Acids

2



- **Nucleic acids** are molecules that allow organisms to **transfer genetic information** from one generation to the next.
- **Nucleic acids**, are linear polymers (chains) made out of units called **nucleotides**.
- Hydrolysis of nucleic acids gives **nucleotides**, which are the building blocks of nucleic acids.



The General Structure of Nucleic Acids




- **Nucleotides** : contain three parts:
 - A Nitrogenous Base (**Nucleobase**)
 - A Five-Carbon Sugar (**Pentose**)
 - A **Phosphate Group**
- **Nucleosides** : contain two parts:
 - A Nitrogenous Base (**Nucleobase**)
 - A Five-Carbon Sugar (**Pentose**)



The General Structure of Nucleic Acids

4


- **The overall structure of the nucleic acid** is a macromolecule with a backbone of sugar molecules connected by phosphate links and with a base attached to each sugar unit.



The General Structure of Nucleic Acids



5

Phosphate group



Pentose Sugar


- **Ribonucleic acid (RNA)** contains the pentose ribose.
- **Deoxyribonucleic acid (DNA)** contains the pentose deoxyribose.




Heterocyclic Base

The sugars also contain a pyrimidine or purine base present on the 1-carbon replacing the hydroxyl group with a base.

the pyrimidines



the purines



The General Structure of Nucleic Acids

Differences Between DNA and RNA Composition

	DNA	RNA
Five-Carbon Sugar	Deoxyribose	Ribose
Nitrogenous Bases	Adenine, Guanine, Cytosine, and Thymine or methyluracil	Adenine, Guanine, Cytosine, and Uracil


The General Structure of Nucleic Acids



7

○ Nucleoside

The **combination of the pentose sugar and** a purine or pyrimidine **base**.



The General Structure of Nucleic Acids


8

- **Nucleotides** are **phosphate esters of nucleosides**.

A hydroxyl group in the sugar part of a nucleoside is esterified with phosphoric acid.

In DNA nucleotides, either the 3' or the 5' hydroxyl group of 2-deoxy-d-ribose is esterified.


- **Nucleotides** can be **hydrolyzed** by aqueous base (or by enzymes) to **nucleosides and phosphoric acid**.



The General Structure of Nucleic Acids

9

- In these abbreviations, **letter d** stands for 2-deoxy-d-ribose, the **next letter** refers to the heterocyclic base, and **MP** stands for monophosphate.



Base	Monophosphate name	Abbreviation
cytosine (C)	2'-deoxycytidine 5'-monophosphate	dCMP
thymine (T)	2'-deoxythymidine 5'-monophosphate	dTMP
adenine (A)	2'-deoxyadenosine 5'-monophosphate	dAMP
guanine (G)	2'-deoxyguanosine 5'-monophosphate	dGMP



The General Structure of Nucleic Acids

- **Nucleic acids** are polynucleotides attached by the phosphate moieties through the 3' and 5' sites on the pentose.
- The **name nucleic acid** is derived from the fact that they are acidic, containing a phosphoric acid moiety, and are found in the nuclei of cells.
- **Pure nucleic acid** was isolated by Levene in the early 1900s.
- He showed that either D-ribose or D-deoxyribose was present in what are now known as **ribonucleic acid (RNA)** and **deoxyribonucleic acid (DNA)**.
- **There are two major types of nucleic acids:**
 - **Deoxyribonucleic acid (DNA)**
 - **Ribonucleic acid (RNA).**

The Primary Structure of DNA


Deoxyribonucleic Acid (DNA)

الملك سعود
King Saud University




11

- In DNA, **2-deoxy-d-ribose** and **phosphate units** alternate in the backbone.
- The 3' hydroxyl of one ribose unit is linked to the 5' hydroxyl of the next ribose unit by a **phosphodiester bond**.
- The **heterocyclic base** is connected to the anomeric carbon of each deoxyribose unit by a **β -N-glycosidic bond**.
- In DNA, there are **no remaining hydroxyl groups** on any deoxyribose unit.
- Each phosphate, however, still has one acidic proton that is usually ionized **at pH 7**, leaving a negatively charged oxygen.
- A **complete description of any particular DNA molecule**, which may contain thousands or even millions of nucleotide units, would have to include the exact **sequence of heterocyclic bases (A, C, G, and T)** along the chain.




A segment of a DNA chain



Secondary DNA Structure; the Double Helix



Two hydrogen bonds



Three hydrogen bonds



A = T pairs have two hydrogen bonds

G = C pairs have three hydrogen bonds

D = deoxyribose
P = phosphate
A = adenine
T = thymine
G = guanine
C = cytosine

Model and schematic representations of the DNA double helix.

The space-filling model at the left shows the base pairs in the helix interior, in planes perpendicular to the main helical axis. The center drawing shows the structure more schematically, including the dimensions of the double helix. At the far right is a schematic method for showing base pairing in the two strands.

Secondary DNA Structure; the Double Helix

- In **1953**, when **Watson and Crick**, working together in Cambridge, England, proposed the **double helix model for DNA**.
- The important features of their model follow:
 1. DNA consists of **two helical polynucleotide chains coiled** around a common axis.
 2. The **helices are right-handed**, and the two strands run in opposite directions with regard to their 3' and 5' ends.
 3. The **purine and pyrimidine bases lie inside the helix**, in planes perpendicular to the helical axis; the **deoxyribose and phosphate groups form the outside** of the helix.
 4. The **two chains are held together** by;
 - **Purine–pyrimidine base pairs** connected by hydrogen bonds.
 - *Adenine is always paired with thymine, and*
 - *Guanine is always paired with cytosine.*
 5. The **diameter of the helix** is 20 Å.

Adjacent base pairs are separated by 3.4 Å and oriented through a helical rotation of 36°.

There are therefore **10 base pairs for every turn of the helix** (360°), and the structure repeats every 34 Å.
 6. There is no restriction on the sequence of bases along a polynucleotide chain. **The exact sequence carries the genetic information.**

Ribonucleic Acid (RNA)

14

○ Ribonucleic acids (RNA) differ from DNA in three important ways:

(1) The sugar is **D-ribose**;

(2) **Uracil replaces thymine** as one of the four heterocyclic bases

(3) Many RNAs are

- **Single-stranded segments,**
- **Combinations of complementary two-stranded helices,**
- **Complex structures.**

