


Monirah A. Al-Alshaikh, Hazem A. Ghabbour, Mohammed S. M. Abdelbaky,  
Santiago García-Granda and Ali A. El-Emam\*

# Crystal structure of 6-(2-fluorophenyl)-3-phenyl-[1,2,4]-triazolo[3,4-*b*][1,3,4]thiadiazole, C<sub>15</sub>H<sub>9</sub>FN<sub>4</sub>S



**Table 1:** Data collection and handling.

|   |  |
|---|--|
| Crystal:  | Colourless, prism,<br>size 0.1467 × 0.1714 × 0.3504 mm |
| Wavelength:   | Cu K $\alpha$ radiation (1.54184 Å)                    |
| $\mu$ :   | 22.9 cm <sup>−1</sup>                                  |
| Diffractometer, scan mode:                              | Xcalibur, Ruby, Gemini, $\omega$ scans                 |
| 2 $\theta_{\max}$ :                                     | 140.94°  |
| $N(hkl)_{\text{measured}}$ , $N(hkl)_{\text{unique}}$ : | 6872, 2510   |
| Criterion for $I_{\text{obs}}$ , $N(hkl)_{\text{gt}}$ : | $I_{\text{obs}} > 2 \sigma(I_{\text{obs}})$ , 2044     |
| $N(\text{param})_{\text{refined}}$ :                    | 190  |
| Programs:   | CrysAlis <sup>PRO</sup> [16], SHELX [17]               |

DOI 10.1515/ncrs-2015-0236

Received January 25, 2016; accepted February 9, 2016; available  
online February 26, 2016

## Abstract

C<sub>15</sub>H<sub>9</sub>FN<sub>4</sub>S, orthorhombic, *Pna*2<sub>1</sub> (no. 33),  $a = 18.9361(2)$  Å,  $b = 11.5248(1)$  Å,  $c = 6.0142(1)$  Å,  $V = 1312.52(3)$  Å<sup>3</sup>,  $Z = 4$ ,  $R_{\text{gt}}(F) = 0.0263$ ,  $wR_{\text{ref}}(F^2) = 0.0706$ ,  $T = 100$  K.

**CCDC no.:** 1416454

The crystal structure is shown in the figure. Tables 1–3 contain details of the measurement method and a list of the atoms including atomic coordinates and displacement parameters.

**Table 2:** Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å<sup>2</sup>).

| Atom  | Site       | <i>x</i> | <i>y</i> | <i>z</i> | <i>U</i> <sub>iso</sub> |
|-------|------------|----------|----------|----------|-------------------------|
| H(12) | 4 <i>a</i> | 0.2635   | 0.6295   | −0.5280  | 0.042                   |
| H(5)  | 4 <i>a</i> | 0.5152   | 0.0662   | 0.9684   | 0.041                   |
| H(13) | 4 <i>a</i> | 0.3812   | 0.6383   | −0.6289  | 0.047                   |
| H(15) | 4 <i>a</i> | 0.4320   | 0.4358   | −0.1120  | 0.037                   |
| H(2)  | 4 <i>a</i> | 0.4707   | 0.2752   | 0.3255   | 0.036                   |
| H(6)  | 4 <i>a</i> | 0.4001   | 0.1159   | 0.8790   | 0.036                   |
| H(3)  | 4 <i>a</i> | 0.5853   | 0.2271   | 0.4230   | 0.042                   |
| H(4)  | 4 <i>a</i> | 0.6076   | 0.1239   | 0.7454   | 0.040                   |
| H(14) | 4 <i>a</i> | 0.4649   | 0.5392   | −0.4229  | 0.047                   |

## Source of material

A mixture of 4-amino-5-phenyl-4*H*-1,2,4-triazole-3-thiol (1.92 g, 0.01 mol), 2-fluorobenzoyl chloride (1.40 g, 0.01 mol) and phosphorous oxychloride (10 mL) was heated under reflux for four hours. On cooling, the reaction mixture was cautiously poured onto crushed ice (50 gm) and the precipitated solid product was filtered, washed with saturated sodium hydrogen carbonate solution and then with water, dried, crystallized from ethanol to yield 2.16 g (73%) of the title compound. M.p. 517–519 K. Colourless prismatic crystals were obtained by slow evaporation of chloroform-ethanol solution (1:1) at room temperature. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500.13 MHz):  $\delta$  7.16–7.33 (m, 6H, aromatic-H), 7.55–7.77 (m, 3H, aromatic-H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 125.76 MHz):  $\delta$  = 115.11, 120.73, 124.27, 127.54, 127.61, 127.77, 133.45, 133.52, 144.57, 158.43 (aromatic-C), 160.05 (C-3), 160.46 (C-8), 163.96 (C-6).

\*Corresponding author: Ali A. El-Emam, Department of Pharmaceutical Chemistry, College of Pharmacy, King Saud University, P. O. Box 2457, Riyadh 11451, Saudi Arabia, e-mail: elemam5@hotmail.com

Monirah A. Al-Alshaikh: Department of Chemistry, College of Sciences, King Saud University, Riyadh 11451, Saudi Arabia

Hazem A. Ghabbour: Department of Pharmaceutical Chemistry, College of Pharmacy, King Saud University, P. O. Box 2457, Riyadh 11451, Saudi Arabia; and Department of Medicinal Chemistry, Faculty of Pharmacy, Mansoura University, Mansoura 35516, Egypt

Mohammed S. M. Abdelbaky and Santiago García-Granda: Departamento de Química Física Analítica, Facultad de Química, Universidad de Oviedo – CINN, C/Julán Clavería, 8, 33006 Oviedo, (Asturias), Spain

**Table 3:** Fractional coordinates and atomic displacement parameters (Å<sup>2</sup>).

| Atom  | Site | <i>x</i>   | <i>y</i>   | <i>z</i>   | <i>U</i> <sub>11</sub> | <i>U</i> <sub>22</sub> | <i>U</i> <sub>33</sub> | <i>U</i> <sub>12</sub> | <i>U</i> <sub>13</sub> | <i>U</i> <sub>23</sub> |
|-------|------|------------|------------|------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|
| S(1)  | 4a   | 0.22196(2) | 0.39409(3) | 0.17939(9) | 0.0203(2)              | 0.0243(2)              | 0.0278(2)              | 0.0004(1)              | −0.0020(2)             | −0.0002(2)             |
| F(1)  | 4a   | 0.20979(6) | 0.5248(1)  | −0.2014(2) | 0.0355(6)              | 0.0347(6)              | 0.0354(7)              | 0.0103(5)              | −0.0101(5)             | −0.0024(5)             |
| N(3)  | 4a   | 0.32725(7) | 0.2993(1)  | 0.3665(3)  | 0.0216(7)              | 0.0203(6)              | 0.0201(8)              | −0.0016(6)             | 0.0010(7)              | 0.0000(6)              |
| N(4)  | 4a   | 0.35789(7) | 0.3513(1)  | 0.1858(3)  | 0.0241(7)              | 0.0211(6)              | 0.0184(7)              | −0.0016(5)             | 0.0000(8)              | 0.0000(7)              |
| N(1)  | 4a   | 0.29497(7) | 0.2115(1)  | 0.6688(3)  | 0.0273(7)              | 0.0224(6)              | 0.0250(8)              | −0.0034(5)             | 0.0043(8)              | 0.0000(7)              |
| N(2)  | 4a   | 0.23441(8) | 0.2597(1)  | 0.5748(3)  | 0.0275(8)              | 0.0250(7)              | 0.0252(9)              | −0.0038(6)             | 0.0004(8)              | −0.0001(7)             |
| C(12) | 4a   | 0.2976(1)  | 0.5912(2)  | −0.4444(4) | 0.062(1)               | 0.0189(8)              | 0.023(1)               | 0.0032(9)              | −0.014(1)              | −0.0028(8)             |
| C(10) | 4a   | 0.3273(1)  | 0.4703(1)  | −0.1265(3) | 0.031(1)               | 0.0186(7)              | 0.021(1)               | −0.0038(7)             | −0.0054(8)             | −0.0013(7)             |
| C(5)  | 4a   | 0.5062(1)  | 0.1091(2)  | 0.8405(4)  | 0.045(1)               | 0.030(1)               | 0.029(1)               | 0.0049(8)              | −0.013(1)              | 0.0051(8)              |
| C(13) | 4a   | 0.3675(1)  | 0.5957(2)  | −0.5047(4) | 0.068(2)               | 0.027(1)               | 0.023(1)               | −0.015(1)              | −0.003(1)              | 0.0035(8)              |
| C(15) | 4a   | 0.3979(1)  | 0.4750(2)  | −0.1939(4) | 0.032(1)               | 0.034(1)               | 0.026(1)               | −0.0076(8)             | −0.0042(8)             | 0.0050(8)              |
| C(11) | 4a   | 0.2789(1)  | 0.5289(2)  | −0.2584(4) | 0.037(1)               | 0.0195(8)              | 0.027(1)               | 0.0018(7)              | −0.0095(9)             | −0.0057(7)             |
| C(7)  | 4a   | 0.3502(1)  | 0.2355(2)  | 0.5453(3)  | 0.030(1)               | 0.0175(8)              | 0.0209(9)              | −0.0016(7)             | −0.0012(8)             | 0.0001(7)              |
| C(2)  | 4a   | 0.4793(1)  | 0.2341(2)  | 0.4558(4)  | 0.028(1)               | 0.0327(9)              | 0.030(1)               | 0.0030(8)              | −0.0001(9)             | 0.0091(8)              |
| C(8)  | 4a   | 0.25608(9) | 0.3114(2)  | 0.3954(3)  | 0.0207(8)              | 0.0206(8)              | 0.028(1)               | −0.0015(7)             | 0.0002(8)              | −0.0028(8)             |
| C(6)  | 4a   | 0.4371(1)  | 0.1384(2)  | 0.7867(4)  | 0.036(1)               | 0.0278(9)              | 0.026(1)               | −0.0001(8)             | 0.0013(9)              | 0.0035(8)              |
| C(1)  | 4a   | 0.4233(1)  | 0.2018(1)  | 0.5933(4)  | 0.0277(9)              | 0.0201(8)              | 0.0250(9)              | −0.0004(7)             | −0.0024(8)             | −0.0022(7)             |
| C(3)  | 4a   | 0.5480(1)  | 0.2048(2)  | 0.5141(4)  | 0.026(1)               | 0.038(1)               | 0.040(1)               | 0.0023(8)              | −0.001(1)              | 0.004(1)               |
| C(4)  | 4a   | 0.5615(1)  | 0.1427(2)  | 0.7067(4)  | 0.0296(9)              | 0.0313(9)              | 0.040(1)               | 0.0044(7)              | −0.010(1)              | −0.0024(9)             |
| C(14) | 4a   | 0.4177(1)  | 0.5370(2)  | −0.3806(4) | 0.045(1)               | 0.044(1)               | 0.030(1)               | −0.016(1)              | 0.004(1)               | 0.0048(9)              |
| C(9)  | 4a   | 0.30898(9) | 0.4043(1)  | 0.0730(3)  | 0.0220(9)              | 0.0191(7)              | 0.024(1)               | −0.0004(6)             | −0.0004(8)             | −0.0031(7)             |

## Experimental details

Cell refinement and data reduction were carried out by CrysAlis PRO [16]. The coordinates of the aromatic H atoms were idealized and refined using a riding model (AFIX 43 option of the SHELX program [17]).

## Discussion

1,2,4-Triazole derivatives and their fused heterocyclic analogues are well known for their different biological activities over 30 years ago, and 1,2,4-triazole rings have been incorporated into ligands used in coordination compounds and polymers. Thus, various 1,2,4-triazole derivatives and their *N*-bridged heterocyclic analogues have been extensively studied [1–6]. Several 3,6-disubstituted[1,2,4]triazolo[3,4-*b*][1,3,4]thiadiazoles were reported to exhibit significant antibacterial [7–10], pesticidal [11], anticancer [12], anti-inflammatory, analgesic and anti-oxidant activities [13]. In continuation to a previous interest in the chemical synthesis of [1,2,4]-triazolo[3,4-*b*][1,3,4]thiadiazole derivatives [14, 15], we report herein the synthesis and the crystal structure of the title compound as potential bioactive agent.

One independent molecule comprises the asymmetric unit. The compound is nearly planar, with respect to the [1,2,4]-triazolo[3,4-*b*][1,3,4]thiadiazole system (N1/N2/C8/S1/C9/N4/N3/C7), the fluorophenyl ring (C10–C15) form dihedral angles of 2.55 (6)° and the phenyl ring (C1–C6) form dihedral angles of 3.83 (3)°. In the crystal

structure, the packing is stabilized by one non-classical intermolecular hydrogen bond, of which the N1 acts as hydrogen bond acceptor and the C12 acts as hydrogen bond donor. The distance of the interaction between C12–H12···N1 is 2.33 Å and the angle is 161°.

**Acknowledgements:** The authors would like to extend their sincere appreciation to the Deanship of Scientific Research at King Saud University for funding this work through the Research Group Project No. PRG-1436–23. We also acknowledge the financial support from Spanish Ministerio de Economía y Competitividad (MINECO-13-MAT2013–40950-R, FPI grant BES-2011–046948 to MSM-A).

## References

- Chai, X.; Zhang, J.; Yu, S.; Hu, H.; Zou, Y.; Zhao, Q.; Dan, Z.; Zhang, D.; Wu, Q.: Design, synthesis, and biological evaluation of novel 1-(1*H*-1,2,4-triazole-1-yl)-2-(2,4-difluorophenyl)-3-substituted benzylamino-2-propanols. *Bioorg. Med. Chem. Lett.* **19** (2009) 1811–1814.
- El-Emam, A. A.; Ibrahim, T. M.: Synthesis, anti-inflammatory and analgesic activity of certain 3-(1-adamantyl)-4-substituted-5-mercapto-1,2,4-triazole derivatives. *Arzneim.-Forsch./Drug Res.* **41** (1991) 1260–1264.
- Aboelmagd, A.; Ali, I. A. I.; Salem, E. M. S.; Abdel-Razik, M.: Synthesis and antifungal activity of some *s*-mercaptotriazolobenzothiazolyl amino acid derivatives. *Eur. J. Med. Chem.* **60** (2013) 503–511.

4. El-Emam, A. A.; Al-Tamimi, A.-M. S.; Al-Omar, M. A.; Al-Rashood, K. A.; Habib, E. E.: Synthesis and antimicrobial activity of novel 5-(1-adamantyl)-2-aminomethyl-4-substituted-1,2,4-triazoline-3-thiones. *Eur. J. Med. Chem.* **68** (2013) 96–102.
5. Luo, Y.; Zhang, S.; Liu, Z.-J.; Chen, W.; Fu, J.; Zeng, Q.-F.: Synthesis and antimicrobial evaluation of a novel class of 1,3,4-thiadiazole: derivatives bearing 1,2,4-triazolo[1,5-*a*]pyrimidine moiety. *Eur. J. Med. Chem.* **64** (2013) 54–61.
6. Plech, T.; Wujec, M.; Kosikowska, U.; Malm, A.; Kaproń, B.: Studies on the synthesis and antibacterial activity of 3,6-disubstituted 1,2,4-triazolo[3,4-*b*]1,3,4-thiadiazoles. *Eur. J. Med. Chem.* **47** (2012) 580–584.
7. Eweiss, N. F.; Bahajaj, A. A.: Synthesis of heterocycles. Part VII. Synthesis and antimicrobial activity of some 7*H*-s-triazolo[3,4-*b*][1,3,4]thiadiazine and s-triazolo[3,4-*b*][1,3,4]thiadiazole derivatives. *J. Heterocycl. Chem.* **24** (1987) 1173–1181.
8. Kotaiah, Y.; Nagaraju, K.; Harikrishna, N.; Rao, C. V.; Yamini, L.; Vijjulatha, M.: Synthesis, docking and evaluation of antioxidant and antimicrobial activities of novel 1,2,4-triazolo[3,4-*b*][1,3,4]thiadiazol-6-yl)selenopheno[2,3-*d*]pyrimidines. *Eur. J. Med. Chem.* **75** (2014) 195–202.
9. Swamy, S. N.; Basappa, Priya, B. S.; Prabhuswamy, B.; Doreswamy, B. H.; Shahidhara, J. S.; Rangappa, K. S.: Synthesis of pharmaceutically important condensed heterocyclic 4,6-disubstituted-1,2,4-triazolo-1,3,4-thiadiazole derivatives as antimicrobials. *Eur. J. Med. Chem.* **41** (2006) 531–538.
10. Mathew, V.; Keshavayya, J.; Vaidya, V. P.: Heterocyclic system containing bridged nitrogen atom: synthesis and pharmacological activities of some substituted 1,2,4-triazolo[3,4-*b*]-1,3,4-thiadiazoles. *Eur. J. Med. Chem.* **41** (2006) 1048–1058.
11. Chaturvedi, B.; Tiwari, N.; Nirupama, N.: A convenient synthesis of 1,2,4-triazolo[3,4-*b*][1,3,4]thiadiazoles as potential pesticides. *Agri. Biol. Chem.* **52** (1988) 1229–1232.
12. Khan, I.; Zaib, S.; Ibrar, A.; Rama, N. S.; Simpson, J.; Iqbal, J.: Synthesis, crystal structure and biological evaluation of some novel 1,2,4-triazolo[3,4-*b*]-1,3,4-thiadiazoles and 1,2,4-triazolo[3,4-*b*]-1,3,4-thiadiazines. *Eur. J. Med. Chem.* **78** (2014) 167–177.
13. Chidananda, N.; Poojary, B.; Sumangala, V.; Kumari, N. S.; Shetty, P.; Arulmoli, T.: Facile synthesis, characterization and pharmacological activities of 3,6-disubstituted 1,2,4-triazolo[3,4-*b*][1,3,4]thiadiazoles and 5,6-dihydro-3,6-disubstituted-1,2,4-triazolo[3,4-*b*][1,3,4]thiadiazoles. *Eur. J. Med. Chem.* **51** (2012) 124–136.
14. Al-Abdullah, E. S.; Shehata, I. A.; Al-Deeb, O. A.; El-Emam, A. A.: Microwave-assisted dehydrosulphurization: an efficient, solvent-free synthesis of 5-(1-adamantyl)-2-arylamino-1,2,4-triazolo[3,4-*b*][1,3,4]thiadiazoles. *Heterocycles* **71** (2007) 379–388.
15. El-Emam, A. A.; Moustafa, M. A.; El-Subbagh, H. I.; El-Ashmawy, M. B.: Triazoles and fused triazoles III: Facile and efficient synthesis of 2,4-disubstituted-s-triazolo[3,4-*b*]-1,3,4-thiadiazoles. *Monatsh. Chem.* **121** (1990) 221–225.
16. Agilent. CrysAlis<sup>PRO</sup>. Agilent Technologies UK Ltd, Yarnton, England, 2014.
17. Sheldrick, G. M.: A short history of SHELX. *Acta Crystallogr. A* **64** (2008) 112–122.