

Stat true or false:-

(i)  $2i - 5j$  is a unit vector.

(ii)  $ax + by + d = 0$ , is orthogonal to the  $xy$ -plane.

(iii)  $(i \times j) \times k = 1$ .

(iv) Area of the circle  $r = 5, 0 \leq \theta \leq 2\pi$ , is  $\frac{1}{2} \int_0^\pi 5^2 d\theta$ .

(v) The graph of  $r = \theta$  is a circle.

(vi)  $\left(\vec{a} \times \vec{b}\right)^2 = a^2 b^2 - \left(\vec{a} \cdot \vec{b}\right)^2$ .

(i) Find the length of the curve,  $x = 2 - \ln t$ ,  $y = 3 \ln t$ , sketch the graph of the curve, and indicate the orientation  $1 \leq t \leq e$ .

(ii) Find a polar equation for  $(x^2 + y^2)^2 + 4xy = 0$ , sketch the graph.

(iii) Find area of the triangle determined by  $P(1, 0, -5)$ ,  $Q(-2, 1, 0)$ , and  $R(3, 2, 1)$ .

(i) Find an equation of the plane through  $A(0, 4, 9)$ , and  $B(0, -2, 6)$  and perpendicular to  $xy$ -plane, Find the distance from  $C(1, 0, -1)$  to the plane.

(ii) Sketch the graph of  $r = 1 + 2 \cos \theta$ , find the slope of the tangent line at  $\theta = \frac{\pi}{2}$ , and find the area of the region bounded by  $r = 1 + 2 \cos \theta$ .

(iii) If  $\vec{a} = 3i - j - 4k$ ,  $\vec{b} = 2i + 5j - 2k$ , and  $\vec{c} = -i + 6k$ , find

\*  $\text{comp}_{\vec{b}} \left( \vec{b} \times \vec{c} \right)$ .

\* A vector having opposite direction of  $\vec{a}$  and twice magnitude.

\* the volume of the box determined by  $\vec{a}$ ,  $\vec{b}$ , and  $\vec{c}$ .

---

**Q1- Prove or disprove:-**

(i)  $\|\vec{a} + \vec{b}\| \leq \|\vec{a}\| + \|\vec{b}\|.$

(ii) The graph of the polar equation  $r = 2\cos 4\theta$ , has 4 loops.

(iii) The distance from the point  $(1,0,-2)$  to the plane  $: 2x - 3y + z = 6$ , is  $\frac{6}{\sqrt{14}}$

(iv)  $r = 4\cos\left(\theta - \frac{\pi}{2}\right)$  is a circle.

(v) The curvature of  $x = 2\cos t$  and  $y = 2\sin t$  is 2 for all  $t$ .

(vi)  $\vec{a} \times (\vec{b} \times \vec{c}) + \vec{b} \times (\vec{c} \times \vec{a}) + \vec{c} \times (\vec{a} \times \vec{b}) = \vec{0}.$

(vii) The line  $x = 1 + 5t, y = 1 - 2t, z = 4 + t$ , and the plane  $2x + 3y - 4z = 5$ , are perpendicular.

(iix) If  $\vec{v}$  is a nonzero vector,  $a = 0$  be scalar, If  $\vec{u}$  is any vector then

$$\text{proj}_{\vec{v}} \vec{u} = \text{proj}_{a\vec{v}} \vec{u}.$$

(ix)  $\oint_c (y - 7e^{x^3})dx + (x + \ln \sqrt{y})dy = 0.$

Where  $c$  is the ellipse,  $\frac{x^2}{9} + \frac{y^2}{4} = 1$

(x)  $\iiint_Q \nabla \cdot \vec{n} dV = S$ , where  $\vec{n}$  is a unit normal vector to any closed surface.

---

**Q2-**

(i) Use Green's Theorem to evaluate  $\oint_c xydx + \sin ydy$ , where  $c$  is the graph of the triangle with vertices  $(1,0)$ ,  $(1,1)$  and  $(0,0)$ .

(ii) Find velocity, acceleration at  $t = 1$ , for the curve,  $\vec{r}(t) = e^{2t}i + e^{-2t}j$ , and sketch them.

(iii) Find the point on the curve at which the tangent line is, horizontal, vertical:  
 $x = t^3 - 4t$ ,  $y = t^2 - 4$ .

-----Q3-

(i) Use the divergence theorem to evaluate  $\iiint_s \vec{F} \cdot \vec{n} ds$ , over the closed surface bounded by  $x^2 + y^2 = 9$ ,  $z = 0$ , and  $z = 5$ .

$$\vec{F} = (x^2 + z^2)i + (y^2 - 2xy)j + (4z - 2yz)k.$$

(ii) If  $\vec{F} = y^2i + 2xj + 5yk$ , verify stokes theorem, where  $s$  : is the hemisphere

$$z = (4 - x^2 - y^2)^{\frac{1}{2}}.$$

(iii) sketch the curve  $r^2 = 4 \cos 2\theta$ . Find the area of the region bounded by the above curve.

-----Q4-(i)

Change the rectangular co-ordinate to a) spherical, b) cylindrical co-ordinates.  
 ( without calculating the integral )

$$\int_{-1}^1 \int_{-\sqrt{1-y^2}}^{\sqrt{1-y^2}} \int_{\sqrt{x^2+y^2}}^1 dz dx dy.$$

(ii) Show that,  $\int_{(2,1,2)}^{(-1,0,4)} (yz + 2)dx + (xz - 3)dy + (xy + 5)dz$ , is independent of path and find its volume.

=====

**M-202**

**FINAL EXAMINATION SEMESTER – II (1430-1431)**

*NOTE: ANSWER ALL QUESTION.*

**Q<sub>1</sub>:**

- a) Use the spherical coordinates to find the volume of the solid bounded by cone  $z = \sqrt{x^2 + y^2}$  and the cylinder  $x^2 + y^2 = 4$ .
- b) Show that: i.  $\text{div}(\text{curl}\vec{F}) = 0$ . ii.  $(\vec{a} \times \vec{b}) \times (\vec{c} \times \vec{d}) = (\vec{a} \times \vec{b} \cdot \vec{d})\vec{c} - (\vec{a} \times \vec{b} \cdot \vec{c})\vec{d}$ .
- c) Find the initial point of  $\vec{V} = -2\vec{i} + 4\vec{j}$ , if terminal point is  $(2,0)$ .
- 
- 

**Q<sub>2</sub>:**

- a) Find the unit vector perpendicular to each of the vectors  $2\vec{i} - \vec{j} + \vec{k}$  and  $3\vec{i} + 4\vec{j} - \vec{k}$  and calculate the  $\sin$  of the angle between the two vectors.
- b) Find an equation of the plane through  $P(4,2,-9)$  with normal vector  $\overrightarrow{OP}$ .
- 
- 

**Q<sub>3</sub>:**

- a) Find the area of the region that inside  $r = 2$ , and outside  $r = 2 + 2\cos\theta$ .
- b) If  $\vec{r}(t) = \sin 2t\vec{i} + \cos t\vec{j}$  is the position vector for a point, find the tangential and normal components of acceleration at the time  $t$ .
- c) Find the slope of the tangent line to the graph  $x = \sin t, y = \cos t$  at  $t = \frac{\pi}{4}$  and sketch the graph.
- 
- 

**Q<sub>4</sub>:**

- a) Show that the line integral  $\int_{(0,0)}^{\left(1, \frac{x}{2}\right)} e^x \sin y dx + e^x \cos y dy$  is independent of paths and find its value.
- b) Use Green's theorem to evaluate  $\oint_C (x^2 + 4)dx + xydy$  where  $C$  is the cardioid  $r = 1 + \cos\theta$ .
- c) Find the surface area of the part of the graph  $z = xy$  that inside the cylinder  $x^2 + y^2 = 4$ .
- 
- 

**Q<sub>5</sub>:**

- a) State the divergence theorem and show that  $\frac{1}{3} \iint_S \vec{r} \cdot \vec{n} ds = V$  where  $\vec{r} = xi + yj + zk$  and  $V$  is the volume enclosed by the surface  $S$ .

Verify the Stoke theorem for  $F = zi + xj + yk$  and  $S$ , where  $S$  is the hemisphere

$$Z = (a^2 - x^2 - y^2)^{\frac{1}{2}}.$$

**1430-1431 H**

**202 Math**

$Q_1$  :-State true or false:-

(i)  $3i - j$  is a unit vector.

(ii)  $\|\vec{u} \times \vec{v}\| = \|\vec{u}\|^2 \|\vec{v}\|^2 - (\vec{u} \cdot \vec{v})^2$

(iii)  $(i \times j) \times k = 1$ .

(iv) Area of the circle  $r = 4, 0 \leq \theta \leq 2\pi$ , is  $\frac{1}{2} \int_0^\pi 16 d\theta$ .

(v) The vectors  $2i - 3j$  and  $6i + 4j$  are perpendicular.

(vi)  $|\vec{r}(t)|$  is constant if and only if  $\vec{r}(t) \cdot \vec{r}'(t) = 0$ .

$Q_2$  :-

(i) Find the length of the curve,  $\vec{r}(t) = \sinh 2ti + \cosh 2tj + 2tk$ ,  $-\frac{1}{2} \leq t \leq 1$ .

(ii) Find apolar equation for  $y^2 = 4px$ , sketch the graph.

(iii) Find the area of the triangle determined by P(3,2,-1), Q(2,4,6), and R(-1,2,7).

(iv) Find the area of the region outside the cardioid  $r = 1 + \cos \theta$  and inside the circle  $r = \sqrt{3} \sin \theta$ .

$Q_3$  :-

- (i) Find the angle between the vectors  $\langle 4, -3, -2 \rangle$  and  $\langle -1, 2, 5 \rangle$ .
- (ii) Find  $\text{proj}_{\vec{v}} \vec{u}$  if  $\vec{u} = -4i + j - 2k$ ,  $\vec{v} = i + 3j - 3k$ .
- (iii) Find the volume of the box determined by  $(-1, 2, 3)$ ,  $(4, -1, 2)$ ,  $(5, 6, 3)$  and  $(1, 1, -2)$ .
- (iv) Find all vectors perpendicular to both vectors  $-2i + j - 4k$  and  $3i - 4j + 5k$ .
- 

**1430-1431 H**

**202 Math**

---

$Q_1$  :- State true or false:-

- (i)  $3i + 4j$  is a unit vector.
- (ii)  $\|\vec{u} \times \vec{v}\| = \|\vec{u}\|^2 \|\vec{v}\|^2 - (\vec{u} \cdot \vec{v})^2$
- (iii)  $i \times (j \times k) = 0$ .
- (iv) Area of the circle  $r = 5, 0 \leq \theta \leq \pi$ , is  $\frac{1}{2} \int_0^\pi 25 d\theta$ .
- (v) The vectors  $2i + 3j$  and  $6i + 4j$  are perpendicular.
- (vi)  $|\vec{r}(t)|$  is constant if and only if  $\vec{r}(t) \cdot \vec{r}'(t) = 0$ .
- 

$Q_2$  :-

- (i) Find the length of the curve,  $\vec{r}(t) = \sinh 2ti + \cosh 2tj + 2tk$ ,  $-\frac{1}{2} \leq t \leq 1$ .
- (ii) Find a polar equation for  $x^2 = 4py$ , sketch the graph.
- (iii) Find the area of the triangle determined by P(2, -1, 1), Q(1, 0, -2), and R(-1, 1, 3).
- (iv) Find the area of one loop of  $r = \sin 3\theta$

---

$Q_3$ :-

- (i) Find the angle between the vectors  $\langle 2, -2, 1 \rangle$  and  $\langle -2, 0, 1 \rangle$ .
- (ii) Find  $\text{proj}_{\vec{v}} \vec{u}$  if  $\vec{u} = 2i + 4j + 2k$ ,  $\vec{v} = -i + 2j + k$ .
- (iii) Find the volume of the box determined by  $(1, -2, -1)$ ,  $(2, 0, -2)$ ,  $(3, 2, -1)$  and  $(-1, 2, -2)$ .
- (iv) Find all vectors perpendicular to both vectors  $-3i + j + 4k$  and  $2i + 4j - 3k$ .
- 

### First Midterm Examination

Second term 1430-1431 H

Course no, 202 (calculus)

Answer all questions :

$Q_1$  : State "true" or "false" to the following statements :-

- Orientation of a curve is determined by decreasing values of the parameter  $t$ .
- The dot product of two parallel vectors is zero.
- $\|\vec{a} + \vec{b}\| = \|\vec{a}\| + \|\vec{b}\|$  when  $\vec{a}$  and  $\vec{b}$  are orthogonal to each other.
- If any two of  $\vec{a}, \vec{b}$  and  $\vec{c}$  are equal, then  $\vec{a} \cdot \vec{b} \wedge \vec{c} = 0$ .

$Q_2$  :

- Sketch the graph of the polar equation  $r = 3 + 2\cos\theta$  and find its area. Also find the slope of the tangent line to the graph at  $\theta = \frac{\pi}{2}$ .
- Find the area of the triangle determined by  $\vec{a}$  and  $\vec{b}$  if  $\vec{a} = 3i - j - 4k$  and  $\vec{b} = 2i + 5j - 2k$ .

$Q_2$  :

- Find the values of  $t$  for  $\vec{u} = t^2i + 6tj + tk$  and  $\vec{v} = 5ti - 5tj + 4t^2k$  are orthogonal.
- Find a vector of magnitude 4 that has the opposite direction of  $\vec{a} = \langle 2, -5, 0 \rangle$ .

Curve  $C$  is given parametrically  $x = t^2 + 1, y = t^2 - 1, -2 \leq t$ . Sketch the graph and indicate the orientation.

First semester

202 Math

King saud university

Second midterm

1430-1431 H

Q<sub>1</sub>

a) Answer true or false :-

- i. The line  $x = 1 + 5t, y = 1 - 2t, z = 4 + t$  and the plane  $2x + 3y - 4z = 1$  are perpendicular
- ii. A particle whose position  $\vec{r}(t) = \cos t \mathbf{i} + \cos t \mathbf{j} + \sqrt{2} \sin t \mathbf{k}$  moves with constant speed.

b) Fill in the blanks

- i. A vector that is normal to the plane  $-6x + y - 7z + 10 = 0$  is .....
- ii. The equation of the plane that has  $x$ -intercept and  $z$ -intercept at 8 at 4,  $y$ -intercept at 2 and  $z$ -intercept at 8 is .....

Q<sub>2</sub>

Evaluate

- i.  $\int \frac{1}{1+t^2} (i + t \mathbf{j} + \tan^{-1} t \mathbf{k}) dt$
- ii.  $\int_0^2 \int_x^{\sqrt{8-x^2}} \frac{1}{5+x^2+y^2} dy dx$
- iii. Find the surface area of that portion of the paraboloid  $z = x^2 + y^2$  that is below the plane  $z = 2$

Q<sub>3</sub>

- i. Graph the curve and the velocity and acceleration vector of  $\vec{r}(t) = t^2 \mathbf{i} + \frac{1}{4} t^4 \mathbf{j}$  at  $t = 1$ .
- ii. Find the curvature and radius of curvature of  $y = e^x - x$  at the point  $(0, 1)$
- iii. Find parametric equations for the line of intersection of

$$2x - 3y + 4z = 1$$

$$x - y - z = 5$$



---

**Q1** Prove or disprove :-

(i) The volume of the sphere  $x^2 + y^2 + z^2 = 1$  , is

$$V = \int_0^{2\pi} \int_0^{\pi} \int_0^1 \rho^2 \sin \varphi \, d\rho \, d\theta \, d\varphi .$$

(ii) Curvature at any point on the circle  $r = 2\cos \theta$  is zero .

(iii) Every straight line has curvatre equal to 1 .

(iv) Let  $\vec{F} = 2x\sin y\vec{i} + x^2\cos y\vec{j}$  then  $\nabla \times \vec{F}$  is  $\vec{0}$  .

(vi) If a particle is moving at a constant speed then the tangential component at acceleration is zero .

---

**Q2**

(a) Convert  $(0, -1, 0)$  to :-

(i) Cylindrical ,                      (ii) Spherical co-ordinates .

(b) Evaluate ,  $\int_0^a \int_0^{\sqrt{a^2-x^2}} \int_0^{\sqrt{a^2-x^2-y^2}} \frac{z^3}{\sqrt{x^2+y^2}} dz \quad dy \quad dx$

(c) Find the surface area of the part of the graph  $z = xy$  , that inside the cylinder  $x^2 + y^2 = 4$  .

(d) The position vector of a particle at time  $t$  is  $\vec{r}(t) = ti + t^2j + t^3k$

for  $1 \leq t \leq 4$ .

Find the tangential and normal components of acceleration at time  $t = 1$ .

---

### Q3

(a) Sketch  $\vec{r}(t), \vec{r}'(t)$  and  $\vec{r}''(t)$  at  $t = \frac{3\pi}{4}$  , where  $\vec{r}(t) = 4\cos ti + 7\sin tj$ , is the curve .

(b) Evaluate  $\int_c (3x^2 - 8y^2)dx + (4y - 6xy)dy$  where  $c$  is the boundary of the region bounded by  $x = 0, y = 0, x + y = 1$  .

(c) Find the curvature , radius of curvature and centre of curvature for  $xy = 1$ , at the point  $\left(2, \frac{1}{2}\right)$ .

**First Semester 1432-1433H**

**King Saud University**

**Final Examination**

**Faculty of Science**

**202 Math**

**Mathematic Department**

Question. No.	1	2	3	4	5	6
Answer						

Q<sub>1</sub>(i) Choose the correct answer:-

1) The vectors  $\langle -4, -6, 10 \rangle$  and  $\langle -10, -15, 25 \rangle$  are:

None of these (iii) Orthogonal (ii) Parallel (i)

2) The line  $x = 1 + 5t$ ,  $y = 1 - 2t$ ,  $z = 4 + t$  and the plane  $2x + 3y - 4z = 1$  are:

None of these (iii) Perpendicular (ii) Parallel (i)

3) If  $\lim_{t \rightarrow a} \vec{r}_1(t) = 2i + j$  and  $\lim_{t \rightarrow a} \vec{r}_2(t) = -i + 2j$ , then  $\lim_{t \rightarrow a} \left( \vec{r}_1(t) \cdot \vec{r}_2(t) \right)$  is equal:

None of these (iii) 1 (ii) 0 (i)

4) If  $f(x, y) = \cos y$ , then:

None of these (iii)  $\nabla f(x, y) = -\sin yj$  (ii)  $\nabla f(x, y) = -\sin y$  (i)

5) If  $\vec{r}(t) = \vec{a} \sin t + \vec{b} \cos t$ , where  $\vec{a}, \vec{b}$  are constant, then  $\vec{r} \times \frac{d\vec{r}}{dt}$  is:

None of these (iii)  $\vec{b} \times \vec{a}$  (ii)  $\vec{a} \times \vec{b}$  (i)

6)  $\oint_C (y - 7e^{-x^2})dx + (x + \ln \sqrt{y})dy$ ,  $C: 2(x-10)^2 + 9(y+13)^2 = 3$  is:

None of these (iii) 0 (ii) 1 (i)

(ii) Prove or disprove:

1) If  $\vec{v}$  is a nonzero vector,  $a \neq 0$  be scalar, If  $\vec{u}$  is any vector then,

$$\text{proj}_{\vec{v}} \vec{u} = \text{proj}_{a\vec{v}} \vec{u}$$

2)  $\iiint_Q \nabla \cdot \vec{n} dV = S$ , where  $\vec{n}$  is a unit normal vector to any closed surface.

3) The distance from the point  $(1,0,-2)$  to the plane  $: 2x - 3y + z = 6$ , is  $\frac{6}{\sqrt{14}}$

4)  $\vec{a} \times (\vec{b} \times \vec{c}) + \vec{b} \times (\vec{c} \times \vec{a}) + \vec{c} \times (\vec{a} \times \vec{b}) = \vec{0}$

Q:

1. If  $\vec{F}(x, y, z) = 4xyz\vec{i} + 2x^2z\vec{j} + 2x^2y\vec{k}$  prove that  $\text{div}(\text{curl } \vec{F}) = 0$

2. If  $\vec{r}(t) = t^2\vec{i} + 2t\vec{j}$  find  $\vec{v}(t)$ ,  $\vec{a}(t)$  and sketch  $\vec{r}(1)$ ,  $\vec{v}(1)$  and  $\vec{a}(1)$

3.

(i) Convert  $\left(8, \frac{\pi}{3}, 7\right)$  in cylindrical coordinates to rectangular coordinates.

- (ii) Convert  $(-\sqrt{2}, \sqrt{2}, 1)$  in rectangular coordinates to cylindrical coordinates.

Q<sub>3</sub>:

1. Determine whether the given pair of lines  $L_1 : x = 1 + t, y = -3 + 2t, z = -2 - t$  and  $L_2 : x = 17 - 3s, y = 4 - s, z = -8 + s$ , has a point of intersection. If so, find it.

2. Find the curvature of the curve described by:  $x = t - \sin t, y = 1 - \cos t$ , at  $t = \frac{\pi}{2}$ .

3. Show that the given line integral is independent of path and find its value

$$\int_{\left(\frac{\pi}{2}, 0\right)}^{(\pi, 1)} \left( e^{3y} - y^2 \sin x \right) dx + \left( 3xe^{3y} + 2y \cos x \right) dy$$

Q.:

- I. Use Green's Theorem to evaluate the line integral  $\int_c (y + e^x)dx + (2x^2 + \cos y)dy$ , where  $c$  is the boundary of the triangle with vertices  $(0,0)$ ,  $(1,1)$ , and  $(1,0)$ .

- II. Use Divergence theorem to evaluate the integral  $\iiint_S \vec{F} \cdot \vec{n} ds$ , where

$\vec{F} = x^3 \vec{i} + y^3 \vec{j} + z^2 \vec{k}$ , and  $S$  is the region enclosed by the cylinder  $x^2 + y^2 = 4$ ,  $z = 0$  and  $z = 2$ .

Q5:

- I. Verify Stoke's Theorem for the vector  $\vec{F}(x, y, z) = 2zi + 3xj + 5yk$ , over the region.  
 $z = 4 - 2x - 4y$

- II. Evaluate  $\iint_S \vec{F} \cdot \vec{n} ds$ , where  $\vec{F}(x, y, z) = xi + yj + zk$  and  $S$  is the paraboloid  $z = 1 - x^2 - y^2$  cut off by  $z = 0$ .

First semester 1432-1433

King Saud University

.III

2nd midterm exam

Mathematical department

.IV

202 Math .V

.VI

Q1. a) Find  $\text{div } \vec{F}$  and  $\text{curl } \vec{F}$  if  $\vec{F}(x, y, z) = x^3 z^4 i + xyz^2 j + x^2 y^2 k$ .

.VII

b) Find the curvature of the curve described by  $y = 2x^2 - x + 2$ .

.VIII

.IX

Q2. a) Evaluate  $\int_c (x - y)dx + xdy$ , where  $c$  is the graph of  $y^2 = x$  from  $(4, -2)$  to  $(4, 2)$ .

.X

b) Find the tangential and normal components of acceleration of a particle moving along the

.XI

curve  $c$  described by  $\vec{r}(t) = (t^2 - 1)i + (t + 1)j + \frac{1}{2}(t^2 + 1)k$ ,  $t = 1$ .



c) Find the surface area of the part of the paraboloid  $z = 4 - x^2 - y^2$ , that lies above the plane  $z = 2$ . .XII

.XIII

Q3. a) Show that the given line integral is independent of path and find its value .XIV

$$\int_{(-1,2)}^{(3,1)} (y^2 + 2xy)dx + (x^2 + 2xy)dy .$$

b) Find the equation for the plane through the point  $(1,4,-5)$  and parallel to the plane  $2x - 5y + 7z = 12$ . .XV