




CHEM 101+103



Second Exam Sample 1

Ahmad Aqel Ifseisi

Assistant Professor of Analytical Chemistry
College of Science, Department of Chemistry
King Saud University

P.O. Box 2455 Riyadh 11451 Saudi Arabia

Building: 05, Office: AA53

Tel. 014674198, Fax: 014675992

Web site: <http://fac.ksu.edu.sa/aifseisi>

E-mail: ahmad3qel@yahoo.com

aifseisi@ksu.edu.sa



كرسي أبحاث
المواد المتقدمة
Advanced Materials
Research Chair



1. The sign of ΔH for the process $\text{CO}_2(\text{s}) = \text{CO}_2(\text{g})$ is: ((the symbol “H” means enthalpy))



A) Positive and $H_{\text{CO}_2}(\text{s}) > H_{\text{CO}_2}(\text{g})$

B) Positive and $H_{\text{CO}_2}(\text{g}) > H_{\text{CO}_2}(\text{s})$

C) Negative and $H_{\text{CO}_2}(\text{s}) > H_{\text{CO}_2}(\text{g})$

D) Negative and $H_{\text{CO}_2}(\text{g}) > H_{\text{CO}_2}(\text{s})$

2. Which of the $\Delta H^\circ_{\text{rxn}}$ of the following equations represents $\Delta H^\circ_{\text{f, K}_3\text{PO}_4(\text{s})}$?



3. A balanced chemical equation with specified value of ΔH and states of substances is called:

- A) A thermochemical equation**
- B) A combustion reaction**
- C) The first law of thermodynamics**
- D) Hess's law**

4. Change in internal energy (ΔE°), in KJ, of the following reaction is:



A) 121.04

B) 134.04

C) 124.04

D) 114.04

5. If 10.0 g of a metal ($C_s = 0.896 \text{ J/g K}$) at 298 K is supplied with 313.5 J of heat, its final temperature, in K, will be:


A) 353

B) 333

C) 323

D) 373

6. From table below, $\Delta H^\circ_{\text{rxn}}$ of the following reaction, in KJ, is:



Compound	$\text{PH}_3(\text{g})$	$\text{PCl}_3(\text{g})$	$\text{HCl}(\text{g})$
$\Delta H_f/\text{KJ mol}^{-1}$	+ 5.40	- 288.07	- 92.30

A) 570.37 B) 507.37 C) 705.37 D) 750.37

7. Knowing that:



the number of kilojoules (KJ) released if 100 g of HCl(g) is produced, is:

A) 235.17

B) 325.17

C) 523.17

D) 253.17

8. The process of surrounding solute particles by solvent particles is known as:

A) Dilution

B) Formation

C) Solvation

D) Osmosis

9. The solubility of?.....in liquid is highly affected by changing pressure

- A) Gases**
- B) Liquids**
- C) Solids**
- D) Salts**

10. If 0.1 mol of solid glucose ($\text{C}_6\text{H}_{12}\text{O}_6$) is dissolved in the same mass of each of the following solvents:

Solvent	Q	X	Y	Z
$K_b/\text{C molal}^{-1}$	0.4	1.53	1.7	0.5

the solvent which its boiling point is elevated more is:

A) Q

B) X

C) Y

D) Z

11. The magnitudes of the molal constant of boiling point elevation (K_b) depend on:

A) Temperature

B) Pressure

C) Nature of solute

D) Nature of solvent

12. The aqueous solution with the highest boiling point is:

A) 0.1 M HI

B) 0.1 M $(\text{NH}_4)_3\text{PO}_4$

C) 0.2 M $\text{C}_2\text{H}_5\text{OH}$

D) 0.1 M NH_4Cl

13. If 1 L carbonated water is bottled under pressure of 2.4 atm of $\text{CO}_2(\text{g})$, and Henry's law constant is $3.36 \times 10^{-2} \text{ mol/L atm}$, the number of grams of dissolved $\text{CO}_2(\text{g})$ is:

A) 5.35

B) 53.5

C) 35.5

D) 3.55

14. At 30 °C, the osmotic pressure, in torr, of 0.108 M aqueous solution of a salt that is assumed to be totally ionized into three ions is:

A) 3.16×10^3

B) 1.63×10^3

C) 6.13×10^3

D) 1.36×10^3

15. The minimum amount of energy required to overcome the energy barrier in a chemical reaction is the:

- A) Activation energy**
- B) Reaction's enthalpy**
- C) Reactant's kinetic energy**
- D) Reactants' heat content**

16. Increasing temperature increases reaction rate because it:

- A) Increases the activation energy
- B) Decreases the activation energy
- C) Increases the number of collisions**
- D) Increases the reaction enthalpy

17. According to the following reaction:



if 0.8 mol of N_2O_5 (g) is initially put in 2L-reaction vessel and is found to be 0.0125 mol after 2 min, the rate of disappearance of N_2O_5 (g), in M/min, is:


A) 0.9169

B) 0.1969

C) 0.6919

D) 0.9961

18. From the following reaction potential energy (PE) diagram:



which of the following is correct for the forward reaction:

	$\Delta H/\text{kJ}$	Activation energy, E_a/kJ	Type of reaction
A)	+ 20	10	exothermic
B)	+ 20	30	endothermic
C)	- 20	10	exothermic
D)	- 20	40	endothermic

19. In a first order reaction, if the concentration of the reactant changes from 0.1 M to 0.025 M in 40 minutes, the reaction rate, in M/min, when the initial concentration is 0.01 M is:

A) 6.634×10^{-4}

B) 6.346×10^{-4}

C) 4.366×10^{-4}

D) 3.466×10^{-4}

20. For the reaction:



if the value of the rate of disappearance of N_2O_5 is $6.25 \times 10^{-3} \text{ mol L}^{-1} \text{ s}^{-1}$, the rate of appearance of NO_2 is:

A) 2.15×10^{-2}

B) 1.25×10^{-2}

C) 2.51×10^{-2}

D) 2.51×10^{-2}

