




CHEM 101+103



Second Exam Sample 2

Ahmad Aqel Ifseisi

Assistant Professor of Analytical Chemistry
College of Science, Department of Chemistry
King Saud University

P.O. Box 2455 Riyadh 11451 Saudi Arabia

Building: 05, Office: AA53

Tel. 014674198, Fax: 014675992

Web site: <http://fac.ksu.edu.sa/aifseisi>

E-mail: ahmad3qel@yahoo.com

aifseisi@ksu.edu.sa



كرسي أبحاث
المواد المتقدمة
Advanced Materials
Research Chair



1. A piece of 155.0 g aluminum metal at 120°C was placed in a constant pressure calorimeter of negligible heat capacity containing 300.0 g of water at 20°C. Calculate the final temperature of the system (the aluminum metal and the water) in °C; given the specific heat of aluminum metal = 0.90 J/g °C, and that of water = 4.184 J/g °C.

A) 24

B) 26

C) 28

D) 30

2. A 2.2 g sample of quinone $\text{C}_6\text{H}_4\text{O}_2$ was burned in a bomb calorimeter for which the total heat capacity is $7850 \text{ J/}^\circ\text{C}$. The temperature of the calorimeter increased from 23.44°C to 30.57°C . What is the molar heat of combustion of quinone (in kJ/mole)?

- A) -2750** B) -2760 C) -2785 D) -2790

3. Hydrazine, $\text{N}_2\text{H}_4(l)$ is used as a rocket fuel. The thermochemical equation for the combustion of hydrazine is:



How many grams of hydrazine are required to obtain 20654 kJ of heat energy?

- A) 800 B) 900 **C) 1000** D) 1100

4. From the enthalpies of the following reactions:



Calculate (in kJ) ΔH for the following reaction:



A) +2174 B) -2174 C) -2298 D) -2486

5. Given $\Delta H^\circ_f \text{NH}_3(\text{g}) = -46 \text{ kJ/mol}$. Calculate (in kJ) the change in internal energy, ΔE° , for the following reaction:

$$\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightarrow 2\text{NH}_3(\text{g})$$

A) -51 B) -41 **C) -87** D) -97

6. The solubility of nitrogen gas in water at 25°C and its partial pressure at 0.76 atm is 5.2×10^{-4} mol/L. What is the partial pressure of nitrogen (in atm) at which its solubility in water is 1.71×10^{-3} mol/L at 25°C?

A) 2.7

B) 2.5

C) 2.3

D) 2.1

7. Benzene, C_6H_6 , and toluene, C_7H_8 , form ideal solution that has a total vapor pressure of 1.0 atm. What is the mole fraction of benzene in this solution, knowing that the vapor pressure of pure benzene is 1.326 atm and that of pure toluene is 0.532 atm? (all vapor pressures given are measured at exactly the same temperature).

A) 0.56

B) 0.59

C) 0.62

D) 0.64

8. A solution containing 10.0 g of CaCl_2 (an electrolyte) in 100.0 g of water freezes at -4.1°C . What is the vant Hoff factor of this solution? (K_f water = 1.86°C/m).

A) 2.85

B) 2.68

C) 2.59

D) 2.45

9. A solution prepared from 100.0 g of non-volatile non-electrolyte solute in 5.5 mole of toluene (C_7H_8) has a vapor pressure of 0.161 atm at 60 °C. What is the molecular weight (in g/mole) of the solute knowing that the vapor pressure of pure toluene at 60 °C is 0.184 atm?

A) 127.3

B) 132.5

C) 145.6

D) 154.8

10. The molar mass of hemoglobin (a non-electrolyte) is 6.8×10^4 g/mole. What is the osmotic pressure (in mmHg) at 27°C of a solution of 8.0 g hemoglobin in 200 mL of water?

A) 9.0

B) 11.0

C) 13.0

D) 14.0

11. For the hypothetical reaction



the rate of appearance of C given by $\frac{\Delta[C]}{\Delta t}$ may be also expressed as:


A) $+\frac{3\Delta[B]}{2\Delta t}$

B) $-\frac{3\Delta[B]}{2\Delta t}$

C) $-\frac{2\Delta[B]}{3\Delta t}$

D) $+\frac{2\Delta[B]}{3\Delta t}$

12. The data in the table below were obtained for the reaction:



Exp.	[NO] ₀ (M)	[H ₂] ₀ (M)	Initial rate (M/s)
1	0.273	0.763	2.83
2	0.273	1.526	2.83
3	0.819	0.763	25.47

the rate law for this reaction is: rate =

A) $k[\text{NO}]^2 [\text{H}_2]$

B) $k[\text{NO}] [\text{H}_2]^2$

C) $k[\text{NO}]^2 [\text{H}_2]^2$

D) $k[\text{NO}]^2$

13. A compound decomposes by a first-order process. If 25% of the compound decomposes in 61.0 minutes, what is the half life (in minutes) of this decomposition reaction?

A) 142

B) 147

C) 152

D) 160

14. The isomerization of cyclopropane to form propene follow a first-order kinetics. At 700 K, 10% of a sample of cyclopropane is isomerized to propene in 170 minutes. How many minutes are required for 35% of a sample of cyclopropane to change to propene at 700 K?

A) 695

B) 675

C) 625

D) 595

