


Primary Isolation and Cultivation of Viruses

Practical Medical Virology 450 MBIO

2017-18

Diagnostic Virology



Learning Outcome


- The methods used for virus cultivation with the common viruses eligible to grow by those methods.
 - Animal laboratory, Eggs or chicken Embryo, and cell cultures.
- Advantages and disadvantages of each method.
- The types of inoculation sites of chicken embryo and usage.
- The procedure for egg embryo inoculation.

Methods for Cultivation of Virus

Since the viruses are obligate intracellular parasites, they cannot be grown on any culture medium. Viruses can be cultivated within suitable hosts, such as a living cell.



Animals



Eggs Embryo



Cell Culture


Methods for Cultivation of Virus

1- Laboratory Animals

- Used for routine cultivation of virus; they play an essential role in studies of viral pathogenesis.
- Live animals such as monkeys, mice, rabbits, guinea pigs, ferrets are widely used for cultivating viruses.
 - **Monkeys** were used for the isolation of **Poliovirus**. But due to their risk to handlers, monkeys find only limited applications in Virology.
 - **Mice** are the most widely employed animals in virology. There are different routes of inoculation in mice such as; intracerebral, subcutaneous, intraperitoneal or intranasal.
- After the animal is inoculated with the virus suspension, the animal is observed for signs of disease like visible lesions or killed so that infected tissues can be examined for virus.

Methods for Cultivation of Virus

1- Laboratory Animals; e.g. Mice



Methods for Cultivation of Virus

1- Laboratory Animals

Advantages

1. A diagnostic procedure for identifying and isolating a virus from a clinical specimen.
2. Mice provide a reliable model for studying viral replication.
3. Gives unique insight into viral pathogenesis and host virus interactions.
4. Study of immune responses, epidemiology and oncogenesis.


Disadvantages

1. Expensive and difficulties in maintenance of animals.
2. Difficulty in choosing of animals for particular virus.
3. Some human viruses cannot be grown in animals, or can be grown but do not cause disease.
4. Mice do not provide models for vaccine development.
5. It will lead to generation of escape mutants.
6. Issues related to animal welfare systems.

Methods for Cultivation of Virus

2- Chicken Embryo Technique


- Good pasture in 1931 AD first used the embryonated hen's egg.
- The egg used for cultivation must be sterile and the shell should be intact and healthy.
- A hole is drilled in the shell of the embryonated egg, and a viral suspension or suspected virus- containing tissue is injected into the fluid of the egg.
- Viral growth and multiplication in the egg embryo is indicated by the death of the embryo, by embryo cell damage, or by the formation of typical pocks or lesions on the egg membranes



Methods for Cultivation of Virus

2- Chicken Embryo Technique

- An embryonated egg offers various sites for the cultivation of viruses.
 - Chorioallantoic membrane(CAM).
 - Amniotic Cavity.
 - Allantoic Cavity.
 - Yolk sac.
 - Embryo.
 - Air sac.



Copyright © 2010 Pearson Education, Inc.


Methods for Cultivation of Virus

2- Chicken Embryo Technique

A-Chorioallantoic Membrane(CAM):

- Is mainly employed in the growth of **poxvirus**.
- Virus growth and replication in the CAM is indicated by visible lesions (pocks); grey white area in transparent CAM.
- **Herpes simplex virus** is also grown. Each pock is derived from a single virion.

The morphology of the pocks may vary depending on the nature of the virus. Under optimal conditions, each infectious virus particle can form one pock. Hence this method is suitable for plaque studies.




Methods for Cultivation of Virus

2- Chicken Embryo Technique

B- Allantoic Cavity:

- Most popular and simple method.
- Allantoic inoculation is employed for the growth and replication of the **influenza virus** for vaccine production. This will provide a rich yield of influenza and some paramyxoviruses.
- Other allantoic vaccines include **Yellow fever** and **rabies vaccines**.

Duck eggs provide a better yield of rabies virus and were used for the preparation of the inactivated non-neural rabies vaccines. But they need a longer incubation period than embryonated hen's egg. Most of avian viruses can be isolated using this method.




Methods for Cultivation of Virus

2- Chicken Embryo Technique

C- Amniotic Cavity:

The amniotic sac is employed inoculated for primary isolation of **influenza a virus** and the **mumps virus**.

Growth and replication of virus in egg embryo can be detected by haemagglutination assay.




Methods for Cultivation of Virus

2- Chicken Embryo Technique

D- Yolk Sac:

It is also a simplest method for growth and multiplication of virus. This method is also used for the cultivation of some bacteria like **Chlamydiae** and **Rickettsiae**.

Mostly mammalian viruses are isolated using this method. Immune interference mechanism can be detected in most of avian viruses.



Methods for Cultivation of Virus

2- Chicken Embryo:

• Advantages

- Popular method for the isolation of virus and growth
 - (grow virus for some vaccine production)
- Ideal substrate for the viral growth and replication.
- Isolation and cultivation of many avian and few mammalian viruses.
- Cost effective and maintenance is much easier.
- Less labor is needed.
- The embryonated eggs are readily available.
- Sterile and wide range of tissues and fluids
- They are free from contaminating bacteria and many latent viruses.
- Specific and non specific factors of defense are not involved in embryonated eggs.

• Disadvantages




The site of inoculation for varies with different virus. That is, each virus have different sites for their growth and replication.

Methods for Cultivation of Virus

2- Chicken Embryo:

• Procedure:

1. For propagation of influenza virus, pathogen-free eggs are used 11-12 days after fertilization.
2. The egg is placed in front of a light source to locate a non-veined area of the allantoic cavity just below the air sac. This is marked with a pencil.
3. After all the eggs have been 'candled' in this way, a small nick is made in the shell at this position using a jeweler's scribe.



Methods for Cultivation of Virus

2- Chicken Embryo:

• Procedure:

4. Next, a hole is drilled at the top of the egg with a Dremel motorized tool.

If this is not done, when virus is injected, the pressure in the air sac will simply force out the inoculum.



Methods for Cultivation of Virus

2- Chicken Embryo:

• Procedure:

5. After all the eggs have been nicked and drilled, they are inoculated with virus using a tuberculin syringe – a 1 ml syringe fitted with a 1/2 inch, 27 gauge needle.
 - The needle passes through the hole in the shell, through the chorioallantoic membrane, and the virus is placed in the allantoic cavity, which is filled with allantoic fluid.
6. The two holes in the shell are sealed with melted paraffin, and the eggs are placed at 37 degrees C for 48 hours.



Methods for Cultivation of Virus

2- Chicken Embryo:

- Procedure: Videos

1- Viral Cultivation in Chicken Embryo

https://www.youtube.com/watch?v=766QH_qaYN8

2- harvesting virus from seven days old chick embryo

<https://www.youtube.com/watch?v=IS6yWcUSsws>