

470 PHCL

Quiz Sample Questions

Questions (Q)	Answers (A)
<p>Q 1. If the primary adverse effect of a drug is bone marrow suppression, which of the following would be the most appropriate laboratory test for monitoring of toxicity?</p> <p>a. Liver panel</p> <p>b. Complete blood count</p> <p>c. Serum glucose</p> <p>d. BUN and creatinine</p>	<p>A1. B</p>
<p>Q 2. A patient has a BUN of 42 mg/dL and a serum creatinine of 2.1 mg/dL. His estimated creatinine clearance is determined to be 19 mL/min. This information would be important for drugs that</p> <p>a. Are eliminated by the liver</p> <p>b. Are eliminated by the kidney</p> <p>c. Are poorly absorbed from the GI tract</p> <p>d. Have a high first-pass effect</p>	<p>A2. B</p>
<p>Q 3. A patient taking a non-steroidal anti-inflammatory drug (NSAID) complains of frequent abdominal pain and GI upset. This is an example of</p> <p>a. Drug allergy</p> <p>b. Drug sensitivity</p> <p>c. Drug intolerance</p> <p>d. Drug variable</p>	<p>A 3. C</p>
<p>Q 4. Which of the following are appropriate questions concerning the chief complaint and the history of present illness?</p> <p>a. How long have the symptoms been occurring?</p> <p>b. Does anything lessen these symptoms or make them go away?</p> <p>c. Are there any associated signs or symptoms?</p> <p>d. All of the above.</p>	<p>A4. D</p>