

Fundamentals of Organic Chemistry

CHEM 109

For Students of Health Colleges

Credit hrs.: (2+1)

King Saud University

College of Science, Chemistry Department


Structure of Carboxylic Acids

2

- The functional group common to all carboxylic acids is the **carboxyl group**.


*The name is a contraction of the parts: the **carbonyl** and **hydroxyl** groups.*

- The **general formula for a carboxylic acid** can be written in expanded or abbreviated forms.




- Depending on whether an **R** or an **Ar** residue is attached to the carboxyl group; **Carboxylic acids are classified** as;

- Aliphatic Carboxylic Acids.**



- Aromatic Carboxylic Acids.**



- Fatty acids.**

Long straight-chain carboxylic acids with even numbers of carbons, which were first isolated from fats and waxes.

Nomenclature of Carboxylic Acids



Common Names

- The **common names** of carboxylic acids all end in *-ic acid*.
- These names usually come from some Latin or Greek word that indicates the original source of the acid.
- **Common name**, substituents are located with Greek letters, beginning with the α -carbon atom.

IUPAC System

- We replace the final **e** in the name of the corresponding alkane with the suffix *-oic* and add the word *acid*.



- **IUPAC system**, the chain is numbered beginning with the carboxyl carbon atom, and substituents are located in the usual way.

Nomenclature of Carboxylic Acids




Carbon atoms	Formula	Source	Common name	IUPAC name
1	HCOOH	ants (Latin, <i>formica</i>)	formic acid	methanoic acid
2	CH ₃ COOH	vinegar (Latin, <i>acetum</i>)	acetic acid	ethanoic acid
3	CH ₃ CH ₂ COOH	milk (Greek, <i>protos pion</i> , first fat)	propionic acid	propanoic acid
4	CH ₃ (CH ₂) ₂ COOH	butter (Latin, <i>butyrum</i>)	butyric acid	butanoic acid
5	CH ₃ (CH ₂) ₃ COOH	valerian root (Latin, <i>valere</i> , to be strong)	valeric acid	pentanoic acid
6	CH ₃ (CH ₂) ₄ COOH	goats (Latin, <i>caper</i>)	caproic acid	hexanoic acid
7	CH ₃ (CH ₂) ₅ COOH	vine blossom (Greek, <i>oenanthe</i>)	enanthic acid	heptanoic acid
8	CH ₃ (CH ₂) ₆ COOH	goats (Latin, <i>caper</i>)	caprylic acid	octanoic acid
9	CH ₃ (CH ₂) ₇ COOH	pelargonium (an herb with stork-shaped seed capsules; Greek, <i>pelargos</i> , stork)	pelargonic acid	nonanoic acid
10	CH ₃ (CH ₂) ₈ COOH	goats (Latin, <i>caper</i>)	capric acid	decanoic acid


Nomenclature of Carboxylic Acids




6



2-bromopropanoic acid
(α -bromopropionic acid)




propenoic acid
(acrylic acid)




3-hydroxybutanoic acid
(β -hydroxybutyric acid)


- The carboxyl group has priority over alcohol, aldehyde, or ketone functionality in naming.
- The prefix **oxo-** is used to locate the carbonyl group of the aldehyde or ketone.



3-oxopropanoic acid



2-bromo-4-oxopentanoic acid



But-2-enoic acid
(2-Butenoic acid)


Nomenclature of Carboxylic Acids




7

➤ Cycloalkane carboxylic acid


When the carboxyl group is attached to a ring, the ending **-carboxylic acid** is added to the name of the parent **cycloalkane**. (i.e. **Cycloalkanecarboxylic acid**)




Cyclopropanecarboxylic acid



Cyclobutanecarboxylic acid




Cyclopentanecarboxylic acid



Cyclohexanecarboxylic acid


➤ **Aromatic acids** are named by attaching the suffix **-oic acid** or **-ic acid** to an appropriate prefix derived from the aromatic hydrocarbon.




Common name:

Salicylic acid

IUPAC name : 2-Hydroxybenzenecarboxylic acid



benzoic acid
(benzenecarboxylic acid)




p-chlorobenzoic acid
(4-chlorobenzenecarboxylic acid)


Nomenclature of Carboxylic Acids




- **Dicarboxylic acids** (acids that contain two carboxyl groups) are known almost exclusively by their common names.



- **Aliphatic dicarboxylic acids** are given the suffix *-dioic acid* in the IUPAC system.




- The three **benzenedicarboxylic acids** are generally known by their common names.



Physical Properties of Acids

9

- **Carboxylic acids** are **polar** and **they form hydrogen bonds with themselves** or **with other molecules**.
- **Carboxylic acids form dimer**, with the individual units held together by **two hydrogen bonds** between the electron-rich oxygens and the electron-poor hydrogens.



Boiling Points

Therefore, they have high boiling points for their molecular weights-higher even those of comparable alcohols.

Solubility in water

Hydrogen bonding also explains the water solubility of the lower molecular weight carboxylic acids.

- The **first four aliphatic acids** (formic through butyric) are **completely miscible in water**.
- **Aromatic acids are insoluble in water**.

Physical Properties of Acids




Structure	Name	Mol. Wt.	b.p. °C	Solubility in H ₂ O at 25°C
HCOOH	Formic acid	46	100	Very soluble
CH ₃ CH ₂ OH	Ethyl alcohol	46	78	Very soluble
CH ₃ COOH	Acetic acid	60	118	Very soluble
CH ₃ CH ₂ CH ₂ OH	<i>n</i> -Propyl alcohol	60	97	Very soluble
CH ₃ (CH ₂) ₃ COOH	Valeric acid	102	187	4.0 g/100 g H ₂ O
CH ₃ (CH ₂) ₄ CH ₂ OH	<i>n</i> -Hexyl alcohol	102	156	0.6 g/100 g H ₂ O
Ph-COOH	Benzoic acid	122	250	Insoluble
Ph-CH ₂ CH ₂ OH	3-Phenylethanol	122	250	Insoluble

Acid Strength and Structure






- **Carboxylic acids** (RCOOH) dissociate in water, yielding a carboxylate anion (RCOO⁻) and hydronium ion.



Why carboxylic acids are more acidic than alcohols?

- In **ethoxide ion**, the negative charge is localized on a single oxygen atom.
- In **acetate ion**, on the other hand, the negative charge can be delocalized through **resonance**.




Acid Strength and Structure



Effect of Structure on Acidity; the Inductive Effect

- Acidities can vary depending on what other groups are attached to the molecule.
- Recall that *electron-withdrawing groups (-I) enhance acidity*, and *electron-releasing groups (+I) reduce acidity*.

This effect relays charge through bonds, by displacing bonding electrons toward electronegative atoms, or away from electropositive atoms.



Acid Strength and Structure




Effect of Structure on Acidity; the Inductive Effect

- **Formic acid is a substantially stronger acid than acetic acid.**

This suggests that the methyl group is more electron-releasing (hence anion-destabilizing and acidity-reducing) than hydrogen.




- **Example:** acetic acid with those of mono-, di-, and trichloroacetic acids.
Comparison of acid strengths of acetic Acid and chlorinated acetic acids



Acid Strength and Structure




Effect of Structure on Acidity; the Inductive Effect



The more chlorines, the greater the effect and the greater the strength of the acid.

- **Comparison of acid strengths** of butyric acid and the monochlorinated acids.




1) Hydrolysis of Cyanides (Nitriles)


15

- The reaction requires either acid or base.


➤ **In acid**, the nitrogen atom of the cyanide is converted to an ammonium ion..



➤ **In base**, the nitrogen atom is converted to ammonia and the organic product is the carboxylate salt, which must be neutralized in a separate step to give the acid.



- Alkyl cyanides** are generally made from the corresponding alkyl halide.




2) Reaction of Grignard Reagents with Carbon Dioxide (Carbonation of Grignard Reagent)


Preparation of Carboxylic Acids

16

- **Grignard reagents** add to the carbonyl group of carbon dioxide to give acids, after protonation of the intermediate carboxylate salt with a mineral acid like aqueous HCl.
- **The acid obtained has one more carbon atom** (the reaction provides a way to increase the length of a carbon chain).






Specific example





1) Reactions with Bases: Salt Formation

17

- Carboxylic acids, when treated with a strong base, form **carboxylate salts**.





- Examples.**



2) Nucleophilic Substitution Reactions


19



Carboxylic Acid Derivatives



- **Carboxylic acid derivatives** are compounds in which the hydroxyl part of the carboxyl group is replaced by various other groups.



- All acid derivatives can be hydrolyzed to the corresponding carboxylic acid.


Acid derivative	HOH (hydrolysis)
$\begin{array}{c} \text{O} \\ \parallel \\ \text{R}-\text{C}-\text{Cl} \\ \text{acyl halide} \end{array}$	$\begin{array}{c} \text{O} \\ \parallel \\ \text{R}-\text{C}-\text{OH} + \text{HCl} \end{array}$
$\begin{array}{c} \text{O} \quad \text{O} \\ \parallel \quad \parallel \\ \text{R}-\text{C}-\text{O}-\text{C}-\text{R} \\ \text{acid anhydride} \end{array}$	$2 \begin{array}{c} \text{O} \\ \parallel \\ \text{R}-\text{C}-\text{OH} \end{array}$
$\begin{array}{c} \text{O} \\ \parallel \\ \text{R}-\text{C}-\text{O}-\text{R}' \\ \text{ester} \end{array}$	$\begin{array}{c} \text{O} \\ \parallel \\ \text{R}-\text{C}-\text{OH} + \text{R}'\text{OH} \end{array}$
$\begin{array}{c} \text{O} \\ \parallel \\ \text{R}-\text{C}-\text{NH}_2 \\ \text{amide} \end{array}$	$\begin{array}{c} \text{O} \\ \parallel \\ \text{R}-\text{C}-\text{OH} + \text{NH}_3 \end{array}$
<i>Main organic product</i>	<i>acid</i>

Acid Chloride

191



- **Acyl chlorides** have the general formula RCOCl.
- **Acyl chlorides** are more common and less expensive than bromides or iodides.
- **Nomenclature:**

Acyl chlorides, or acid chlorides, are named by replacing the -ic acid ending of the parent acid by -yl chloride.



- **Preparation:**


They can be prepared from acids by reaction with thionyl chloride or phosphorous pentachloride.






Acid Chloride

201

- **Reactions:** They can react rapidly with most nucleophile.








- **Examples:**



Esters


211

- **Esters** are derived from acids by replacing the $-\text{OH}$ group by an $-\text{OR}$ group and have the general formula $\text{R}'\text{COOR}$.
- **Nomenclature:**
 - They are named in a manner analogous to carboxylic acid salts.
 - The **R part of the $-\text{OR}$ group is name first**, followed by the name of the acid, with the **$-\text{ic acid}$** ending changed to **$-\text{ate}$** .




- **Preparation:**

When a carboxylic acid and an alcohol are heated in the presence of an acid catalyst (HCl or H_2SO_4), an equilibrium is established with the ester and water.




○ Reactions


- **Cyclic esters (lactones)** can be prepared from hydroxy acids if these groups can come in contact through bending of the chain.



- **Saponification;** esters are commonly hydrolyzed with base.




- Ammonia converts esters to **amides**.




- **Amides** are the least reactive of the common carboxylic acid derivatives.
- Primary amides have general formula **RCNH₂**.

➤ **Nomenclature:**


Amides are named by replacing the *-ic* or *-oic acid* ending of the acid name, either the common or the IUPAC name, with the *-amide* ending.




formamide
(methanamide)




acetamide
(ethanamide)




butanamide




benzamide
(benzenecarboxamide)




formamide



N,N-dimethylformamide




acetamide



N,N-dimethylacetamide


○ Preparation:

- They can be prepared by the reaction of ammonia with esters, with acyl halides, or with acid anhydrides.
- Amides can also prepared by heating the ammonium salts of acids.




○ Reactions

- **Amides** react with nucleophiles and they can be hydrolyzed by water.



- **Amides** can be reduced by lithium aluminum hydride to give amines.





Acid Anhydrides

31


➤ **Acid anhydrides** have general formula **RCOOCOR**.

➤ **Nomenclature:**


The name of an anhydrides is obtained by naming the acid from which is derived and replacing the word acid with anhydride.



IUPAC name: Propanoic anhydride
Common name: Propionic anhydride



Benzoic anhydride




Succinic anhydride

Acid Anhydrides


32

➤ Preparation

- **Acid anhydrides** are derived from acids by removing water from two carboxyl groups and connecting the fragments.



- **Anhydrides** can also be prepared from acid chlorides and carboxylate salts. *This method is used for preparing anhydrides derived from two different carboxylic acids (mixed anhydrides).*



Acid Anhydrides

33

➤ Reactions

- **Anhydrides** undergo **nucleophilic acyl substitution reactions** (They are more reactive than esters, but less reactive than acyl halides).

