


Titration curve of amino acids

Titration Curves :

- Titration Curves are produced by monitoring the pH of a **given volume** of a sample solution after successive **addition of acid or alkali**.
- The curves are usually plots of pH against the volume of titrant added (acid or base).
- Each **dissociation group** represent **one stage** in the titration curve.

Amino acid general formula:

- Amino acids consist of:
 - A basic amino group (—NH_2)
 - An acidic carboxyl group (—COOH)
 - A hydrogen atom (—H)
 - A distinctive side chain (—R).



Titration of amino acid:


- When an amino acid is dissolved in water it exists predominantly in the **isoelectric form**.
- Amino acid is an **amphoteric** compound → It act as either an acid or a base:
 - **Upon titration with acid** → it acts as a BASE (accept a proton).
 - **Upon titration with base** → it acts as an ACID (donate a proton)

Titration of amino acid cont':

- Amino acids are example of **weak acid** which contain **more than one dissociate group**.
- **Examples:**

(1) Alanine:


- Contain COOH ($\text{pK}_{\text{a}1}= 2.34$) and NH_3^+ ($\text{pK}_{\text{a}2}= 9.69$) groups (it has one pI value =6.010). [Diprotic]
- The COOH will dissociate first then NH_3^+ dissociate later . (Because $\text{pK}_{\text{a}1}<\text{pK}_{\text{a}2}$)



Full protonated alanine


(2) Arginine:

- Contain COOH ($\text{pK}_{\text{a}1}= 2.34$) , NH_3^+ ($\text{pK}_{\text{a}2}= 9.69$) groups and basic group ($\text{pK}_{\text{a}3}=12.5$) (it has one pI value=11). [Triprotic]



Arginine

Titration curve of Alanine



pK_1 carboxylic acid = 2.34

pK_2 amino group = 9.69

$\text{pI} = (\text{pK}_1 + \text{pK}_2)/2$

Titration curve of alanine or glycine [diprotic]:

[1] In starting point:


- Alanine is **full protonated**.
- $[\text{NH}_3^+-\text{CH}-\text{CH}_3-\text{COOH}]$.

[2] COOH will dissociate **first**:

- $[\text{NH}_3^+-\text{CH}-\text{CH}_3-\text{COOH}] > [\text{NH}_3^+-\text{CH}-\text{CH}_3-\text{COO}^-]$
- $\text{pH} < \text{pK}_{\text{a}1}$.

[3] In this point the component of alanine act as **buffer**:

- $[\text{NH}_3^+-\text{CH}-\text{CH}_3-\text{COOH}] = [\text{NH}_3^+-\text{CH}-\text{CH}_3-\text{COO}^-]$.
- $\text{pH} = \text{pK}_{\text{a}1}$



Titration curve of alanine or glycine [diprotic]:

[4] In this point:


- $[\text{NH}_3^+-\text{CH}-\text{CH}_3-\text{COOH}] < [\text{NH}_3^+-\text{CH}-\text{CH}_3-\text{COO}^-]$.
- $\text{pH} > \text{pKa}_1$.

[5] Isoelectric point:

- The COOH is **full dissociate** to COO^- .
- $[\text{NH}_3^+-\text{CH}-\text{CH}_3-\text{COO}^-]$.
- Con. of -ve charge = Con. of +ve charge.
- The amino acid present as **Zwitter ion** (neutral form) .
- Remember that :PI (isoelectric point) is the pH value at which the net charge of amino acid equal to zero.
- $\text{pI} = (\text{pKa}_1 + \text{pKa}_2) / 2 = (2.32 + 9.96) / 2 = 6.01$

[6] The NH_3^+ **start dissociate**:

- $[\text{NH}_3^+-\text{CH}-\text{CH}_3-\text{COO}^-] > [\text{NH}_2-\text{CH}-\text{CH}_3-\text{COO}^-]$.
- $\text{pH} < \text{pKa}_2$.



Titration curve of alanine or glycine [diprotic]:

[7] In this point the component of alanine act as **buffer**:


- $[\text{NH}_3^+-\text{CH}-\text{CH}_3-\text{COO}^-] = [\text{NH}_2-\text{CH}-\text{CH}_3-\text{COO}^-]$.
- $\text{pH} = \text{pK}_a_2$.

[8] In this point:

- $[\text{NH}_3^+-\text{CH}-\text{CH}_3-\text{COO}^-] < [\text{NH}_2-\text{CH}-\text{CH}_3-\text{COO}^-]$.
- $\text{pH} > \text{pK}_a_2$

[9] End point:

- The alanine is **full dissociated**.
 - $[\text{NH}_2-\text{CH}-\text{CH}_3-\text{COO}^-]$
 - $\text{pOH} = (\text{pK}_b + \text{P}[\text{A}^-])/2$
- $\text{pK}_b = \text{pK}_w - \text{pK}_a_2$



Calculating the pH at different point of the titration curve :

The pH calculated by different way :

[1] at starting point :

$$\text{pH} = (\text{pK}_a + \text{P}[\text{HA}]) / 2$$

[2] At any point within the curve (befor or in or after middle titration):

$$\text{pH} = \text{pK}_a + \log([A^-] / [\text{HA}])$$

[3] At end point:

$$\text{pOH} = (\text{pK}_b + \text{P}[A^-]) / 2$$

$$\text{pH} = \text{pK}_w - \text{pOH}$$

$$\text{pK}_b = \text{pK}_w - \text{pK}_a$$

Practical Part

Objectives:

- To study titration curves of amino acid.
- To use this curve to estimate the pK_a values of the ionizable groups of the amino acid.
- To determine pI .
- To determine the buffering region.
- To understand the acid base behaviour of an amino acid.

Method:

- a) You are provided with 10 ml of a 0.1M alanine solution, titrate it with 0.1M NaOH adding the base drop wise mixing, and recording the pH after each 0.5 ml NaOH added until you reach a pH=11.

Measured pH value	Amount of 0.1M NaOH added [ml]

- b) Take another 10 ml of a 0.1M alanine solution, titrate it with 0.1 M HCL adding the acid drop wise mixing, and recording the pH after each 0.5 ml HCL added until you reach a pH=2.17.

Measured pH value	Amount of 0.1M HCl added [ml]

Results:

- Record the titration table and plot a curve of pH versus ml of titrant added.
- Calculate the pH of the alanine solution after the addition of 0 ml, 5ml, of 0.2M NaOH, and calculate pH after addition of 0.5 ml , 2ml of HCl.
- Determine the pKa of ionizable groups of amino acids.
- Compare your calculated pH values with those obtained from Curve.
- Determine the PI value from your result .

