

Acta Crystallographica Section E

Structure Reports

Online


ISSN 1600-5368

(*E*)-2-(1,1-Dicyclohexyl-3-phenylallyl)-5,5-dimethyl-1,3,2-dioxaborinane

Gamal A. El-Hiti, Keith Smith, Mark C. Elliott, Dyfyr Heulyn Jones and Benson M. Kariuki

Acta Cryst. (2013). **E69**, o1403

This open-access article is distributed under the terms of the Creative Commons Attribution Licence <http://creativecommons.org/licenses/by/2.0/uk/legalcode>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original authors and source are cited.



Acta Crystallographica Section E: Structure Reports Online is the IUCr's highly popular open-access structural journal. It provides a simple and easily accessible publication mechanism for the growing number of inorganic, metal-organic and organic crystal structure determinations. The electronic submission, validation, refereeing and publication facilities of the journal ensure very rapid and high-quality publication, whilst key indicators and validation reports provide measures of structural reliability. The journal publishes over 4000 structures per year. The average publication time is less than one month.

Crystallography Journals **Online** is available from journals.iucr.org

(E)-2-(1,1-Dicyclohexyl-3-phenylallyl)-5,5-dimethyl-1,3,2-dioxaborinaneGamal A. El-Hiti,^{a*} Keith Smith,^b Mark C. Elliott,^b
Dyfyrr Heulyn Jones^b and Benson M. Kariuki^{b*}^aDepartment of Optometry, College of Applied Medical Sciences, King Saud University, PO Box 10219, Riyadh 11433, Saudi Arabia, and ^bSchool of Chemistry, Cardiff University, Main Building, Park Place, Cardiff CF10 3AT, Wales
Correspondence e-mail: gelhiti@ksu.edu.sa, kariukib@cardiff.ac.uk


Received 29 July 2013; accepted 3 August 2013

Key indicators: single-crystal X-ray study; $T = 150$ K; mean $\sigma(\text{C}—\text{C}) = 0.002$ Å;
 R factor = 0.050; wR factor = 0.128; data-to-parameter ratio = 20.0.

The crystal structure of the title compound, $\text{C}_{26}\text{H}_{39}\text{BO}_2$, which contains no strong hydrogen bond donors, displays only long $\text{C}—\text{H} \cdots \text{O}$ contacts between inversion-related pairs of molecules. The structure contains layers rich in oxygen and boron parallel to the ac plane. The dioxaborinane ring adopts an envelope conformation with the C atom attached to the two methyl groups as the flap.

Related literature

For the synthesis and applications of allylboronic esters, see: Lombardo *et al.* (2002); Carosi & Hall (2007); Althaus *et al.* (2010); Fandrick *et al.* (2010); Clary *et al.* (2011); Hesse *et al.* (2012); Incerti-Pradillos *et al.* (2013). For the X-ray structure of a boronic ester, see: Sopková-de Oliveira Santos *et al.* (2003).

**Experimental***Crystal data* $\text{C}_{26}\text{H}_{39}\text{BO}_2$
 $M_r = 394.38$
Triclinic, $P\bar{1}$
 $a = 9.4967$ (3) Å
 $b = 11.2837$ (2) Å
 $c = 12.0297$ (4) Å
 $\alpha = 109.897$ (2)°
 $\beta = 96.388$ (2)° $\gamma = 102.048$ (2)°
 $V = 1161.90$ (6) Å³
 $Z = 2$
Mo $K\alpha$ radiation
 $\mu = 0.07$ mm⁻¹
 $T = 150$ K
 $0.38 \times 0.30 \times 0.28$ mm*Data collection*Nonius KappaCCD diffractometer
Absorption correction: multi-scan
(*DENZO/SCALEPACK*;
Otwinowski & Minor, 1997)
 $T_{\min} = 0.975$, $T_{\max} = 0.981$ 8277 measured reflections
5287 independent reflections
4303 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.027$ *Refinement* $R[F^2 > 2\sigma(F^2)] = 0.050$
 $wR(F^2) = 0.128$
 $S = 1.03$
5287 reflections264 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.32$ e Å⁻³
 $\Delta\rho_{\min} = -0.20$ e Å⁻³**Table 1**

Hydrogen-bond geometry (Å, °).

$D—H \cdots A$	$D—H$	$H \cdots A$	$D \cdots A$	$D—H \cdots A$
$\text{C22}—\text{H22B} \cdots \text{O2}^i$	0.99	2.69	3.5773 (18)	150

Symmetry code: (i) $-x + 1, -y, -z$.

Data collection: *COLLECT* (Nonius, 2000); cell refinement: *SCALEPACK* (Otwinowski & Minor, 1997); data reduction: *DENZO* (Otwinowski & Minor, 1997) and *SCALEPACK*; program(s) used to solve structure: *SIR92* (Altomare *et al.*, 1994); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP99* for Windows (Farrugia, 2012); software used to prepare material for publication: *WinGX* (Farrugia, 2012) and *CHEMDRAW Ultra* (Cambridge Soft, 2001).

The authors would like to extend their appreciation to the Deanship of Scientific Research at King Saud University for its funding for this research through the research group project RGP-VPP-239.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GO2094).

References

- Althaus, M., Mahmood, A., Suárez, J. R., Thomas, S. P. & Aggarwal, V. K. (2010). *J. Am. Chem. Soc.* **132**, 4025–4028.
- Altomare, A., Casciarano, G., Giacovazzo, C., Guagliardi, A., Burla, M. C., Polidori, G. & Camalli, M. (1994). *J. Appl. Cryst.* **27**, 435.
- Cambridge Soft (2001). *CHEMDRAW Ultra*. Cambridge Soft Corporation, Cambridge, Massachusetts, USA.
- Carosi, L. & Hall, D. G. (2007). *Angew. Chem. Int. Ed.* **46**, 5913–5915.
- Clary, J. W., Rettenmaier, T. J., Snelling, R., Bryks, W., Banwell, J., Wipke, W. T. & Singaram, B. (2011). *J. Org. Chem.* **76**, 9602–9610.
- Fandrick, K. R., Fandrick, D. R., Gao, J. J., Reeves, J. T., Tan, Z., Li, W., Song, J. J., Lu, B., Yee, N. K. & Senanayake, C. H. (2010). *Org. Lett.* **12**, 3748–3751.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Hesse, M. J., Butts, C. P., Willis, C. L. & Aggarwal, V. K. (2012). *Angew. Chem. Int. Ed.* **51**, 12444–12448.
- Incerti-Pradillos, C. A., Kabeshov, M. A. & Malkov, A. V. (2013). *Angew. Chem. Int. Ed.* **52**, 5338–5341.
- Lombardo, M., Morganti, S., Tozzi, M. & Trombini, C. (2002). *Eur. J. Org. Chem.* pp. 2823–2830.
- Nonius (2000). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sopková-de Oliveira Santos, J., Lancelot, J.-C., Bouillon, A. & Rault, S. (2003). *Acta Cryst.* **C59**, o111–o113.

supplementary materials

Acta Cryst. (2013). E69, o1403 [doi:10.1107/S1600536813021739]

(*E*)-2-(1,1-Dicyclohexyl-3-phenylallyl)-5,5-dimethyl-1,3,2-dioxaborinane

Gamal A. El-Hiti, Keith Smith, Mark C. Elliott, Dyfyr Heulyn Jones and Benson M. Kariuki

1. Comment

The title compound I, a useful synthetic intermediate, was synthesized by the reaction of (*E*)-dicyclohexylstyrylborane with the anion of dichloromethyl methyl ether followed by esterification with 2,2-dimethyl-1,3-propanediol. Allylboronic esters have been synthesized from the reaction of lithiated carbamates with vinylboranes [Althaus *et al.* (2010)], and from the reaction of primary allyl halides with pinacolborane and magnesium [Incerti-Pradillos *et al.* (2013), Clary *et al.* (2011)]. Allylboronic esters are important synthetic intermediates, which have been shown to react with aldehydes to give homoallylic alcohols [Lombardo *et al.* (2002)], with the control of the newly-generated stereogenic centre possible through use of a chiral catalyst [Carosi *et al.* (2007)]. Allylboronic esters take part in a zinc alkoxide catalysed reaction with ketones to give the corresponding homoallylic alcohol products [Fandrick *et al.* (2010)], and also take part in a proto-deboronation reaction which has been used to synthesize the pheromone of the Californian red scale beetle [Hesse *et al.* (2012)]. For the X-ray structure of a boronic ester, see: Sopková-de Oliveira Santos *et al.* (2003).

In the molecule (Figure 1), the two cyclohexyl groups assume a chair conformation and an envelope conformation is observed for the dioxaborinane ring. The phenylallyl group is not planar as the plane through the double bond makes an angle of 20.84 ° with the phenyl group. There are no strong hydrogen bond donors in the structure. Long contacts of C—H···O type occur between pairs of molecules to form loosely bound dimers (Figure 2). The dimers are stacked along the *a*-axis to form a structure with layers rich in oxygen and boron parallel to the *ac* plane (Figure 3).

2. Refinement

H atoms were positioned geometrically and refined using a riding model with $U_{\text{iso}}(\text{H}) = 1.2$ times U_{eq} for the atom they are bonded to except for the methyl groups where 1.5 times U_{eq} was used with free rotation about the C—C bond. Of the low angle reflections not included in the refinement only (0 0 1) and (0 1 0) were omitted due to low intensities consistent with being obscured by the beamstop. The rest were eliminated automatically during data processing possibly as overloads.

Computing details

Data collection: *COLLECT* (Nonius, 2000); cell refinement: *SCALEPACK* (Otwinowski & Minor, 1997); data reduction: *DENZO* and *SCALEPACK* (Otwinowski & Minor, 1997); program(s) used to solve structure: *SIR92* (Altomare *et al.*, 1994); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP99* for Windows (Farrugia, 2012); software used to prepare material for publication: *WinGX* (Farrugia, 2012) and *CHEMDRAW Ultra* (Cambridge Soft, 2001).




Figure 1

A molecule showing atom labels and 50% probability displacement ellipsoids for non-H atoms.




Figure 2

A pair of molecules showing C—H...O interactions as dotted lines.




Figure 3

Molecular packing in the crystal structure showing oxygen and boron rich layers.

(E)-2-(1,1-Dicyclohexyl-3-phenylallyl)-5,5-dimethyl-1,3,2-dioxaborinane

Crystal data

$C_{26}H_{39}BO_2$

$M_r = 394.38$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 9.4967$ (3) Å

$b = 11.2837$ (2) Å

$c = 12.0297$ (4) Å

$\alpha = 109.897$ (2)°

$\beta = 96.388$ (2)°

$\gamma = 102.048$ (2)°

$V = 1161.90$ (6) Å³

$Z = 2$

$F(000) = 432$

$D_x = 1.127$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 4303 reflections

$\theta = 2.2$ – 27.5°

$\mu = 0.07$ mm⁻¹

$T = 150$ K

Block, colourless

$0.38 \times 0.30 \times 0.28$ mm

Data collection

Nonius KappaCCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

CCD slices, ω and ϕ scans

Absorption correction: multi-scan

(*DENZO/SCALEPACK*; Otwinowski & Minor, 1997)

$T_{\min} = 0.975$, $T_{\max} = 0.981$

8277 measured reflections

5287 independent reflections

4303 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.027$

$\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 2.1^\circ$

$h = -12 \rightarrow 12$

$k = -14 \rightarrow 14$

$l = -15 \rightarrow 12$

Refinement

Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.050$
 $wR(F^2) = 0.128$
 $S = 1.03$

5287 reflections

264 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0488P)^2 + 0.4651P]$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.004$
 $\Delta\rho_{\max} = 0.32 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.20 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.24096 (15)	0.23892 (13)	0.62817 (11)	0.0235 (3)
C2	0.22913 (18)	0.35166 (15)	0.71836 (13)	0.0348 (3)
H2	0.2828	0.4349	0.7232	0.042*
C3	0.14007 (19)	0.34351 (17)	0.80085 (14)	0.0411 (4)
H3	0.1319	0.4211	0.8607	0.049*
C4	0.06330 (17)	0.22337 (17)	0.79648 (14)	0.0382 (4)
H4	0.0029	0.2179	0.8534	0.046*
C5	0.07497 (17)	0.11101 (16)	0.70869 (14)	0.0348 (3)
H5	0.0228	0.0280	0.7055	0.042*
C6	0.16250 (16)	0.11876 (13)	0.62506 (13)	0.0281 (3)
H6	0.1689	0.0407	0.5648	0.034*
C7	0.33581 (15)	0.25145 (13)	0.54118 (12)	0.0255 (3)
H7	0.4098	0.3312	0.5644	0.031*
C8	0.32693 (14)	0.16134 (12)	0.43362 (11)	0.0221 (3)
H8	0.2530	0.0818	0.4117	0.027*
C9	0.42193 (14)	0.17080 (12)	0.34118 (11)	0.0205 (3)
C10	0.58492 (14)	0.24810 (12)	0.40612 (11)	0.0218 (3)
H10	0.5813	0.3293	0.4725	0.026*
C11	0.65750 (15)	0.16852 (13)	0.46484 (12)	0.0260 (3)
H11A	0.5978	0.1466	0.5210	0.031*
H11B	0.6589	0.0856	0.4014	0.031*
C12	0.81407 (16)	0.24166 (14)	0.53369 (13)	0.0330 (3)
H12A	0.8121	0.3195	0.6031	0.040*
H12B	0.8576	0.1844	0.5655	0.040*
C13	0.90896 (17)	0.28434 (15)	0.45299 (15)	0.0380 (4)
H13A	0.9200	0.2064	0.3883	0.046*

H13B	1.0079	0.3364	0.5010	0.046*
C14	0.83863 (17)	0.36592 (15)	0.39717 (15)	0.0361 (4)
H14A	0.8996	0.3908	0.3430	0.043*
H14B	0.8349	0.4471	0.4618	0.043*
C15	0.68323 (15)	0.29066 (13)	0.32573 (13)	0.0283 (3)
H15A	0.6875	0.2125	0.2577	0.034*
H15B	0.6400	0.3466	0.2918	0.034*
C16	0.35441 (15)	0.23557 (12)	0.25946 (12)	0.0229 (3)
H16	0.4152	0.2331	0.1963	0.027*
C17	0.19670 (16)	0.16006 (14)	0.19300 (13)	0.0309 (3)
H17A	0.1928	0.0671	0.1500	0.037*
H17B	0.1315	0.1645	0.2522	0.037*
C18	0.14178 (19)	0.21631 (17)	0.10238 (15)	0.0404 (4)
H18A	0.2011	0.2042	0.0387	0.048*
H18B	0.0384	0.1683	0.0635	0.048*
C19	0.1523 (2)	0.36161 (18)	0.16395 (16)	0.0436 (4)
H19A	0.1255	0.3970	0.1021	0.052*
H19B	0.0814	0.3726	0.2187	0.052*
C20	0.30661 (19)	0.43826 (15)	0.23583 (14)	0.0364 (4)
H20A	0.3073	0.5301	0.2804	0.044*
H20B	0.3754	0.4379	0.1797	0.044*
C21	0.35828 (17)	0.37906 (13)	0.32525 (13)	0.0287 (3)
H21A	0.4597	0.4290	0.3689	0.034*
H21B	0.2940	0.3855	0.3852	0.034*
C22	0.51831 (16)	−0.10974 (13)	0.09454 (13)	0.0290 (3)
H22A	0.6131	−0.1146	0.1338	0.035*
H22B	0.5259	−0.1158	0.0115	0.035*
C23	0.39668 (15)	−0.22485 (12)	0.08888 (12)	0.0256 (3)
C24	0.37886 (18)	−0.20513 (13)	0.21769 (12)	0.0301 (3)
H24A	0.2915	−0.2720	0.2158	0.036*
H24B	0.4659	−0.2183	0.2607	0.036*
C25	0.25412 (17)	−0.23168 (15)	0.01216 (14)	0.0363 (4)
H25A	0.1768	−0.3060	0.0092	0.055*
H25B	0.2688	−0.2427	−0.0697	0.055*
H25C	0.2250	−0.1507	0.0477	0.055*
C26	0.44291 (19)	−0.35104 (14)	0.03502 (14)	0.0373 (4)
H26A	0.5375	−0.3438	0.0822	0.056*
H26B	0.4524	−0.3656	−0.0486	0.056*
H26C	0.3685	−0.4246	0.0369	0.056*
B1	0.42494 (16)	0.02685 (14)	0.25703 (13)	0.0210 (3)
O1	0.36243 (11)	−0.07729 (8)	0.28350 (8)	0.0261 (2)
O2	0.49222 (11)	0.01427 (9)	0.16035 (8)	0.0273 (2)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0250 (7)	0.0294 (6)	0.0191 (6)	0.0096 (5)	0.0060 (5)	0.0110 (5)
C2	0.0404 (9)	0.0318 (7)	0.0276 (7)	0.0065 (6)	0.0118 (6)	0.0056 (6)
C3	0.0444 (9)	0.0484 (9)	0.0257 (8)	0.0151 (8)	0.0147 (7)	0.0040 (7)
C4	0.0306 (8)	0.0644 (10)	0.0285 (8)	0.0168 (7)	0.0142 (6)	0.0232 (7)

C5	0.0303 (8)	0.0443 (8)	0.0394 (8)	0.0094 (6)	0.0119 (6)	0.0263 (7)
C6	0.0299 (7)	0.0298 (7)	0.0296 (7)	0.0115 (6)	0.0099 (6)	0.0138 (6)
C7	0.0291 (7)	0.0248 (6)	0.0240 (7)	0.0063 (5)	0.0094 (5)	0.0102 (5)
C8	0.0238 (7)	0.0240 (6)	0.0219 (6)	0.0079 (5)	0.0073 (5)	0.0108 (5)
C9	0.0243 (6)	0.0216 (6)	0.0188 (6)	0.0083 (5)	0.0072 (5)	0.0092 (5)
C10	0.0229 (6)	0.0225 (6)	0.0210 (6)	0.0072 (5)	0.0066 (5)	0.0081 (5)
C11	0.0286 (7)	0.0282 (6)	0.0230 (7)	0.0092 (5)	0.0049 (5)	0.0106 (5)
C12	0.0309 (8)	0.0338 (7)	0.0318 (8)	0.0121 (6)	0.0013 (6)	0.0085 (6)
C13	0.0264 (8)	0.0372 (8)	0.0450 (9)	0.0078 (6)	0.0061 (7)	0.0093 (7)
C14	0.0276 (8)	0.0352 (8)	0.0442 (9)	0.0035 (6)	0.0120 (7)	0.0145 (7)
C15	0.0285 (7)	0.0301 (7)	0.0295 (7)	0.0067 (6)	0.0111 (6)	0.0140 (6)
C16	0.0265 (7)	0.0262 (6)	0.0219 (6)	0.0114 (5)	0.0088 (5)	0.0123 (5)
C17	0.0295 (8)	0.0356 (7)	0.0295 (7)	0.0114 (6)	0.0042 (6)	0.0136 (6)
C18	0.0355 (9)	0.0569 (10)	0.0353 (8)	0.0190 (8)	0.0016 (7)	0.0228 (7)
C19	0.0482 (10)	0.0627 (11)	0.0443 (9)	0.0370 (9)	0.0182 (8)	0.0340 (8)
C20	0.0511 (10)	0.0379 (8)	0.0391 (8)	0.0276 (7)	0.0211 (7)	0.0244 (7)
C21	0.0378 (8)	0.0276 (7)	0.0282 (7)	0.0158 (6)	0.0107 (6)	0.0142 (5)
C22	0.0338 (8)	0.0268 (7)	0.0280 (7)	0.0129 (6)	0.0142 (6)	0.0067 (5)
C23	0.0303 (7)	0.0246 (6)	0.0224 (7)	0.0103 (5)	0.0080 (5)	0.0068 (5)
C24	0.0475 (9)	0.0219 (6)	0.0245 (7)	0.0128 (6)	0.0112 (6)	0.0096 (5)
C25	0.0369 (9)	0.0403 (8)	0.0265 (7)	0.0126 (7)	0.0036 (6)	0.0054 (6)
C26	0.0498 (10)	0.0284 (7)	0.0339 (8)	0.0173 (7)	0.0133 (7)	0.0064 (6)
B1	0.0235 (7)	0.0238 (7)	0.0179 (7)	0.0080 (5)	0.0047 (5)	0.0094 (5)
O1	0.0400 (6)	0.0207 (4)	0.0199 (5)	0.0090 (4)	0.0112 (4)	0.0082 (4)
O2	0.0363 (6)	0.0229 (4)	0.0255 (5)	0.0100 (4)	0.0153 (4)	0.0083 (4)

Geometric parameters (Å, °)

C1—C6	1.3902 (19)	C16—C17	1.531 (2)
C1—C2	1.3983 (18)	C16—C21	1.5316 (17)
C1—C7	1.4777 (17)	C16—H16	1.0000
C2—C3	1.388 (2)	C17—C18	1.532 (2)
C2—H2	0.9500	C17—H17A	0.9900
C3—C4	1.379 (2)	C17—H17B	0.9900
C3—H3	0.9500	C18—C19	1.528 (2)
C4—C5	1.381 (2)	C18—H18A	0.9900
C4—H4	0.9500	C18—H18B	0.9900
C5—C6	1.3879 (19)	C19—C20	1.524 (3)
C5—H5	0.9500	C19—H19A	0.9900
C6—H6	0.9500	C19—H19B	0.9900
C7—C8	1.3264 (18)	C20—C21	1.5338 (19)
C7—H7	0.9500	C20—H20A	0.9900
C8—C9	1.5251 (16)	C20—H20B	0.9900
C8—H8	0.9500	C21—H21A	0.9900
C9—C16	1.5672 (17)	C21—H21B	0.9900
C9—C10	1.5717 (18)	C22—O2	1.4424 (15)
C9—B1	1.6034 (18)	C22—C23	1.5238 (19)
C10—C11	1.5369 (18)	C22—H22A	0.9900
C10—C15	1.5386 (17)	C22—H22B	0.9900
C10—H10	1.0000	C23—C24	1.5227 (18)

C11—C12	1.525 (2)	C23—C25	1.523 (2)
C11—H11A	0.9900	C23—C26	1.5285 (18)
C11—H11B	0.9900	C24—O1	1.4408 (15)
C12—C13	1.523 (2)	C24—H24A	0.9900
C12—H12A	0.9900	C24—H24B	0.9900
C12—H12B	0.9900	C25—H25A	0.9800
C13—C14	1.524 (2)	C25—H25B	0.9800
C13—H13A	0.9900	C25—H25C	0.9800
C13—H13B	0.9900	C26—H26A	0.9800
C14—C15	1.528 (2)	C26—H26B	0.9800
C14—H14A	0.9900	C26—H26C	0.9800
C14—H14B	0.9900	B1—O1	1.3565 (17)
C15—H15A	0.9900	B1—O2	1.3682 (16)
C15—H15B	0.9900		
C6—C1—C2	117.89 (12)	C17—C16—H16	106.8
C6—C1—C7	122.74 (12)	C21—C16—H16	106.8
C2—C1—C7	119.37 (12)	C9—C16—H16	106.8
C3—C2—C1	120.89 (14)	C18—C17—C16	111.14 (12)
C3—C2—H2	119.6	C18—C17—H17A	109.4
C1—C2—H2	119.6	C16—C17—H17A	109.4
C4—C3—C2	120.37 (14)	C18—C17—H17B	109.4
C4—C3—H3	119.8	C16—C17—H17B	109.4
C2—C3—H3	119.8	H17A—C17—H17B	108.0
C3—C4—C5	119.46 (13)	C19—C18—C17	111.29 (13)
C3—C4—H4	120.3	C19—C18—H18A	109.4
C5—C4—H4	120.3	C17—C18—H18A	109.4
C4—C5—C6	120.36 (14)	C19—C18—H18B	109.4
C4—C5—H5	119.8	C17—C18—H18B	109.4
C6—C5—H5	119.8	H18A—C18—H18B	108.0
C5—C6—C1	121.03 (13)	C20—C19—C18	111.51 (12)
C5—C6—H6	119.5	C20—C19—H19A	109.3
C1—C6—H6	119.5	C18—C19—H19A	109.3
C8—C7—C1	125.85 (12)	C20—C19—H19B	109.3
C8—C7—H7	117.1	C18—C19—H19B	109.3
C1—C7—H7	117.1	H19A—C19—H19B	108.0
C7—C8—C9	127.42 (12)	C19—C20—C21	111.18 (13)
C7—C8—H8	116.3	C19—C20—H20A	109.4
C9—C8—H8	116.3	C21—C20—H20A	109.4
C8—C9—C16	109.58 (10)	C19—C20—H20B	109.4
C8—C9—C10	110.43 (10)	C21—C20—H20B	109.4
C16—C9—C10	111.93 (10)	H20A—C20—H20B	108.0
C8—C9—B1	109.36 (10)	C16—C21—C20	110.77 (12)
C16—C9—B1	108.39 (10)	C16—C21—H21A	109.5
C10—C9—B1	107.06 (10)	C20—C21—H21A	109.5
C11—C10—C15	109.13 (11)	C16—C21—H21B	109.5
C11—C10—C9	110.54 (10)	C20—C21—H21B	109.5
C15—C10—C9	115.16 (11)	H21A—C21—H21B	108.1
C11—C10—H10	107.2	O2—C22—C23	112.27 (10)

C15—C10—H10	107.2	O2—C22—H22A	109.2
C9—C10—H10	107.2	C23—C22—H22A	109.2
C12—C11—C10	112.66 (11)	O2—C22—H22B	109.2
C12—C11—H11A	109.1	C23—C22—H22B	109.2
C10—C11—H11A	109.1	H22A—C22—H22B	107.9
C12—C11—H11B	109.1	C24—C23—C25	111.00 (12)
C10—C11—H11B	109.1	C24—C23—C22	107.48 (11)
H11A—C11—H11B	107.8	C25—C23—C22	110.22 (12)
C13—C12—C11	111.29 (12)	C24—C23—C26	108.90 (12)
C13—C12—H12A	109.4	C25—C23—C26	110.06 (12)
C11—C12—H12A	109.4	C22—C23—C26	109.12 (11)
C13—C12—H12B	109.4	O1—C24—C23	112.89 (11)
C11—C12—H12B	109.4	O1—C24—H24A	109.0
H12A—C12—H12B	108.0	C23—C24—H24A	109.0
C12—C13—C14	110.08 (12)	O1—C24—H24B	109.0
C12—C13—H13A	109.6	C23—C24—H24B	109.0
C14—C13—H13A	109.6	H24A—C24—H24B	107.8
C12—C13—H13B	109.6	C23—C25—H25A	109.5
C14—C13—H13B	109.6	C23—C25—H25B	109.5
H13A—C13—H13B	108.2	H25A—C25—H25B	109.5
C13—C14—C15	111.34 (12)	C23—C25—H25C	109.5
C13—C14—H14A	109.4	H25A—C25—H25C	109.5
C15—C14—H14A	109.4	H25B—C25—H25C	109.5
C13—C14—H14B	109.4	C23—C26—H26A	109.5
C15—C14—H14B	109.4	C23—C26—H26B	109.5
H14A—C14—H14B	108.0	H26A—C26—H26B	109.5
C14—C15—C10	111.18 (12)	C23—C26—H26C	109.5
C14—C15—H15A	109.4	H26A—C26—H26C	109.5
C10—C15—H15A	109.4	H26B—C26—H26C	109.5
C14—C15—H15B	109.4	O1—B1—O2	122.35 (11)
C10—C15—H15B	109.4	O1—B1—C9	119.72 (11)
H15A—C15—H15B	108.0	O2—B1—C9	117.93 (11)
C17—C16—C21	108.63 (11)	B1—O1—C24	120.49 (10)
C17—C16—C9	112.83 (11)	B1—O2—C22	119.36 (10)
C21—C16—C9	114.57 (11)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$C22-H22B\cdots O2^i$	0.99	2.69	3.5773 (18)	150

Symmetry code: (i) $-x+1, -y, -z$.