

STRUCTURE  
REPORTS

ISSN 1600-5368

[journals.iucr.org/e](http://journals.iucr.org/e)

## Crystal structure of 2-[4-(methylsulfanyl)quinazolin-2-yl]-1-phenylethanol

**Gamal A. El-Hiti, Keith Smith, Amany S. Hegazy, Mansour D. Ajarim and  
Benson M. Kariuki**

*Acta Cryst.* (2014). **E70**, o1101



**IUCr Journals**  
CRYSTALLOGRAPHY JOURNALS ONLINE

This open-access article is distributed under the terms of the Creative Commons Attribution Licence <http://creativecommons.org/licenses/by/2.0/uk/legalcode>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original authors and source are cited.



# Crystal structure of 2-[4-(methylsulfanyl)-quinazolin-2-yl]-1-phenylethanol

Gamal A. El-Hiti,<sup>a\*</sup> Keith Smith,<sup>b</sup> Amany S. Hegazy,<sup>b</sup>  
Mansour D. Ajarim<sup>c</sup> and Benson M. Kariuki<sup>b\*</sup>

<sup>a</sup>Cornea Research Chair, Department of Optometry, College of Applied Medical Sciences, King Saud University, PO Box 10219, Riyadh 11433, Saudi Arabia,

<sup>b</sup>School of Chemistry, Cardiff University, Main Building, Park Place, Cardiff CF10 3AT, Wales, and <sup>c</sup>Criminal Evidence, Ministry of Interior, Riyadh 11632, PO Box 86985, Saudi Arabia. \*Correspondence e-mail: gelhiti@ksu.edu.sa, kariukib@cardiff.ac.uk

Received 17 August 2014; accepted 4 September 2014

Edited by J. F. Gallagher, Dublin City University, Ireland


In the molecule of the title compound, C<sub>17</sub>H<sub>16</sub>N<sub>2</sub>OS, the almost planar methylsulfanylquinazoline group [the methyl C atom deviates by 0.032 (2) Å from the plane through the ring system] forms an interplanar angle of 76.26 (4)° with the plane of the phenyl group. An intramolecular O—H...N hydrogen bond is present between the quinazoline and hydroxy groups. In the crystal, molecules are stacked along the *b*-axis direction.

**Keywords:** crystal structure; 4-(methylsulfanyl)quinazoline derivative; hydrogen bonding.

**CCDC reference:** 1022918

## 1. Related literature

For the synthesis of 4-(methylsulfanyl)quinazoline derivatives, see: Smith *et al.* (2005a,b); Leonard & Curtin (1946); Meerwein *et al.* (1956). For the crystal structures of related compounds, see: Alshammari *et al.* (2014a,b).



## 2. Experimental

### 2.1. Crystal data

C<sub>17</sub>H<sub>16</sub>N<sub>2</sub>OS  
*M<sub>r</sub>* = 296.38

Monoclinic, *P*2<sub>1</sub>/*n*  
*a* = 15.6142 (3) Å

*b* = 5.6142 (1) Å  
*c* = 17.2355 (3) Å  
*β* = 101.138 (2)°  
*V* = 1482.43 (5) Å<sup>3</sup>  
*Z* = 4

Cu *Kα* radiation  
*μ* = 1.93 mm<sup>−1</sup>  
*T* = 293 K  
0.32 × 0.19 × 0.14 mm

### 2.2. Data collection

Agilent SuperNova (Dual, Cu at zero, Atlas) diffractometer  
Absorption correction: multi-scan (*CrysAlis PRO*; Agilent, 2014)  
*T*<sub>min</sub> = 0.867, *T*<sub>max</sub> = 1.000

9606 measured reflections  
2938 independent reflections  
2688 reflections with *I* > 2σ(*I*)  
*R*<sub>int</sub> = 0.014

### 2.3. Refinement

*R* [*F*<sup>2</sup> > 2σ(*F*<sup>2</sup>)] = 0.032  
*wR* (*F*<sup>2</sup>) = 0.094  
*S* = 1.06  
2938 reflections

192 parameters  
H-atom parameters constrained  
Δρ<sub>max</sub> = 0.16 e Å<sup>−3</sup>  
Δρ<sub>min</sub> = −0.28 e Å<sup>−3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O1—H1...N1	0.82	2.12	2.7531 (15)	134

Data collection: *CrysAlis PRO* (Agilent, 2014); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS2013* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2013* (Sheldrick, 2008); molecular graphics: *Mercury* (Macrae *et al.*, 2006), *ORTEP-3 for Windows* (Farrugia, 2012) and *CHEMDRAW Ultra* (Cambridge Soft, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

## Acknowledgements

The authors thank the Cornea Research Chair, Department of Optometry, College of Applied Medical Sciences, King Saud University, for funding this research.

Supporting information for this paper is available from the IUCr electronic archives (Reference: GG2141).

## References

- Agilent (2014). *CrysAlis PRO*. Agilent Technologies, Yarnton, Oxfordshire, England.
- Alshammari, M. B., Smith, K., Hegazy, A. S., Kariuki, B. M. & El-Hiti, G. A. (2014a). *Acta Cryst. E* **70**, o919–o920.
- Alshammari, M. B., Smith, K., Hegazy, A. S., Kariuki, B. M. & El-Hiti, G. A. (2014b). *Acta Cryst. E* **70**, o953.
- Cambridge Soft (2001). *CHEMDRAW Ultra*. Cambridge Soft Corporation, Cambridge, Massachusetts, USA.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Leonard, N. J. & Curtin, D. Y. (1946). *J. Org. Chem.* **11**, 349–352.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Meerwein, H., Laasch, P., Mersch, R. & Nentwig, J. (1956). *Chem. Ber.* **89**, 224–238.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Smith, K., El-Hiti, G. A. & Hegazy, A. S. (2005a). *J. Sulfur Chem.* **26**, 121–131.
- Smith, K., El-Hiti, G. A. & Hegazy, A. S. (2005b). *Synthesis*, pp. 2951–2961.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

## supporting information

*Acta Cryst.* (2014). E70, o1101 [doi:10.1107/S1600536814019990]

**Crystal structure of 2-[4-(methylsulfanyl)quinazolin-2-yl]-1-phenylethanol**

**Gamal A. El-Hiti, Keith Smith, Amany S. Hegazy, Mansour D. Ajarim and Benson M. Kariuki**

**S1. Comment**


In  $C_{17}H_{16}N_2OS$  (Fig. 1), the methylsulfanylquinazoline group is planar and is oriented at an angle of  $76.26(4)^\circ$  with the phenyl group. The hydroxyl group forms an intramolecular hydrogen bond to one of the quinazoline ring nitrogen atoms (Table 1) whereas the second N atom is not involved in hydrogen bonding. In the crystal structure, molecules are stacked along the *b*-axis direction (Fig. 2).

**S2. Experimental**


Synthesis and crystallization: 2-(2-hydroxy-2-phenylethyl)-4-(methylsulfanyl)quinazoline was obtained in 83% yield from lithiation of 2-methyl-4-(methylsulfanyl)quinazoline with *n*-butyllithium at  $78^\circ\text{C}$  in anhydrous THF under nitrogen followed by reaction with benzaldehyde (Smith *et al.*, 2005*b*). Crystallization from a mixture of ethyl acetate and diethyl ether (1:3 by volume) gave the title compound as colorless crystals. The NMR and low and high resolution mass spectra for the title compound were consistent with those reported (Smith *et al.*, 2005*b*).

**S3. Refinement**

H atoms were positioned geometrically and refined using a riding model with  $U_{\text{iso}}(\text{H})$  constrained to be 1.2 times  $U_{\text{eq}}(\text{C})$  except for the methyl group where it was 1.5 times with free rotation about the C—C bond. For the OH group,  $U_{\text{iso}}(\text{H})$  1.5 times  $U_{\text{eq}}(\text{O})$  was used with free rotation about the C—O bond.

**Figure 1**

A molecule of  $C_{17}H_{16}N_2OS$  with atom labels and 50% probability displacement ellipsoids for nonhydrogen atoms.

**Figure 2**

Crystal structure packing viewed down the *b* axis.

### 2-[4-(Methylsulfonyl)quinazolin-2-yl]-1-phenylethanol

#### Crystal data

$C_{17}H_{16}N_2OS$

$M_r = 296.38$

Monoclinic,  $P2_1/n$

$a = 15.6142(3) \text{ \AA}$

$b = 5.6142(1) \text{ \AA}$

$c = 17.2355(3) \text{ \AA}$

$\beta = 101.138(2)^\circ$

$V = 1482.43(5) \text{ \AA}^3$

$Z = 4$

$F(000) = 624$

$D_x = 1.328 \text{ Mg m}^{-3}$

Cu  $K\alpha$  radiation,  $\lambda = 1.54184 \text{ \AA}$

Cell parameters from 5816 reflections

$\theta = 3.5\text{--}73.7^\circ$

$\mu = 1.93 \text{ mm}^{-1}$

$T = 293 \text{ K}$

Block, colourless

$0.32 \times 0.19 \times 0.14 \text{ mm}$

*Data collection*

Agilent SuperNova (Dual, Cu at 0, Atlas)  
diffractometer

$\omega$  scans

Absorption correction: multi-scan  
(*CrysAlis PRO*; Agilent, 2014)

$T_{\min} = 0.867$ ,  $T_{\max} = 1.000$

9606 measured reflections

2938 independent reflections

2688 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.014$

$\theta_{\max} = 74.0^\circ$ ,  $\theta_{\min} = 3.5^\circ$

$h = -19 \rightarrow 19$

$k = -6 \rightarrow 6$

$l = -11 \rightarrow 21$

*Refinement*

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.032$

$wR(F^2) = 0.094$

$S = 1.06$

2938 reflections

192 parameters

0 restraints

Hydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0507P)^2 + 0.2343P]$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.16 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.28 \text{ e } \text{\AA}^{-3}$

*Special details*

**Experimental.** Absorption correction: *CrysAlisPro*, Agilent Technologies, Version 1.171.36.28 (release 01-02-2013 *CrysAlis171 .NET*) (compiled Feb 1 2013, 16:14:44) Empirical absorption correction in *SCALE3 ABSPACK*.

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.67633 (8)	0.0261 (2)	0.10297 (7)	0.0458 (3)
C2	0.68632 (8)	−0.1600 (2)	0.04845 (7)	0.0450 (3)
C3	0.62090 (9)	−0.3225 (3)	0.01440 (8)	0.0553 (3)
H3	0.5652	−0.3110	0.0258	0.066*
C4	0.63923 (10)	−0.4967 (3)	−0.03523 (9)	0.0601 (4)
H4	0.5961	−0.6048	−0.0569	0.072*
C5	0.72249 (10)	−0.5136 (3)	−0.05366 (8)	0.0569 (3)
H5	0.7342	−0.6336	−0.0873	0.068*
C6	0.78660 (9)	−0.3561 (3)	−0.02273 (8)	0.0523 (3)
H6	0.8413	−0.3669	−0.0363	0.063*
C7	0.76994 (8)	−0.1775 (2)	0.02961 (7)	0.0444 (3)
C8	0.81859 (8)	0.1354 (2)	0.11131 (7)	0.0445 (3)
C9	0.59215 (12)	0.3019 (3)	0.19721 (11)	0.0734 (5)
H9A	0.6392	0.2626	0.2400	0.110*
H9B	0.5401	0.3312	0.2176	0.110*
H9C	0.6070	0.4421	0.1708	0.110*
C10	0.88942 (8)	0.3057 (2)	0.14745 (8)	0.0479 (3)
H10A	0.8870	0.3261	0.2029	0.058*
H10B	0.8775	0.4595	0.1220	0.058*
C11	0.98167 (8)	0.2298 (2)	0.14133 (7)	0.0429 (3)

H11	0.9941	0.0764	0.1683	0.051*
C12	1.04767 (8)	0.4094 (2)	0.18105 (7)	0.0414 (3)
C13	1.06970 (9)	0.6065 (2)	0.14024 (8)	0.0483 (3)
H13	1.0441	0.6280	0.0873	0.058*
C14	1.12977 (9)	0.7713 (2)	0.17802 (9)	0.0538 (3)
H14	1.1449	0.9011	0.1499	0.065*
C15	1.16716 (9)	0.7448 (2)	0.25659 (9)	0.0544 (3)
H15	1.2075	0.8556	0.2815	0.065*
C16	1.14426 (9)	0.5519 (3)	0.29825 (8)	0.0541 (3)
H16	1.1681	0.5350	0.3517	0.065*
C17	1.08585 (8)	0.3841 (2)	0.26030 (8)	0.0477 (3)
H17	1.0720	0.2526	0.2883	0.057*
N1	0.83603 (7)	−0.0253 (2)	0.06177 (6)	0.0475 (3)
N2	0.74020 (7)	0.1686 (2)	0.13399 (6)	0.0475 (3)
O1	0.99198 (7)	0.2036 (2)	0.06187 (6)	0.0606 (3)
H1	0.9563	0.1070	0.0393	0.091*
S1	0.57374 (2)	0.05921 (7)	0.12865 (2)	0.06185 (14)

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0408 (6)	0.0551 (7)	0.0426 (6)	0.0077 (5)	0.0110 (5)	0.0052 (5)
C2	0.0415 (6)	0.0531 (7)	0.0399 (6)	0.0044 (5)	0.0071 (5)	0.0040 (5)
C3	0.0463 (7)	0.0683 (9)	0.0517 (7)	−0.0049 (6)	0.0101 (6)	−0.0006 (6)
C4	0.0609 (8)	0.0649 (8)	0.0529 (8)	−0.0122 (7)	0.0070 (6)	−0.0060 (6)
C5	0.0654 (9)	0.0565 (8)	0.0487 (7)	0.0011 (7)	0.0105 (6)	−0.0077 (6)
C6	0.0504 (7)	0.0597 (8)	0.0474 (7)	0.0065 (6)	0.0107 (5)	−0.0063 (6)
C7	0.0414 (6)	0.0508 (7)	0.0405 (6)	0.0059 (5)	0.0065 (5)	0.0007 (5)
C8	0.0398 (6)	0.0514 (7)	0.0418 (6)	0.0076 (5)	0.0072 (5)	−0.0004 (5)
C9	0.0777 (11)	0.0691 (10)	0.0845 (11)	0.0061 (8)	0.0435 (9)	−0.0105 (8)
C10	0.0431 (6)	0.0510 (7)	0.0492 (7)	0.0052 (5)	0.0078 (5)	−0.0062 (5)
C11	0.0433 (6)	0.0459 (6)	0.0410 (6)	0.0028 (5)	0.0123 (5)	0.0009 (5)
C12	0.0376 (6)	0.0434 (6)	0.0451 (6)	0.0061 (5)	0.0129 (5)	−0.0006 (5)
C13	0.0508 (7)	0.0475 (7)	0.0482 (6)	0.0057 (5)	0.0139 (5)	0.0037 (5)
C14	0.0538 (7)	0.0434 (6)	0.0691 (8)	0.0010 (6)	0.0244 (6)	0.0017 (6)
C15	0.0451 (7)	0.0510 (7)	0.0677 (8)	−0.0014 (6)	0.0123 (6)	−0.0110 (6)
C16	0.0479 (7)	0.0604 (8)	0.0517 (7)	0.0045 (6)	0.0038 (6)	−0.0037 (6)
C17	0.0468 (6)	0.0489 (7)	0.0477 (6)	0.0037 (5)	0.0098 (5)	0.0039 (5)
N1	0.0390 (5)	0.0554 (6)	0.0482 (5)	0.0046 (4)	0.0084 (4)	−0.0067 (5)
N2	0.0429 (5)	0.0546 (6)	0.0462 (5)	0.0072 (5)	0.0116 (4)	−0.0021 (5)
O1	0.0655 (6)	0.0740 (7)	0.0477 (5)	−0.0177 (5)	0.0247 (4)	−0.0148 (5)
S1	0.0459 (2)	0.0755 (3)	0.0698 (2)	0.00317 (16)	0.02499 (17)	−0.00692 (17)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

C1—N2	1.3092 (17)	C9—H9C	0.9600
C1—C2	1.4340 (18)	C10—C11	1.5252 (16)
C1—S1	1.7521 (13)	C10—H10A	0.9700

C2—C7	1.4083 (17)	C10—H10B	0.9700
C2—C3	1.4100 (19)	C11—O1	1.4173 (14)
C3—C4	1.365 (2)	C11—C12	1.5082 (17)
C3—H3	0.9300	C11—H11	0.9800
C4—C5	1.400 (2)	C12—C17	1.3880 (17)
C4—H4	0.9300	C12—C13	1.3894 (18)
C5—C6	1.365 (2)	C13—C14	1.3873 (19)
C5—H5	0.9300	C13—H13	0.9300
C6—C7	1.4065 (18)	C14—C15	1.375 (2)
C6—H6	0.9300	C14—H14	0.9300
C7—N1	1.3713 (17)	C15—C16	1.384 (2)
C8—N1	1.3067 (16)	C15—H15	0.9300
C8—N2	1.3678 (16)	C16—C17	1.3843 (19)
C8—C10	1.5032 (18)	C16—H16	0.9300
C9—S1	1.7902 (17)	C17—H17	0.9300
C9—H9A	0.9600	O1—H1	0.8200
C9—H9B	0.9600		
N2—C1—C2	122.82 (11)	C11—C10—H10A	108.5
N2—C1—S1	119.58 (10)	C8—C10—H10B	108.5
C2—C1—S1	117.60 (10)	C11—C10—H10B	108.5
C7—C2—C3	119.24 (12)	H10A—C10—H10B	107.5
C7—C2—C1	115.14 (11)	O1—C11—C12	108.24 (10)
C3—C2—C1	125.62 (12)	O1—C11—C10	112.40 (10)
C4—C3—C2	120.08 (13)	C12—C11—C10	110.70 (10)
C4—C3—H3	120.0	O1—C11—H11	108.5
C2—C3—H3	120.0	C12—C11—H11	108.5
C3—C4—C5	120.48 (13)	C10—C11—H11	108.5
C3—C4—H4	119.8	C17—C12—C13	118.57 (12)
C5—C4—H4	119.8	C17—C12—C11	120.28 (11)
C6—C5—C4	120.70 (13)	C13—C12—C11	121.12 (11)
C6—C5—H5	119.7	C14—C13—C12	120.31 (12)
C4—C5—H5	119.7	C14—C13—H13	119.8
C5—C6—C7	120.01 (13)	C12—C13—H13	119.8
C5—C6—H6	120.0	C15—C14—C13	120.65 (13)
C7—C6—H6	120.0	C15—C14—H14	119.7
N1—C7—C6	119.03 (12)	C13—C14—H14	119.7
N1—C7—C2	121.49 (11)	C14—C15—C16	119.52 (13)
C6—C7—C2	119.47 (12)	C14—C15—H15	120.2
N1—C8—N2	126.28 (12)	C16—C15—H15	120.2
N1—C8—C10	118.73 (11)	C15—C16—C17	119.99 (13)
N2—C8—C10	114.98 (11)	C15—C16—H16	120.0
S1—C9—H9A	109.5	C17—C16—H16	120.0
S1—C9—H9B	109.5	C16—C17—C12	120.92 (12)
H9A—C9—H9B	109.5	C16—C17—H17	119.5
S1—C9—H9C	109.5	C12—C17—H17	119.5
H9A—C9—H9C	109.5	C8—N1—C7	117.30 (11)
H9B—C9—H9C	109.5	C1—N2—C8	116.95 (11)



C8—C10—C11	115.00 (10)	C11—O1—H1	109.5
C8—C10—H10A	108.5	C1—S1—C9	102.08 (7)

*Hydrogen-bond geometry (Å, °)*

<i>D—H···A</i>	<i>D—H</i>	<i>H···A</i>	<i>D···A</i>	<i>D—H···A</i>
O1—H1···N1	0.82	2.12	2.7531 (15)	134