

# General Botany (Bot 102)

علم النبات العام  
( ١٠٢ نبت )

د. سعود العمري مكتب أب ٣٥

٤٦٧٥٨٦٦

# Course specification- an overview

## عرض توصيف المقرر

### A. Basic information:

- Lecture: 2hour
- Labs: 2 (1x2) hours
- Total credit hours: 3**

### B. Professional information:

## (1) Goals & Objectives

- The course provides background material for students who have not previously been exposed thoroughly to basic Botany at university level.
- Knowledge of the fundamental and applied aspects of different botany fields of study, considering the levels of organization, plant structure and function, classification of the plant kingdom and interactions between plants and their environment.

## **(2) Intended Learning Outcomes (ILO'o)**

### **المخرجات التعليمية المستهدفة**

#### **a. Knowledge and understanding:**

- **Knowledge of fundamental and concepts of Botany.**
- **Demonstrate the ability of how to utilize the theoretical concepts in applicable form.**
- **Knowledge of the major fields of Botany.**
- **What characters link plants to each other and to their environment? How flexible are these links? And how intricate or intermeshed? How do plants evolve and function in their environment? ....their reproduction? ...acquisition of energy and nutrients? .....?**

## **b. Intellectual skills:**

- **Knowledge integration and evaluation of Botanical processes at different levels of organization and classification from molecules to biomes (zonobiomes).**
- **Testing hypothesis and solve problems with self-direction and originality.**
- **What can organisms tell us about the past, present and future course of their existence?**



### **c. Professional and practical skills:**

- **Ability to work in laboratory and field either independently or as a member of a team.**
- **Ability to do research and report on many areas of Botanical and biological sciences**

## **d. General and transferable skills:**

- **Problem analysis and solving at theoretical and practical levels.**
- **Learn in familiar and unfamiliar situation with open mentality and in the spirit of critical enquiry**
- **Know how to cope with situations (bad or good) and accept to live with others.**

### **(3) Course content**

**Course content covers the basic principles of Botany that deal with:**

- **levels of organization from molecules to biomes**
- **Plant cell structure and division**
- **Basics of Mendel's Genetics**
- **Plant structure (Morphological & Anatomical perspectives) and function (basics of plant physiology)**
- **Classification of plant kingdom**
- **Plant environment and interactions among plants.**

## **(4) Teaching and Learning Methods**

- **Lectures**
- **Laboratory studies**
- **Student group assignment**
- **Class discussion and reading materials**
- **Term papers (internet search)**

## **(5) Student Assessment Methods**

- **Written exams and quizzes:** to assess the ability to manage and present the understanding in appropriate manner.
- **Practical exams and assignments:** to assess the skills and abilities and evaluate the outreach of used methodology.
- **Feedback questionnaires:** to assess the student satisfaction, efficiency of instructors and suitability of course contents.
- **Oral exam or group discussion (optional):** to assess the ability to present knowledge and understanding.

## ❖ **Assessment schedule:**

**-Monthly exams and assignments+ class attendance 30**

**-Laboratory exam and assignment 30**

**-Final written exam 40**

**Total Marks**

**100**

# List of references

## **(A) Course notes:**

**Lecture and laboratory hand outs and assigned reading materials**

## **(2) Text books**

- 1-Plant Biology in brief. Mohammed H. Al-whaibi (2011). King Saud University, Academic publishing & Press.**
- 2- Biology of plants 5th Ed. by Raven, P.H., Evert, R.F. and Eichhorn, S.E.(1992) W.H. Freeman and company, Worth Publishers. New York**

## **Supporting books:**

- 3-Biology of Plants 5th ed. by Raven et al. 1992 Worth Publishers. (Translation into Arabic by Al-Whaibi, M. H. and A. S. Al-Khalil, 2002., (2005 second Ed.) Scientific Publications, King Saud University Press, Riyadh, Saudi Arabia. (In Arabic).**
- 4-Manual of Biology of Plants.2002. (2005 second Ed.) Arif, I. A., A. S. Al-Khalil, Al-Whaibi, M. H., R. M. Al-Summ and K. M. Zayed. Scientific Publications, King Saud University Press, Riyadh, Saudi Arabia. (In Arabic).**
- 5- Study Guide to Plant Biology (In Brief). 2008. Al-Whaibi, M. H. and A. S. Al-Khalil) Scientific Publications, King Saud University Press, Riyadh, Saudi Arabia. (In Arabic).**
- 6- Raven, P.H., Evert, R.F. and Eichhorn, S.E.(1999). Biology of plants 6th. E. W.H. Freeman and company, Worth Publishers. New York.**
- 7. Raven, P.H., Evert, R.F. and Eichhorn, S.E.(2005). Biology of plants 7th. E. W.H. Freeman and company, Worth Publishers. New York**

## **(C) Periodicals & Websites:** To be listed and handed out during lecture time.




# Why study plants?






# **We could not live without plants**

- Plants produce most of the oxygen we breathe.
- Plants produce most of the chemically stored energy we consume as food and burn for fuel.
- Plants produce an assortment of useful chemicals.




# Plants are diverse





Plants have evolved the ability to thrive in diverse land habitats.

**Plants fix carbon dioxide into energy- rich molecules where human & animals can use as food**



**Plants convert  $\text{CO}_2$  gas into sugars through the process of photosynthesis.**

# Plants can produce an amazing assortment of chemicals





# More reasons .....?



- ❖ **To help conserve endangered plants and threatened environments**
- ❖ **To learn more about the natural world**
- ❖ **To enhance the abilities of plants to provide us with food, medicines, and energy**

# **Mendel's studies of peas revealed the laws of inheritance**



# **Plant scientists can contribute to the alleviation of hunger**


**By developing plants that**

- **are drought or stress tolerant**
- **require less fertilizer or water**
- **are resistant to pathogens**
- **are more nutritious**



# Identification of resistance genes

Geneticists have identified the gene conferring resistance and are introducing it into edible varieties.



The plant on the left carries the resistance gene and is free from disease symptoms.



# **Plant biologists study ways to keep plants fresh after harvesting**



**After harvesting, fruits soften, ripen, and eventually rot.**

**These processes make the fruit less appealing and affect the nutritional qualities.**




# Genetically biofortified foods

**Iron-enriched rice**



**Vitamin A–enriched rice**



**Wild-type (top) and  
antioxidant-enriched  
tomatoes**

# Plants provide us with more than food




## Plants:

- are sources of novel **therapeutic drugs**
- provide better **fibers** for paper or fabric
- are sources of **biorenewable** products
- provide **renewable energy** sources

# **Plants can replace petroleum for many products and purposes**


**Petroleum is NOT a renewable resource**



**Unfortunately, it takes millions and millions of years to convert dead organic material into petroleum...and we are running out of it.**



# Plants can be a source of biofuels



# **What is Life?**

**Life is a characteristic that distinguishes objects that have self-sustaining biological processes, other things are classified as inanimate.**

# **Properties of life .....**

- **Cellular structure (unit of life)**
- **Metabolism (perform function)**
- **Movement (intracellular)**
- **Homeostasis**
- **Organization**
- **Growth**
- **Adaptation**
- **Behavior (response to stimuli)**
- **Reproduction (avoid extinction)**
- **Pass on their traits to offspring (heredity)**


# **PLANT CELL**

**CELLULAR COMPONENTS & PROCESSES**



# **The cell theory states:**


- 1. The cell is the basic unit of structure and function of all living things.**
- 2. All living things are composed of one or more cells.**
- 3. All cells come from pre-existing cells.**
- 4. The cells of all living things carry on similar chemical activities.**
- 5. All cells carry on their metabolic activities in organelles.**



(a) A T-even bacteriophage


Copyright © 2001 Benjamin Cummings, an imprint of Addison Wesley Longman, Inc.

## Animal Virus Structure



**A virus and a prion are not considered cellular nor living organisms because of their simplicity (only Nucleic acid surrounded by a protein coat in the case of a virus or only a single strand of protein in the case of a prion). Neither exhibit characteristics of life unless they are in a host cell and cannot replicate outside the host cell.**

# types of cells:



- Which **ONE** is more complicated?

# There are two types of cells:


1. **Prokaryotic-cells** that DO NOT have a well-defined NUCLEUS or other cell ORGANELLES
  2. **Eukaryotic-cells** have a NUCLEUS with nuclear membrane & cell ORGANELLES
- Which is more complicated?

# Prokaryote cells

1. Oldest of cell types, first appeared 3.5 billion years ago.
2. Cells that do not contain a nucleus.
3. DNA is not contained in an internal structure.
4. Have a cell membrane.
5. Do not have membrane-bound organelles.
6. Generally smaller and simpler than eukaryotic cells.
7. Can live in hostile environments. Halophiles and thermophiles that are archeabacteria.
8. Very diverse in their metabolic process: **obligate aerobes** (require  $O_2$ ), **obligate anaerobes** (killed by  $O_2$ ), and **facultative anaerobes** (can survive with or without  $O_2$ ).
9. Example: Bacteria



# Prokaryotic Cell Structure




# Eukaryote cells

1. First appeared in the fossil record 1.5 billion years ago.
2. Eukaryotes are organisms that have a nucleus in each cell.
3. The nucleus contains that cell's DNA.
4. Have a cell membrane.
5. Generally larger and more complex than prokaryotic cells.
6. Have complex membrane bound organelles (mitochondrion, chloroplast, Golgi apparatus, etc.)
7. Many eukaryotic cells are highly specialized.
8. Examples: Plants, animals, fungi, and protists.



# Eukaryotic Cell Structure and Function

- Plasma Membrane
- Nucleus
- Ribosomes
- Nucleolus
- Endoplasmic Reticulum
- Golgi Bodies
- Plastids
- Chloroplasts
- Mitochondria
- Vacuoles
- Microtubules
- Cytoplasm





# Cell Wall



# **Cell wall consists of:**

**(1) Middle lamella** – mostly pectin, cements adjacent cells together


**(2) Primary cell wall**

- Found in all plant cells
- Cellulose matrix with hemicellulose, proteins, pectin, lignin, cutin, and wax
- Characteristic of undifferentiated cells or ones that still are growing


**(3) Secondary cell wall**

- Just inside primary cell wall
- Characteristic of mature cells
- Comprised of hemicellulose and lignin

## Connections between Cells: *Plasmodesmata*



# Cell or Plasma Membrane Structure






# Cell or Plasma Membrane

- The cell membrane's function is to form a **barrier between the cell's inner and outer environment**. It is **selectively permeable** meaning that it allows certain materials to pass through and prevents the movement of other through it.
- It is composed of a **phospholipid bilayer with protein molecules** (integral proteins) embedded within in the bilayer. Some of these proteins pass completely through both layers of phospholipids. There are also other types of molecules such as **cholesterol and carbohydrates** that are associated with the cell membrane's outer surface.
- The **phospholipids and proteins** are not in a **static state**, but have the ability to **move from one location to another** or change positions within the bilayer. Therefore the molecules which make up the membrane are described as being in a **fluid state**. The structure of the membrane is called the "fluid-mosaic model." **The membrane is literally a mosaic of molecules that have the ability to move from area to area on the surface of the membrane.**

# Cytoplasm

A watery solution made of cytosol that contains the **cell organelles**. Cytoplasm includes salts, an assortment of organic molecules, including many enzymes that catalyze reactions, as well as water







# Cytoskeleton

## Cytoskeleton:

The cytoskeleton is a “framework” that supports the cell membrane and other cell structures within the cytoplasm.



Microtubule



Microfilaments

0.25  $\mu\text{m}$

# Cytoskeleton

- Eukaryotic cells are given their **shape** and internal organization by the cytoskeleton.
- The cytoskeleton is made up of:
- **Microfilaments** and **microtubules**

## Microfilaments


- are **threadlike** structures made up of the protein **actin**.
- form extensive networks in some cells.
- produce a tough, flexible framework that **supports** the cell.
- help some cells **move**.

## Microtubules

- are **hollow** structures made up of proteins known as **tubulins**.  
maintain cell shape.
- are important in cell **division**.
- build projections from the cell surface—**cilia and flagella**—that enable some cells to swim rapidly through liquids.


The Nucleus is enclosed in an envelope which is a double membrane structure. It has pore complexes in the membranes which allow the movement of materials in and out of the structure. It contains DNA and proteins in the form of loose threads called chromatin. During mitosis or meiosis the chromatin super coils to form chromosomes. Self duplicating structure divides when the cell divides. The nucleolus is a structure composed of RNA located in the nucleoplasm. There maybe be more than one present and it functions in the production of ribosomes. The overall function of the nucleus is the regulation of cellular activities.

# Nucleus



# Cell Organelles

# Plastids





# Plastids

Plastids are structures that function in storage and photosynthesis according to their type.

Amyloplasts (Leucoplastides) are large white structure where starch is stored. They are responsible for the color of an Irish potato.


Chromoplasts contain pigments and are responsible for the orange and yellow colors of fruits and flowers.

Chloroplasts are double membrane structures where the process of photosynthesis occurs.




# Plastids

The inner membrane is arranged in flattened sacs called **thylakoids**. The thylakoids are stacked one on top of another. A stack of thylakoids is called a **granum** or **grana** (pl). The space in between the grana is called the **stroma**



# Mitochondria


The mitochondria is a **double membrane** structure with an outer membrane which surrounds a highly folded inner membrane. It is the site of aerobic cellular respiration in which **ATP is produced**. The inner membrane has finger like projection called **cristae** which increase the surface area. The inner space within the mitochondrion is called the **matrix** and contains cytoplasm, ribosomes, and DNA. Mitochondrion are self replicating. They are found in both plant and animal cells and are sometimes called “**the powerhouse of the cell**”.



Copyright © Pearson Education, Inc., publishing as Benjamin Cummings.

# Golgi Apparatus (Dictyosome)


The Golgi apparatus appears as a series of **flattened, stacked, membrane sacs**. The Golgi apparatus is the **center for manufacturing, modifying, and packaging of materials for transport**. It receives secretory proteins from the RER and modifies and packages the materials in small secretory vesicles. It is found in both animal and plant cells. In plant cells it may be referred to as **Dictyosome**.





# Ribosomes

Ribosomes are the structures within the cell which **read mRNA and assemble amino acids into polypeptide chains**. They are found **free floating** in the cytoplasm **or attached** to the nuclear envelope or the rough endoplasmic reticulum. They are found **in all prokaryotic and eukaryotic cell types**. They are composed of two subunits. Prokaryotic cells have smaller ribosomes (70s) and eukaryotic cells have the larger (80s) form.




# Endoplasmic Reticulum

The endoplasmic reticulum (ER) is a series of **single membrane channels** which run throughout the cytoplasm of the cell.


The smooth endoplasmic reticulum (SER) is free of ribosomes and functions in **lipid synthesis, metabolism of carbohydrates, and as a detoxification center** of the cell.

The rough endoplasmic reticulum (RER) has ribosomes bound to its outer membrane layer and is the active site of **protein synthesis**. These are **secretory proteins** which will be released by the cell. Both forms of endoplasmic reticulum are found in plant and animal cells.



Vacuoles are **storage areas** and can also serve as the **site of chemical digestion** within the cell itself. Vacuoles in animal cells are often small. However, plant cells often have a large centrally located vacuole which contains **water and dissolved solutes**, surrounded by a membrane called the **tonoplast**. Freshwater **Protists** contain specialized vacuoles which act as “**water pumps**” to remove excess water that enters their cytoplasm. These specialized vacuoles are called **contractile vacuoles**.

# Vacuoles



Copyright © Pearson Education, Inc., publishing as Benjamin Cummings.



# Self-Replicating Organelles

## • Mitochondria

- Involved in energy release



## • Plastids

- Involved in energy capture and storage.

– Chloroplasts

– Amyloplasts

– Chromoplasts



# **The Cellular Basis of Reproduction and Inheritance**

# Cell Reproduction

## The Cell Cycle

### A. Growth

- Increase in cell size.

### B. Division


- Production of new cells
- Two overlapping processes
  - Karyokinesis – nuclear division
  - Cytokinesis – cytoplasm division

# Methods of Reproduction

## Asexual reproduction

- Chromosomes are duplicated and cell divides
- One copy of each chromosome is placed in each cell
- Each “daughter” cell is genetically identical to the parent and the other daughter
  - Type of Cellular Division required: mitosis

**Advantage = fast and convenient**  
**Disadvantage = very little genetic variation**



# **Sexual reproduction**

- **Offspring inherit DNA from both of their parents**
  - **Type of Cellular Division required: meiosis**
- **Offspring can show great variation**
  - **Advantage = lots of genetic variation**
  - **Disadvantage = metabolically expensive**



# Related Terms

- **Chromatin**


- **Material in an active nucleus.**
  - Submicroscopic “threads” consisting of 50% DNA and 50% supporting proteins.
  - Abundant water and dissolved chemicals.

- **Gene**

- a unit of heredity information determining the nature of a specific trait and have specific places on chromosomes.
- a section of DNA that codes for a protein, tRNA or rRNA molecule

- **DNA Replication**

- Conversion of one strand/piece of DNA into two identical strands/pieces.



- **Chromosome Set**

- One copy of each of the different chromosomes in the nucleus containing one copy of each different gene.

- **Haploid Number ( $n$ )**

- The number of chromosomes comprising one set.
  - For humans,  $n=23$
  - For some ferns,  $n=250$
- A haploid individual has one set of chromosomes per cell.

- **Diploid Number ( $2n$ )**

- The number of chromosomes in a cell containing two sets.
- A diploid individual has 2 sets per cell.
- (Triploid is 3 sets, Tetraploid is 4 sets, etc.)

# Structure of the Chromosome

**Chromosome** – a package of hereditary material with supporting proteins visible in condensed form during cell division.




**Chromatid** – a single strand of DNA

During most of the life of a cell the chromosomes exist as a single strand called a “monad”.

At the beginning of karyokinesis the single strand is replicated forming two identical chromatids attached to one another, a “dyad”.

Sister chromatids have identical DNA  
Centromere  
Kinetochore on centromere provides binding site for microtubules

**Centromere**




**Kinetochore**

- **Genome:** Complete complement of an organism's DNA.
  - Includes genes (control traits) and non-coding DNA organized in chromosomes.



- **Eukaryotes & DNA:**

- Many eukaryotes have 1000 times as much DNA as prokaryotes.
- DNA is located in the nucleus in the form of chromosomes.
- Chromosomes are DNA wound tightly around proteins called histones.





# •Homologues

- Chromosomes exist in homologous pairs in diploid cells.




Exception: **Sex** chromosomes in human (X, Y).

Other chromosomes are known as **autosomes**, they have homologues.

# **Cell Division**

**Binary Fission, Mitosis & Meiosis**

**Prokaryotic cells reproduce asexually  
by a type of cell division called binary  
fission**



## **binary fission**

- The circular DNA molecule replicates to form 2 chromosomes
- The chromosome copies move apart
- The cell elongates
- The plasma membrane grows inward, dividing the parent into two daughter cells


# Mitosis

- Eukaryotes divide by a more complicated system called **Mitosis**
- This is because:
  1. They have a nucleus which must be broken up and then reformed
  2. They have their DNA “packaged” in the form of **Chromosomes**
  3. Chromosomes are composed of **Chromatin**
  4. Also contain **Nucleosomes** containing **Histones** - Proteins the DNA is wrapped around Name for the DNA/Protein complex is **Chromatin**
  5. They usually have more than 1 chromosome (Humans have 23 pairs)
  6. They have numerous organelles to equally share




# The Cell Cycle

- Most of the cell's life is spent doing its regular function.
- Cells divide along a rough time frame called its Cell Cycle.
- The Cell cycle consists of the following steps:
  - G1 (Gap 1) Phase** - Cell performs its **normal function** (cells which do not divide stay in this stage for their entire life span)
  - S (Synthesis) Phase** - Here the cell actively **duplicates its DNA** in preparation for division
  - G2 (Gap 2) Phase** - Amount of cytoplasm (including organelles) increases in **preparation for division**.
  - Mitosis** - Actual division occurs




# Interphase

- **Cell Replicates its DNA/Chromosomes in preparation of upcoming division**




# Prophase

1. Chromosomes Shorten and become visible.
2. Centrosome move to opposite sides of the cell
3. Nuclear envelope disappears
4. Spindle Fibers & Astral Fibers both together are known as the Spindle Apparatus begin to form



# Metaphase


- Chromosomes line up along center of cell called the Metaphase Plate
- Chromosomes attach to spindle fibers
- Spindle & Astral fibers are now clearly visible





# Anaphase


- Centromeres break up separating chromosome copies
- Chromosomes are pulled apart to opposite sides of cell
- Spindle & Astral fibers begin to break down





# Telophase (cytokinesis)


- Nuclear envelope forms around both sets of chromosomes
- DNA uncoils
- Spindle & Astral fibers completely disappear
  - Cytokinesis happens with most (but not all) cells
  - Cytoplasm & organelles move (mostly equally) to either side of the cell. Cell Membrane “pinches” to form 2 separate cells















# Plant Cytokinesis- division of the cytoplasm

- With Plants, a cell wall must be formed between the 2 daughter cells.
- Vesicles containing Cellulose form and fuse between the two daughter cells, eventually forming a complete cell wall.

- ❑ Vesicles containing cell wall material line up across middle of cell
- ❑ Vesicles merge and form cell plate
- ❑ Cell plate grows until it divides the cell in 2



# Overview of Mitosis

Prophase	Prometaphase	Metaphase	Anaphase	Telophase	Cytokinesis
					
<ul style="list-style-type: none"> <li>Chromosomes condense and become visible</li> <li>Spindle fibers emerge from the centrosomes</li> <li>Nuclear envelope breaks down</li> <li>Centrosomes move toward opposite poles</li> </ul>	<ul style="list-style-type: none"> <li>Chromosomes continue to condense</li> <li>Kinetochores appear at the centromeres</li> <li>Mitotic spindle microtubules attach to kinetochores</li> </ul>	<ul style="list-style-type: none"> <li>Chromosomes are lined up at the metaphase plate</li> <li>Each sister chromatid is attached to a spindle fiber originating from opposite poles</li> </ul>	<ul style="list-style-type: none"> <li>Centromeres split in two</li> <li>Sister chromatids (now called chromosomes) are pulled toward opposite poles</li> <li>Certain spindle fibers begin to elongate the cell</li> </ul>	<ul style="list-style-type: none"> <li>Chromosomes arrive at opposite poles and begin to decondense</li> <li>Nuclear envelope material surrounds each set of chromosomes</li> <li>The mitotic spindle breaks down</li> <li>Spindle fibers continue to push poles apart</li> </ul>	<ul style="list-style-type: none"> <li>Animal cells: a cleavage furrow separates the daughter cells</li> <li>Plant cells: a cell plate, the precursor to a new cell wall, separates the daughter cells</li> </ul>
					
MITOSIS					



# Overview of Mitosis

- ❑ Occurs in somatic cells
- ❑ Longitudinal division of replicated chromosomes in one nucleus to form two genetically identical daughter nuclei.
- ❑ Each “daughter” nucleus has the same number of chromosomes (and sets) that the “parent” nucleus had.
- ❑ Mitosis requires One division.
  - 1 cell → 2 cells (*called daughter cells*)
  - Daughter cells are genetically identical
  - Chromosome number does not change.


# Meiosis

- Similar in many ways to mitosis
- Several differences
- Involves 2 cell divisions
- Results in 4 cells with  $\frac{1}{2}$  the normal genetic information



# Meiosis Phases


- Meiosis occurs in 2 phases; **Meiosis I**, & **Meiosis II**.
- **Meiosis I.**
  - Prior to division, amount of DNA doubles



# Crossing Over

- During metaphase 1 homologous chromosomes line-up along the metaphase plate
- Areas of homologous chromosomes

Chiasmata  
called cniasmata





# Crossing over (contd.)


Exchange of genetic material  
between Homologous Chromosomes

- During Prophase I


*occurs at CHIASMA*



Produces new genetic combinations  
--Chromosomes with both  
Maternal & Paternal components







Crossing over occurs  
in meiosis I


Homologous chromosomes  
separate in meiosis I

2 cells,  $N = 1$  for  
each

Sister chromatids  
separate in meiosis  
II

4 cells,  $N = 1$  for each.  
Chromosomes are different  
due to crossing over





# **Classic (Mendel) Genetics**






















# Gregor Mendel: Father of Genetics

- Genetics is the scientific study of heredity.
- Gregor Mendel (1860's) an Austrian Monk, was interested in figuring out how heredity was determined in plants and animals.
  - used pea plants
  - quantitative approach to collect data.
- Mendel studied seven different pea plant traits.
  - Seed shape & color, pod shape & color, plant height, flower color and seed coat color
- A trait is a specific characteristic, such as seed color or plant height, that varies from one individual to another.

# Gregor Mendel's Experiment

- He called the offspring of the **P-generation**, the **F1**, or “first filial,” generation. *Filius* is the Latin word for “son.”
  - These pea plants were cross pollinated.
  - In **cross-pollination**, male sex cells in pollen from the flower on one plant fertilize the egg cells of a flower on another plant.
- The offspring of crosses between parents with different traits are called **hybrids**.
- The F2 generation was allowed to **self-pollinate** (on the same plant).
- Out of 929 F2 Generation plants, 705 were **purple** and 224 were **white**.
  - Ratio of **3** purple to **1** white flowers

### Mendel's Seven F<sub>1</sub> Crosses on Pea Plants

	Seed Shape	Seed Color	Seed Coat Color	Pod Shape	Pod Color	Flower Position	Plant Height
P	Round  X 	Yellow  X 	Gray  X 	Smooth  X 	Green  X 	Axial  X 	Tall  X 
F <sub>1</sub>	 Round	 Yellow	 Gray	 Smooth	 Green	 Axial	 Tall

**Mendel's F<sub>1</sub> Crosses** When Mendel crossed plants with contrasting characters for the same trait, the resulting offspring had only one of the characters. 🌱 From these experiments, Mendel concluded that some alleles are dominant and others are recessive.



## **Mendel's conclusions: Rules**

### **1. Rule of Unit Factors**

Each organism has 2 factors for each of its traits  
(alleles: gene alternatives)

### **2. Rule of Dominance**

For each trait there exists 2 possible factors that are expressed in physical characters, one that may be dominant, and the other recessive.


### **3. Law of Segregation**

The two alleles for each trait must separate when gametes form.

# Expression of Traits

1. **Phenotype**: physical expression of a gene
2. **Genotype**: a make of genes on a chromosome
3. **Homozygous**: alleles for a trait are the same
4. **Heterozygous**: alleles for a trait are opposite

# Dihybrid cross



# Punnett Square for Dihybrid Cross

(Cross between 2 parents that are Heterozygous for two traits)

Law of Independent assortment – different traits are passed independently of each other. All possible combinations of gametes with the two traits must be considered possible.

Round Yellow Seeds X Round Yellow Seeds

RrYy      X      RrYy



# Did this mean that the two dominant alleles would always stay together?


When Mendel let the  $F_1$  self pollinate, he got a definite ratio of visible phenotypes:

**9** with both dominant  
(RY)-Round Yellow

**3** one dominant and one recessive  
(Ry)-round green

**3** one recessive and one dominant  
(rY)-wrinkled yellow

**1** both recessive  
(rryy)-wrinkled green





## **Mendel's Dihybrid Crosses**

**(cross involving two different traits)**

- A. experimenting with plants that had two different traits that differed from each other
  - 1. used true-breeding pea plants that had round yellow seeds (RRYY)
  - 2. crossed with true-breeding pea plants that had wrinkled green seeds (rryy)
  - 3. already knew that round was dominant and wrinkled was recessive and yellow was dominant and green was recessive

**B.  $F_1$  generation produced all round yellow seeds**

**C.  $F_2$  generation produced**

**9 round yellow**

**3 round green**

**3 wrinkled yellow**

**1 wrinkled green**

**D. The Law of Independent Assortment**

**Genes for different traits are inherited independently of each other.**

# **Plant Structure (Morphology & Anatomy)**


# Plant Morphology

## Shoot system


- **Stem**
  - Supports and places leaves
  - Transports H<sub>2</sub>O and nutrients
- **Leaves**
  - Photosynthesis
- **Reproductive structures - Flowers**

## Root system

- Anchors the plant
- Absorbs water and minerals
- Storage (CHO) & synthesis of some hormones
- Propagation



# Root & Shoot Tips





# Dicots & Monocots

Flowering plants possess three kinds of vegetative (non-reproductive) organs: roots, stems, and leaves.


The flower is the reproductive organ of the Angiosperms.

**Shoots consist of:**


**Stems and leaves;**

**Functions are photosynthesis, support, reproduction, storage and transport**

**Roots Functions are anchorage, and absorption of water and Minerals, provides nutrients for the shoot and can be an area of storage**




# Seed Structure



- A **seed** is a sporophyte embryo with its own food supply in a protective coat
- Seed plants (gymnosperms and angiosperms) retain their spores


# Plant Seedling




- **Monocotyledons (Monocots)- have a single seed leaf**
- **Dicotyledons (Dicots)- have double seed leaves**

# Seed Germination

## Monocotyledon (corn)



## Dicotyledon (bean)



# Roots

**Function** - absorption, anchorage

**Structure** – root cap, root hairs, endodermis


**Adventitious roots** - arise from non-root organ to perform specific function

**Lateral roots** - arise from another root (1°, 2°, etc.)





# Roots: Specialized for H<sub>2</sub>O & Nutrient Absorption

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.




# Root Types: Adventitious & Tap Root System

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



# Fibrous Root System


- Develops when the secondary roots become the main roots.
- Shallow roots but spread over a broad area.
- Helps prevent erosion.



FIBROUS

# Tap Root System

- Develops from the primary root.
- Reaches deep into the ground
- Helps the plant during periods of drought.





# Root Modifications:

## Storage Roots




Storage roots *Raphanus sativus*, radish



Storage roots, Sweet potato



# Carrot: Tap Root modification



Function: Storage of water.

Carrot plants are often associated with very sandy soils.

The enlarged root is familiar to those who have eaten the vegetable.

The root modification allows the storage of water in the cortex and central stele.

The mass of the root stabilizes the plant in the loose sandy soils.

## buttress roots



buttress roots, rusty-leaved fig



# Respiratory roots



Pneumatophores (respiratory  
roots)

*Avicennia marina*, black mangrove

# Prop roots



**Prop roots (also adventitious)**



# Haustoria – parasitic roots

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.





# Root Adaptations .....??



(a)



(b)



(c)



(d)



(e)

# **Shoot**

## **(Stem + Associated Leaves)**


# Stems

The two main functions of the stems are **conduction** and **support**


- **Conduction** involves moving substances manufactured in the leaves through the phloem to other parts of the plant including developing leaves, stems, roots, developing flowers, seeds and fruits and the xylem carries water from the roots to the leaves, where water is transpired
- **Support** involves holding the plant off the ground - supporting the principal photosynthetic organs of the plant (the leaves) as well as flowers, seeds and fruits

# Stem Habit

=relative position of stem (+ growth, structure)



**acaulescent**




**caulescent**



**cespitose**




**prostrate**




**repent**



**arborescent**




**suffrutescent**



**decumbent**


# Stem Branching






# Apical Dominance

Usually the growing terminal bud inhibits the development of the lateral buds, a phenomenon known as **apical dominance** – as the influence of the apical meristem lessens the growth of the lateral buds which proceed with their development.



**strong apical dominance:**  
**Jeffrey pine**




Waddington family arboretum website

**weak apical dominance:**  
**silver maple**


# Stem (Shoot) Types & Modifications

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.


## Types of Modified Stems (1)



## Types of Modified Stems (2)




**Runners (strawberry)**



**Tubers (potato)**




# Types of Modified Stems (3)



**Tendrils**

**Tendrils (grape)**


**Leaves (modified as spines)**



**Cladophyll**

**Cladophylls  
(prickly pear)**

# Onion Bulb






## Compare .....???

Kingsley R. Stern, Botany Visual Resource Library © 1997 The McGraw-Hill Companies, Inc. All rights reserved.

# Types of Specialized Stems



# Leaves




# Leaves

- **The leaf- Is the main photosynthetic organ of most vascular plants**
- **Leaves generally consist of a flattened blade and a petiole, which joins the leaf to a node of the stem**
- **Leaves may appear different based on whether they grow in shade or full sun**
- **Some plant species have evolved **modified leaves** that serve various functions**
  - **Climbing, pollinator attraction, storage, digestion, prevention of water loss, etc.**

# Leaves


## External Parts of the Leaf:

- **Petiole** عنق
  - Leaf stalk or part that connects the leaf to the stem.
- **Blade** نصل
  - The large, flat part of a leaf.
- **Midrib** عرق وسطى
  - The large center vein.





# Leaf Types




**(a) Simple leaf.** ورقة بسيطة A simple leaf is a single, undivided blade. Some simple leaves are deeply lobed.

**(b) Compound leaf (Pinnate).** مركبة ريشية In a compound leaf, the blade consists of multiple leaflets. Note that a leaflet has no axillary bud at its base.


**(c) Doubly compound leaf (Bipinnate).** مركبة ريشية مزدوجة In a doubly compound leaf, each leaflet is divided into smaller leaflets.



# Compound Leaves



palmately  
compound



pinnately  
compound




Bi-Pinnately Compound Leaf

# Leaf Venation

- **Parallel-** متوازي veins extend the entire length of the leaf with little or no cross-linking
- **Pinnate-** ريشي leaves have one major vein from which others branch
- **Palmate-** راحي leaves have several veins which branch

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

## Dicot and Monocot Leaves



**Reticulate**  
شبكة



**Parallel**  
متوازي



# Leaf Adaptations/ Modifications

Some plant species have evolved modified leaves to serve various functions.

(a) **Tendrils.** The tendrils by which this pea plant clings to a support are modified leaves. After it has “lassoed” a support, a tendril forms a coil that brings the plant closer to the support. Tendrils are typically modified leaves, but some tendrils are modified stems, as in grapevines.



(b) **Spines.** The spines of cacti, such as this prickly pear, are actually leaves, and photosynthesis is carried out mainly by the fleshy green stems.



(c) **Storage leaves.** Most succulents, such as this ice plant, have leaves modified for storing water.



(d) **Bracts.** Red parts of the poinsettia are often mistaken for petals but are actually modified leaves called bracts that surround a group of flowers. Such brightly colored leaves attract pollinators.



(e) **Reproductive leaves.** The leaves of some succulents, such as *Kalanchoe daigremontiana*, produce adventitious plantlets, which fall off the leaf and take root in the soil.





# Tendrils





# Spiny leaf- Cacti spines





# Succulent leaves




# Brightly-colored leaves- to attract pollinators






# Leaf Modifications

(d) Poinsettia leaves attract pollinators.




Copyright © 2008 Pearson Benjamin Cummings. All rights reserved.

(e) Pitcher plant leaves trap insects.



(f) Flowerpot plant leaves collect soil.



# Sharp things

- **Thorn** - sharp-pointed stem/shoot (fr. axillary bud)
- **Spine** - sharp-pointed leaf or leaf part  
leaf spine (also leaflet spine)  
stipular spine  
petiolar spine
- **Prickle** - sharp pointed epidermal appendage



Prickles

Thorns

Spines



# Leaf Structural/ Functional Types



**Tentacular Leaf - *Drosera* spp Sundew (Droseraceae)**



# Carnivorous plants

- **Insect-Trapping Leaves in areas with low soil N**
- **Insect digested by enzymes to release N from proteins**

**Trap Leaf - *Dionaea muscipula* Venus Fly Trap**



Copyright © McGraw-Hill Companies  
Permission  
Required for Reproduction or Display



# A Carnivorous Plant

Pitcher plant  
leaf



## **Reproductive Leaves - New plants at tips.**



# Sun & Shade Leaves

## Sun and Shade Leaves



**Shade Leaves**



**Sun Leaves**

**Grown in shade**

**Grown in sun**



# **Plant Tissues & organs**



# Plant Tissues

- A **tissue** is an organization of cells that work together as a functional unit.
- **Parenchyma** cells make up parenchyma tissue, which is a simple tissue.
- **Xylem** and **phloem** are complex tissues; they are composed of a number of different cell types (**Vascular tissue**).
- Tissues are grouped into **tissue systems** that extend throughout the body of the plant from to form the various **organs** of the plant.
- There are four plant tissue systems: *vascular, dermal, ground and meristematic tissue.*

# Tissue Systems

## Tissue Systems:

### **1. Ground tissue includes:**

- Parenchyma tissue
- Collenchyma tissue
- Sclerenchyma tissue

### **2. Vascular tissue includes:**

- Xylem tissue
- Phloem tissue

### **3. Dermal tissue:**

- Epidermis

### **4. Meristematic tissue**

# Plant Tissues

include

**Dermal tissue**

**Meristematic tissue**

**Vascular tissue**

**Ground tissue**

includes

includes

includes

**Epidermal cells**

**Xylem**

**Phloem**

**Parenchyma cells**

**Collenchyma cells**

**Schlerenchyma cells**

includes


includes

**Tracheids**

**Vessel elements**

**Sieve tube elements**

**Companion cells**



# Dermal Tissue:

- Covers the plant body and consists of epidermis in young plants & non-woody plants that is replaced later by **periderm** in woody plant
- **Epidermis** is made of parenchyma cells in a single layer
- Epidermis on stem and leaves prevents water loss by transpiration & produces a waxy material called **cuticle**





# Dermal Tissue:




Guard Cells  
surrounding stoma


- Openings in the epidermis on the underside of a leaf where gases are exchanged are called **stomata** (stoma, singular)
- **Sausage-shaped guard** cells are found on each side of the stoma to help open and close the pore to prevent water loss
- **Dead cork** cells replace epidermis in woody stems & roots

# Dermal Tissue:

Open:



Closed:



# Ground Tissue

- The ground tissue system makes up the rest of a plant and consists primarily of **parenchyma tissue**.
- Ground tissue functions primarily in storage, support, photosynthesis, and the production of defensive and attractant substances (oils and toxins).

# Vascular Tissue

- The vascular tissue system includes the **xylem and phloem**; it is the conductive or “plumbing” system of the plant.
- The **phloem** transports carbohydrates from sites of production (sources such as leaves) to sites of utilization for energy or where it is being stored (sinks) elsewhere in the plant.
- The **xylem** distributes water and mineral ions taken up by the roots to the stem and leaves.

# Meristems (Plant Stem Cells)

- **There are 3 main types of meristematic tissue in vascular seed plants ----  
apical, intercalary, & lateral meristems**




# Meristems (Plant Stem Cells)

- There are 3 main types of meristematic tissue in vascular seed plants ----  
**apical, intercalary, & lateral meristems**

**TABLE 31-2** *Types of Meristems*




Type	Location	Function
Apical meristem	tips of stems and roots	growth; increase length at tips
Intercalary meristem	between the tip and base of stems and leaves	growth; increase length between nodes
Lateral meristem	sides of stems and roots	growth; increase diameter

# Location of Meristematic Tissues




# Plant Cell Types (Support and Storage)

- **Parenchyma cells** are the most numerous type of cell in young plants.
- Parenchyma cells usually have thin walls and large central vacuoles.
- The photosynthetic cells in leaves or stems are parenchyma cells filled with chloroplasts. These cells are called **chlorenchyma** cells.
- Some parenchyma cells store lipids or starch (potatoes).
- Other parenchyma cells serve as “packing material” and play a vital role in supporting the stem especially in nonwoody stems.
- **Collenchyma cells** are supporting cells that lay down primary cell walls that are thick in the corners.
- Collenchyma cells provide support to leaf petioles, nonwoody stems, and growing organs.
- These cell types compose the cortex and pith tissues of the root and stems.








**(a) In leaves, parenchyma cells function in photosynthesis and gas exchange.**



**(b) In roots, parenchyma cells function in carbohydrate storage.**






**Parenchyma cells**

**Collenchyma cells**





LM 61x






# Plant Cell Types (Support)

- **Sclerenchyma cells** are the main supporting cells of a plant. They have a thick secondary cell wall that contains a substance called **lignin**, a component of wood. Therefore they are found in woody plants.
- There are **two types** of sclerenchyma cells: elongated fibers and variously shaped sclereids.
- **Fibers** often organize into bundles. (They are common components of xylem.)
- **Sclereids** may pack together very densely. (Sclereids are found in fruits such as pears and are what give them their gritty texture.) They are often referred to as “**stone cells**”.



**(a) Fibers**




**(b) Sclereids**





**Thick secondary cell walls**

# Sclerenchyma & Sclereids



**Sclerynchyma**

**Sclerynchyma  
Fibers**



**Sclereid  
Fibers**



# Plant Cell Types: Vascular Tissue (Transport)

## Xylem

- The xylem conducts water from roots to above ground plant parts. It contains conducting cells called *tracheary elements*.
- Tracheids are evolutionarily more ancient tracheary elements found in gymnosperms.
- Both tracheary elements: *Vessel elements* and *tracheids* undergo *apoptosis* (die) and do their jobs as empty cells (only the cell walls remain).
- *Vessel elements* are the water “*pipeline*” system in flowering plants, also formed from dead cells. Flowering plants have both tracheids and vessel elements.
- Vessel elements are generally larger in diameter than tracheids and are laid down end-to-end to form hollow tubes.




# Xylem

dead cells →  
water-conducting  
cells of xylem



tracheids

vessel  
elements



(b) Vessel elements  
with partially  
perforated end walls

(a) Tracheids



(c) Tracheids and vessels (colorized SEM)


# Plant Cell Types: Vascular Tissue (Transport)

## Phloem


- Cells of the phloem are alive when they do their job, unlike those of the xylem.
- The characteristic cell of the phloem is the *sieve tube member* in **flowering plants**.
- Cells of the phloem are arranged end-to-end and form long sieve tubes, which transport carbohydrates and other materials.
- The plasmodesmata in sieve tube members enlarge as they mature, resulting in end walls that look like sieves. called **sieve plate**
- At functional maturity, a sieve tube is filled with sieve tube sap (water, sugars, and other solutes).
- The sieve tube members have *adjacent companion cells*.
- *Companion cells* retain all their organelles and may regulate the performance of and support the sieve tube members.

# Phloem

## ■ sieve tube elements & companion cells




(a) Longitudinal view



(b) Transverse section (LM)

# Plant Organs






# **Structure of Primary Plant Organs**


# Root Tip Zones

- *The root tip*
  - covered by a root cap, which protects the delicate apical meristem as the root pushes through soil during primary growth
- *Zone of cell division*
  - Actively dividing, including root apical meristem, produces root cap cells
- *Zone of elongation*
  - Root cells elongate, pushes root tip further into soil
- *Zone of maturation*
  - Cells complete maturation, become fully functional




# Structure of the Root:

- Root cap
- Apical meristem
- Epidermis



# Structure of the Root:






# Structure of the Root:

- **Root cap** covers the apical meristem (growth tissue) at the tip of the root & produces a slimy substance so roots can more easily grow through the ground
- **Apical meristem** replaces cells of the root cap as they are damaged
- **Epidermis** covers the outside of the root & has extensions called root hairs that absorb water & minerals and increase the surface area of the root

# Structure of the Root:

- The **core** of the root is called the vascular cylinder, contains **xylem & phloem**
- A band of ground tissue called **cortex** surrounds the vascular cylinder
- A single cell layer called **endodermis** separates the cortex & vascular tissue
- Endodermal cells are coated with a waxy layer called the **Casparian strip** so water is channeled into the vascular tissue
- The **Pericycle** is the outermost layer of central vascular tissue & forms lateral roots

# Root Structure (Monocot vs Eudicot)






**Key  
to labels**

Dermal


Ground

Vascular



(a) Root with xylem and phloem in the center (typical of eudicots)

(b) Root with parenchyma in the center (typical of monocots)




**Endodermis**

**Pericycle**

**Xylem**

**Phloem**






# ***Stem Structure & Function:***

- Adapted to **support** leaves
- **Transport** water & minerals
- **Transport** sugars (usually sucrose) from Source (where they're made) to Sink (where they're stored)
- Movement of sugars is called **translocation**
- **Store** food and/or water
- Tubers (potatoes) underground **food storage** stems
- Stems grow from the **tip** or **apical meristem**
- Stems increase in circumference by **lateral meristems**
- Leaves are attached to stems at **nodes** & have lateral buds that can develop into new stems or branches
- **Internode** is space between nodes on a stem

- The tip of each stem usually has a Terminal Bud enclosed by specialized leaves called **Bud Scales**
- Vascular Tissue is arranged in **bundles** with xylem toward the inside & phloem toward the outside
- Vascular bundles are **scattered** throughout monocot stems
- Vascular bundles are arranged in **rings** in dicot stems


# Stem Structure

## Dicot (woody plants)



(a) Dicot

## Monocot (grasses)



(b) Monocot

### Key

- Dermal
- Ground
- Vascular



# Leaf Structure

- Leaf anatomy is adapted to carry out **photosynthesis**, limit evaporative water loss, and transport the products of photosynthesis to the rest of the plant.
- The two zones in leaf parenchyma that photosynthesize are the **palisade mesophyll** and the **spongy mesophyll**.
- Within the mesophyll is **air space** through which CO<sub>2</sub> can diffuse to the photosynthesizing cells.
- **Veins** (vascular bundles) supply mesophyll cells with water and minerals, and they **transport** the products of photosynthesis to the rest of the plant.
- The epidermis of the leaf is the outermost cell layer, which is covered by a **waxy cuticle**. The epidermis functions to keep water and photosynthetic products in the leaf.
- Guard cells allow controlled gas exchange through pores in the leaf (**the stomata**).






# Stomata:

- Openings called stomata on the underside of leaves for **gas exchange** ( $\text{CO}_2$  &  $\text{O}_2$ )
- Two **guard cells** on either side of the stomata open & close the openings
- When guard cells **LOSE** water, the stoma **CLOSE**, while the stoma **OPEN** when guard cells **GAIN** water & swell




# Leaf Structure



**Figure 35.17a-c**


**(b) Surface view of a spiderwort (*Tradescantia*) leaf (LM)**



**(c) Transverse section of a lilac (*Syringa*) leaf (LM)**



# Plant Secondary Growth




**These tissues form a vascular bundle. In woody plants they grow together and fuse to form a continuous ring creating annual growth rings.**

# Mature Dicot Stem

- Secondary growth increases the diameter of stems and roots.
- Secondary growth results from the activity of vascular and cork cambium.
- Vascular rays connect storage parenchyma to the sieve tubes of the phloem.
- Only eudicots have a vascular cambium and a cork cambium and thus undergo secondary growth.
- Cross sections of most tree trunks in temperate zone forests have annual rings.
- Annual rings form due to differential rates of growth in spring (when water is plentiful) and in summer.
- Wood that is no longer conducting water is known as heartwood.
- Sapwood is wood that is actively conducting water and minerals in the tree.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

## Continued Secondary Growth









Monocot

Dicot


Stem



Root



Leaf



# Monocots vs. Dicots

## Monocots



One  
cotyledon



Veins  
usually  
parallel



Vascular bundles  
usually complexly  
arranged



Fibrous  
root  
system



Floral parts  
usually in  
multiples  
of three

Embryos

Leaf  
venation

Stems

Roots

Flowers

## Dicots



Two  
cotyledons



Veins  
usually  
netlike



Vascular bundles  
usually arranged  
in ring



Taproot  
usually  
present



Floral parts  
usually in  
multiples of  
four or five

<b>Plant tissue</b>	<b>Cell type</b>	<b>Site</b>
<b>Epidermis</b>	<b>Ground cells; guard cells; cells forming trichomes; Sclerenchyma cells</b>	<b>Outer layer of plant body</b>
<b>Periderm</b>	<b>Cork cells; cork cambium cells; parenchyma cells of phelloderm; Sclerenchyma cells</b>	<b>Outer layer; mainly stems</b>
<b>Xylem</b>	<b>Tracheids; vessel members; Sclerenchyma cells; Parenchyma cells</b>	<b>All organs</b>
<b>Phloem</b>	<b>Sieve cells or sieve-tube members; albuminous cells or companion cells; other parenchyma cells; Sclerenchyma cells</b>	<b>All organs</b>
<b>Parenchyma</b>	<b>Parenchyma cells</b>	<b>Cortex, pith</b>
<b>Collenchyma</b>	<b>Collenchyma cells</b>	<b>Mainly stems</b>
<b>Sclerenchyma</b>	<b>Sclerenchyma cells</b>	<b>All organs</b>


# **Plant Physiology**



# Levels of Tissue Organization

- **cell** - unit of structure of all live organisms
- **tissue** - composed of groups of similar cells
- **organs** - composed of groups of tissues functioning together
- **organ systems** - composed of groups of organs functioning together


# Cell wall and Turgor



- When comparing two solutions there are three possible relationships, We identify the relationships by determining what would happen if a cell were placed in the solution.
- **Hypotonic**- A solution that causes a cell to **swell** because of osmosis meaning water rushes into the cell.
- **Isotonic**- A solution that causes **no change** in cell size. Meaning there is no movement of water.
- **Hypertonic**- A solution that causes a cell to **shrink** because of osmosis. Meaning water leaves the cell.

# Cell wall and Turgor

- Cell walls is made of neutral and charged polysaccharides  $\longrightarrow$  absorbs  $H_2O$  because it has a **hypotonic** environment.
- Increased  $H_2O$  in the cell  $\longrightarrow$  Turgor Pressure
- If a plant cell is turgid, It is very firm, a **healthy state** in most plants
- If a plant cell is flaccid, It is in an **isotonic** or **hypertonic** environment





# PROCESSES OF THE PLASMA MEMBRANE


- There are two types of passive transport: Diffusion and Osmosis
- The goal of both diffusion and osmosis is to reach **EQUILIBRIUM** within the cell
- **Equilibrium** is a condition in which the movement in one direction is equal to the movement in another direction

# Diffusion

- the tendency of molecules to move from an area of **higher concentration** to an area of **lower concentration**
- A difference in concentration between two adjacent regions is called a **concentration gradient**.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.


# Diffusion



# Osmosis

- movement of water through a membrane from a region of higher to lower concentration
- **Solute** - substance being dissolved in a liquid (e.g. salt)
- **Solvent** - substance doing the dissolving (e.g. water)
- **Permeability** - the extent to which a membrane will allow particular sized molecules to pass
- **Semi-permeable membrane** (selectively permeable)- allows some molecules to pass but not others





# Types of transport in Cells

- **Passive transport** - movement of substances through a membrane from a region of **high** to a region of **low** concentration - **no energy needed (ATP)** - diffusion and **osmosis** are examples of this
- **Active transport** - movement of substances through a membrane from a region of **low** concentration to a region of **high** concentration - **requires cellular energy (ATP)**

- Whether **passive** or **active** transport is needed depends on the **CONCENTRATION GRADIENT**
- The concentration gradient is the **difference** in the concentration of a substance in two different spaces
- **Concentration** - the amount of a particular substance in a contained area compared with the amount of the same substance in another area

# Active Transport

- The movement of a substance against the concentration gradient. (uphill)
- Active transport requires cell to **USE ENERGY**
- **Sodium pump** - transports three sodium ions out of the cell and two potassium ions into the cell
- Both are against the concentration gradient
- The energy needed to perform this activity is supplied by ATP (adenosine tri-phosphate)
- ATP is a unit of energy made by the cell



# Active and Passive Transport

extracellular fluid

*Passive*

$O_2$

$Na^+$

diffusion through membrane

aqueous channel

$O_2$

$Na^+$

Passive transport proceeds as diffusion from areas of high concentration to areas of low concentration

cytoplasm

**Active**

$Na^+$

$K^+$

sodium potassium pump

ATP

ADP +  $P_i$

$Na^+$

$K^+$


Active transport proceeds against the concentration gradient, and requires an input of energy

# Photosynthesis

- the conversion of light energy to chemical energy by chlorophyll in chloroplasts
- Overall Net equation for photosynthesis:
  - 6 Water + 6 Carbon dioxide yields glucose + 6 oxygen (when catalyzed by chlorophyll in the presence of sunlight)
$$6\text{CO}_2 + 6\text{H}_2\text{O} = \text{C}_6\text{H}_{12}\text{O}_6 + 6\text{O}_2$$

## Respiration






# Plant Water Relations

# Why is water important to (plant) cells?

- **Water constitutes about 70% by weight of annual plants**
- **Water has multiple roles in plant cells**
  1. **Thermal property: a liquid!**
    - **High heat potential: can absorb energy changes without large temperature changes (slows heating and cooling)**
  2. **“Universal” solvent required for mineral uptake and transport**
  3. **It is a requirement for biochemical reactions to proceed**
    - **Most enzymes are water soluble**



# How plants get their food ?





# Photosynthesis:

Green plants take in **carbon dioxide** ( $\text{CO}_2$ ) from the air


They take up **water** ( $\text{H}_2\text{O}$ ) from the soil

The plants combine the  $\text{CO}_2$  with the  $\text{H}_2\text{O}$  to make the sugar (Glucose ( $\text{C}_6\text{H}_{12}\text{O}_6$ ))


- **Photosynthesis** is a chemical process that energy from light is harvested to provide carbohydrates.
- It is the major path through which carbon re-enters the biosphere (from  $\text{CO}_2$ ).
- **Photosynthesis** is also the major source of oxygen ( $\text{O}_2$ ) in the earth's atmosphere



# All the reactions to combine $\text{CO}_2$ and $\text{H}_2\text{O}$ take place in the chloroplast







e.g. seed germination




fruits

other sugars


**GLUCOSE**

protein



cytoplasm


starch




storage e.g. starch in potato


cellulose

cell walls



# Visible Light Passing Through a Prism






# Capturing Energy

## أقتناص الطاقة

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

### Leaf— Levels of Organization (1)





# Types of photosynthesis

- **C3**

- The majority of plants In the case of C3 photosynthesis, the first organic product of carbon fixation is a three-carbon compound (3-phosphoglycerate), which is the reason these plants are termed the C3 plants.

- **C4**

- CO<sub>2</sub> temporarily stored as 4-C organic acids resulting in more efficient C exchange rate
- Advantage in high light, high temperature, low CO<sub>2</sub>
- Many grasses and crops (e.g., corn, sorghum, millet, sugar cane)

- **CAM**

- Stomata open during night
- Advantage in arid climates
- Many succulents (e.g. cacti, euphorbs, bromeliades, agaves)

# **TO SUM UP .....**

**Plants combine carbon dioxide from the air, and water from the soil to make glucose.**

**The energy needed for this process comes from sunlight**

**The sunlight is absorbed by chlorophyll contained in the chloroplasts of the leaf.**

**The glucose can be used for energy or to make other substances.**


**To make other substances, the glucose must be combined with other chemical elements such as nitrogen and potassium.**

**These chemical elements are present as ions in the soil and are taken up in solution by the roots.**

# **Classification of Organisms**

# Classification of Organisms

## Taxonomic Kingdoms of Life



- The most basic category of organisms is called a **kingdom**.
- Most scientists divide organisms into **five** major kingdoms.

# Systematics

1. **Species:** Organisms sharing a set of biological traits and reproducing only their exact kind.  
(Species is the fundamental unit in taxonomy)
  - a. **strains:** organisms within the species varying in a given quality
  - b. **types:** organisms within the species varying immunologically.
2. **Genus:** closely related species
3. **Family :** closely related genera
4. **Order:** closely related families
5. **Class :** closely related order
6. **Phylum:** related classes



# **Survey of Microorganisms**

- 1. Viruses**
- 2. Bacteria**
- 3. Cyanobacteria**
- 4. Algae**
- 5. Fungi**
- 6. protozoa**


# **VIRUSES**

- **Obligated intracellular parasite.**
- **host specific:**
  - **bacteriophage**
  - **animal virus**
  - **plant virus**
- **according to its genetic material**
  - **DNA virus**
  - **RNA virus**
- **Shape:**


**Most common shape is icosahedral , some are helical shape**
- **Structure:**

**Protein capsid and genetic material some animal virus have envelope with glycoprotein spikes**
- **Life cycle: lytic infection lysogenic infection**
- **Some animal viruses are closely associated with certain cancers**

# Virus structure



(a) A T-even bacteriophage



# Biological Kingdoms

## 2 Kingdoms

Traditional view



A diagram illustrating the traditional two-kingdom view of biological classification. It consists of two adjacent squares. The left square is green and contains the word 'plants'. The right square is orange and contains the word 'animals'.

plants


animals



# Biological Kingdoms

## 5 Kingdoms

Whittaker, 1969



# Biological Kingdoms

Five kingdom system:

Monera

Protista

Plantae

Fungi

Animalia

Six kingdom system:

Eu-  
bacteria

Archae-  
bacteria

Protista

Plantae

Fungi

Animalia

Three domain system:

Eu-  
bacteria

Archae-  
bacteria

E U K A R Y A

Eight kingdom system:

Eu-  
bacteria

Archae-  
bacteria

Archezoa

Chromista

Protista

Plantae

Fungi

Animalia

# **Six-Kingdom System**

**Kingdom Archaeobacteria**

**Kingdom Eubacteria**

**Kingdom Fungi**


**Kingdom Protista**

**Kingdom Plantae**

**Kingdom Animalia**


# KINGDOM:ARCHAEBACTERIA

- They Are **UNICELLULAR PROKARYOTES** with distinctive Cell Membranes as well as Biochemical and Genetic Properties that differ from ALL other kinds of life.
- Some are autotrophic, producing food by chemosynthesis. Includes Chemosynthetic Bacteria
- Most are heterotrophic.
- Many Archaeobacteria live in **harsh environments** such as Sulfurous Hot Springs, very salty lakes, and in anaerobic environments, such as the intestines of mammals.




# KINGDOM EUBACTERIA

- They are **UNICELLULAR PROKARYOTES**.
- Most of the Bacteria (Germs) that affect your life are members of the Kingdom Eubacteria.
- Eubacteria are both **autotrophic** and **heterotrophic**.
- Includes the disease-causing bacteria such as tooth decay or food poisoning.
- The Combined Kingdoms, Archaeobacteria and Eubacteria include the **greatest number** of living things on Earth.
- All of the prokaryotes are in these two kingdoms (**Archaeobacteria and Eubacteria**).
- Both reproduce by **binary fission**, but they do have some ways to recombine genes, allowing evolution to occur.



*Escherichia coli*




*Staphylococcus*




# KINGDOM PROTISTA

- Kingdom Protista contains all eukaryotes that are **NOT** Plants, Animal, or Fungi, more than 50,000 species in all.
- Kingdom Protista includes unicellular and a few simple multicellular **EUKARYOTES**.
- The cells of multicellular protists are not specialized to perform specific functions in the organisms.



Amoeba



**Pediatrum**

# KINGDOM FUNGI

- Fungi are **eukaryotes**, and most are **multicellular**.
- The cells of fungi have cell walls that contain a material called **chitin**.
- These organisms are heterotrophic and obtain their nutrients by releasing digestive enzymes into a food source.
- Fungi act either as **decomposers** or as **parasites** in nature.



mushroom

# KINGDOM PLANTAE

- Plants are **eukaryotic**, **multicellular** and carry out **photosynthesis**. They are autotrophs.
- The cells of plants have cell walls, that contain the **polysaccharide cellulose**.
- Plant cells are specialized for different functions.
- Kingdom Plantae includes mosses, ferns, cone-bearing plants (gymnosperms), and flowering plants (angiosperms).



fern



Flowering plant

# KINGDOM ANIMALIA

- Animals are **multicellular**, **eukaryotic**, and **heterotrophic**.
- Animal cells have **NO** CELL WALLS.
- Most members of the Animal Kingdom can move from place to place.
- Fish, Birds, Reptiles, Amphibians, and mammals-including humans belong to the Kingdom Animalia.
- This Kingdom also includes sponges, jellyfish, worms, sea stars, and insects.



jellyfish



elephant



# Six-Kingdom System

**TABLE 18-2** *Six Kingdoms of Life*

<u>Kingdom</u>	<u>Cell type</u>	<u>Number of cells</u>	<u>Nutrition</u>
Archaeobacteria	prokaryotic	unicellular	autotrophy and heterotrophy
Eubacteria	prokaryotic	unicellular	autotrophy and heterotrophy
Protista	eukaryotic	unicellular and multicellular	autotrophy and heterotrophy
Fungi	eukaryotic	unicellular and multicellular	heterotrophy
Plantae	eukaryotic	multicellular	autotrophy and (rarely) heterotrophy
Animalia	eukaryotic	multicellular	heterotrophy

prokaryotic  
eukaryotic


unicellular  
multicellular

autotrophic  
heterotrophic



عدد الانواع	Group
٥,٠٠٠	بكتيريا
٤٠,٠٠٠	طحالب
٤٠,٠٠٠	احياء وحيدة الخلية
٣٠٣,٠٠٠	لا فقاريات (دون الحشرات)
٨٤٠,٠٠٠	لافقاريات (حشرات)
٤٦,٥٠٠	فقاريات
٧٠,٠٠٠	فطريات
٢٥٦,٠٠٠	نباتات متطورة
١,٦٠٠,٥٠٠	المجموع

Category	Species	Totals
<b>Vertebrate Animals</b>		
Mammals	5,513	
Birds	10,425	
Reptiles	10,038	
Amphibians	7,302	
Fishes	32,900	
<b>Total Vertebrates</b>		<b>66,178</b>
<b>Invertebrate Animals</b>		
Insects	1,000,000	
Spiders and scorpions	102,248	
Molluscs	85,000	
Crustaceans	47,000	
Corals	2,175	
Others	68,827	
<b>Total Invertebrates</b>		<b>1,305,250</b>
<b>Plants</b>		
Flowering plants (angiosperms)	268,000	
Conifers (gymnosperms)	1,052	
Ferns and horsetails	12,000	
Mosses	16,236	
Red and green algae	10,386	
<b>Total Plants</b>		<b>307,674</b>
<b>Others</b>		
Lichens	17,000	
Mushrooms	31,496	
Brown algae	3,127	
<b>Total Others</b>		<b>51,623</b>
<b>TOTAL SPECIES</b>		<b>1,730,725</b>



# BACTERIA

## ❑ Typical prokaryotes.

- Three shapes: cocci, bacilli and spiral
- Can be autotroph or heterotroph

Autotroph: photoautotroph or chemoautotroph


## ❑ Heterotroph: parasite or saprophyte

- Type of reproduction: binary fission
- Some genetic material transfer:  
transformation, transduction and conjugation


# Bacterial morphology



Coccus




Rod



Spirillum




Spirochete



Stalk

Hypha

Budding and appendaged bacteria



Filamentous



# Algae

1. **Euglenoids**  
eg. *Euglena*  
**food storage** - lipoid polysaccharide - paramylum
2. **Green algae**  
eg. *Chlamydomonas*  
**food storage** - starch
3. **Golden Brown algae**  
eg. *Diatoms*  
**food storage** – oil and leucosin (a polysaccharide)  
have fucoxanthin, a brownish pigment
4. **Brown Algae**  
Mainly marine water algae  
**food storage** – laminarin, a polysaccharide and mannitol, a sugar alcohol
5. **Fire Algae**  
**Dinoflagellate**  
eg. *Peridinium*  
**food storage** – starch, fat, oils

# Fungi

The fungi are **not true plants ...!!** you see they do not contain chlorophyll! Indeed many scientists today place fungi as more closely related to animals than plants (**chitin** – which also forms the arthropod exoskeleton - is the main component of fungal cell wall).

Fungi and animals are descended from a common ancestor: A unicellular eukaryote with a flagellum.

# Unicellular

Unicellular members of the zygomycetes, ascomycetes, and basidiomycetes.

**e.g. , Yeasts:**  
**Budding: mitosis**  
**followed by**  
**asymmetrical**  
**cell division.**




*Saccharomyces* sp.

# Multicellular

## Multicellular fungi:

- Body is a **mycelium**—composed of tubular filaments called **hyphae**. (singular hypha)
- Hyphae cell walls have **chitin**.
- Some hyphae have incomplete cross walls or **septa**, and are called **septate**.
- Hyphae without septa are called **coenocytic**.

# Most Hyphae Are Incompletely Divided into Separate Cells




**LIFE 8e, Figure 30.4**



# Lichens: Nature's perfect marriage

**Lichens: fungus + photosynthetic organism**

- Fungi—mostly ascomycetes
- Photosynthetic partner—cyanobacterium or alga, or both.
- Species are named for fungal component.
- Can survive the harshest environments on Earth.
- Very sensitive to toxic compounds—good indicators of air pollution.
- Lichens are great ecological markers
- Early medicinal remedy



# **Plant systematics**

# WHAT'S THE PLANT ?

- Plants are members of the kingdom plantae. Plants are photosynthetic multicellular eukaryotes  
**PHOTOAUTOTROPHS.**
- Cell walls are made of **CELLULOSE** - the material that bacteria and protists in our small intestine digest for us. Cellulose is a kind of complex sugar or polysaccharide.
- Although cellulose plays an important role in structural support in the cell walls of plants, cellulose is found in other forms - such as cotton.
- The green of plants comes from their photosynthetic pigments (chlorophyll a & b)

# What are characteristics of plants?

- Sessile
- Multicellular & cell specialization
- Eukaryotic
- cellulose cell walls
- autotrophic (photosynthetic)
- Chlorophylls *a* and *b* in thylakoid membranes
- Store reserve food as amylose (starch)
- Alternation of Generations
  - Sporophyte
  - gametophyte

# What do plants need to survive?

- **Sunlight** - energy of sun captured by chlorophyll and used to join  $\text{CO}_2$  and  $\text{H}_2\text{O}$  to form glucose ( $\text{C}_6\text{H}_{12}\text{O}_6$ ); **plants need broad leaves to maximize light absorption**
- **Water and minerals** - **roots** to absorb these
- **Gas Exchange** – **stomata** in leaves
- **Protective structures** - were required to protect the developing embryos.
- **Movement of water and nutrients**
  - Most plants have tubes – **phloem** (nutrients down) and **xylem** (water up)
  - Some small plants use **diffusion**



# Plant Adaptations to Land

## Problems:

- Need minerals
- Gravity
- Increase in Height for Light
- Adaptations for Drier environment
- Reproduction


## Solutions:

- Roots absorb  $H_2O$  & minerals
- Lignin & cellulose in cell walls
- Vascular Transport System
- Waxy cuticle & stomata with guard cells
- Pollen containing sperm

# GENERAL LIFE CYCLE OF PLANTS

- The life cycle of plants has two different phases. This is called; **ALTERATION OF GENERATIONS**. In this alteration of generations, the plant takes turns undergoing mitosis and meiosis to produce haploid (n) gametes and diploid (2n) phase.
- The diploid (2n) phase is called the **sporophyte** - or spore producing plant. The haploid (n) phase is called the **gametophyte** - or gamete producing plant.
- The **spores** are haploid (n) and produced through meiosis in the sporophyte plant - each spore can grow into a new plant; the gametophyte!
- A **gamete** is a reproductive cell produced by mitosis and fuses during fertilization with another gamete to produce a new individual - the diploid sporophyte.


# Alternation of Generations



- **Alternation of generations:**
  - **Sporophyte** –  $2n$  – Diploid – produce haploid spores by meiosis
  - **Gametophyte** - ( $1n$ ) – haploid undergoes mitosis to produce eggs and sperm – the eggs and sperm (gametes)
  - **Zygote** - merge to grow into a  $2n$  sporophyte (cycle continues)

# Plant Cladogram

## Relationships among the various groups of plants






# The four main groups of plants:

- **Mosses** (nonvascular plants)– Bryophytes (15,600 species)
- **Ferns** (vascular & seedless plants)– Pterophytes (11,000 species)
- **Gymnosperms** (vascular & naked seeds)- Cone-bearing Plants (760 species)
- **Angiosperms** (vascular & covered seeds)- Flowering Plants (235,000 species)

# Diversity of plants today



▲ **Figure 22-7** The great majority of plants alive today are angiosperms, which are also known as flowering plants. **Interpreting Graphics** What is the second largest group of plants?

# Plant kingdom

(Land Plants)



**Hepatophyta (Liverworts)**



**Anthocerotophyta (Hornworts)**



**Bryophyta Mosses**

**Nonvascular plants**



**Lycophytes (spike mosses)**

**Sphenophyta (horsetails)**

**Psilotophyta (whisk ferns)**

**Pterophyta (Ferns)**

**Seedless plants**

**Vascular plants**



**Gymnosperms**




**Angiosperms**

**Seed plants**

# Nonvascular Plants

- Do not have vascular tissue for support or conduction of materials
- Require a constantly moist environment
- Plants can't grow as tall



**Moss Gametophytes & Sporophytes**



# **Nonvascular Plants (contd.)**

- **Cells must be in direct contact with moisture**
- **Materials move by diffusion cell-to-cell**
- **Sperm must swim to egg through water droplets**



# Vascular Plants

## Vascular System:

- **Xylem** tissue carries **water** and **minerals** upward from the roots
- **Phloem** tissue carries **sugars** made by photosynthesis from the leaves to where they will be stored or used
- **Sap** is the fluid carried inside the xylem or phloem

# **Seed producing plants**

- **Major adaptations**
  - Pollen (male gametophyte)
  - Seeds (embryonic plant)(male and female gametophyte are greatly reduced in size)
- **Two types**
  - Gymnosperms (lack flowers, naked seeds)
  - Angiosperms (flowering plants, seeds enclosed in fruits)

# Types of Nonvascular Plants

- (1) Bryophyta**
- (2) Hepatophyta (liverwort)**
- (3) Anthoceroophyta (Hornworts)**

# Types of Nonvascular Plants

## (1) BRYOPHYTA



*Sphagnum*

**Sporophytes**

**Gametophyte**



## (2) Hepatophyta (liverwort)




**Sporophytes**


**Gametophyte**



### (3) AnthoceroPHYta (Hornworts)



# Simplified Lifecycle of a Bryophyte



# **Types of Seedless Vascular Plants**

---

- (1) Psilophyta (Whisk Ferns)**
- (2) Lycophyta (Spike Mosses)**
- (3) Sphenophyta (Horsetails)**
- (4) Pterophyta (True Ferns)**

# **Types of Seedless Vascular Plants**

## **(1) PSILOPHYTA (whisk ferns)**



- **Epiphytes**
- **Rootless and leafless**



## **(2) LYCOPHYTA (spike mosses)**




- **Produces a sporangia bearing strobilus.**



# (3) SPHENOPHYTA (Horsetails)



- Genus: *Equisetum*
- Large deposits of silica in their leaves.
- Jointed stems with whorled leaf arrangement.





## **(4) PTEROPHYTA (True ferns)**



# PTEROPHYTA (True ferns)

- Broad leaves called **fronds**
- Leaflets called **pinnae**
- Sporangium (**sorus p/sori**) are formed on the underside of the fronds.
- Unfolding fronds are called **fiddleheads**.
- Spores are dispersed by the wind.

# Seedless Vascular Plants (reproduce via spores)

**TABLE 30-2** *Seedless Vascular Plants*

Example plant	Phylum	Features	Size	Location
Whisk ferns	Psilotophyta	<ul style="list-style-type: none"> <li>• produce reproductive structures on the ends of forked branches</li> <li>• no roots or leaves</li> </ul>	<ul style="list-style-type: none"> <li>• about 30 cm (1 ft) tall</li> </ul>	<ul style="list-style-type: none"> <li>• tropical and temperate regions, as far north as South Carolina</li> </ul>
Club mosses	Lycophyta	<ul style="list-style-type: none"> <li>• evergreens that produce spores in cones</li> <li>• have roots</li> </ul>	<ul style="list-style-type: none"> <li>• about 5 cm (2 in.) tall</li> </ul>	<ul style="list-style-type: none"> <li>• tropical and temperate regions, on forest floors, in swamps, or as epiphytes</li> </ul>
Horsetails	Sphenophyta	<ul style="list-style-type: none"> <li>• jointed stems</li> <li>• outer cells of stems contain silica, the major component of sand</li> </ul>	<ul style="list-style-type: none"> <li>• about 60–90 cm (2–3 ft) tall</li> </ul>	<ul style="list-style-type: none"> <li>• tropical and temperate regions, usually in moist soil</li> </ul>
Ferns	Pterophyta	<ul style="list-style-type: none"> <li>• leaves</li> <li>• most have an underground stem</li> <li>• most produce spores on the underside of their leaves</li> </ul>	<ul style="list-style-type: none"> <li>• range from less than 1 cm (0.4 in.) to 25 m (82 ft) tall</li> </ul>	<ul style="list-style-type: none"> <li>• all climates, on forest floors, as epiphytes, some in full sun, some aquatic</li> </ul>

# **Types of Gymnosperms**



# Plant kingdom

(Land Plants)



**Hepatophyta (Liverworts)**



**Anthocerotophyta (Hornworts)**



**Bryophyta Mosses**

**Nonvascular plants**



**Lycophytes (spike mosses)**

**Sphenophyta (horsetails)**

**Psilotophyta (whisk ferns)**

**Pterophyta (Ferns)**

**Seedless plants**

**Vascular plants**



**Gymnosperms**



**Angiosperms**

**Seed plants**

# **(1) CYCADOPHYTA**

## **(Cycads)**

- **Large fernlike leaves**
- **Plants are either males or females**
- **Plants produce gametes in large strobilus**





## (2) GINKGOPHYTA (*Ginkgo biloba*)





## (2) GINKGOPHYTA (*Ginkgo biloba*)


- Only one species in present day
- Bear male and female cones on separate plants.
- Male produces pollen in strobilus-like cones.
- Female bears seed which develop a fleshy outer covering
- Plants are resistant to air pollution



# **(3) CONIFEROPHYTA**

## **(Cone-bearing trees)**

- **Cone bearing**
- **needles with thick waxy covering**
- **Stomata in cavities below surface**
- **Seeds are carried on the surfaces of cones**
- **Evergreen and adapted to cold and dry habitats**




**Male & female cones**





# What is alternation of generations in gymnosperms?



*Gnetum*



## (4) GNETOPHYTA

There are three genera:

1. *Gnetum*: A tropical climbing plant
2. *Ephedra* (Shrub-like plants)
3. *Welwitschia* (Desert dweller with large tuberous root). Has only two leaves and may live 1000 yr.

*Ephedra*



*Welwitschia*





# **Angiospermae**


## **(Anthophyta – flowering plants)**



# Key


➡ Haploid ( $n$ )

➡ Diploid ( $2n$ )



# What are the ANTHOPHYTA?


- Flowering
- Seeds are within a layer of protective tissue
- Flowers, ovaries, pollinators (insects, etc.)





# Inflorescence types

- An inflorescence is an arrangement of one or more flowers on a floral axis



# **Inflorescence types**

- **Inflorescence type determined by:**
  - **Number of flowers**
  - **Positional relationships**
  - **Degree of the development of their pedicels**
  - **Nature of their branching pattern**

# Simple Inflorescences

- **Terminal: flower at the tip of a stem**



Scarlet rose-mallow (*Hibiscus coccineus*)

# Compound Inflorescences

- Two or more flowers per inflorescence





# Compound Inflorescences


- **Spike:** elongate inflorescence; flowers are sessile, dense, or remote from one another



Spiked blazing star (*Liatris spicata*)


# Compound Inflorescences

- **Catkin:** A spikelike inflorescence of unisexual flowers; found only in woody plants.



# Compound Inflorescences


- **Raceme**: an elongate inflorescence of pedicellate flowers on an unbranched rachis





# Compound Inflorescences


- **Umbel:** a flat-topped or somewhat rounded inflorescence in which all of the pedicels arise from a common point at the tip of the peduncle






# Compound Inflorescences

- **Corymb**: a flat-topped or somewhat rounded inflorescence in which the pedicels of varying length are inserted along the rachis



# Compound Inflorescences

- **Panicle:** a much-branched inflorescence with a central rachis which bears branches which are themselves branched



panicle  
(compound raceme)



# Compound Inflorescences











- **Head, (Capitulum) :**  
is a short dense spike in which the flowers are borne directly on a broad, flat peduncle, giving the inflorescence the appearance of a single flower.




# **Dicots v Monocots – other differences**




## Characteristics of Monocots and Dicots

	Monocots	Dicots
Seeds	Single cotyledon 	Two cotyledons 
Leaves	Parallel veins 	Branched veins 
Flowers	Floral parts often in multiples of 3 	Floral parts often in multiples of 4 or 5 
Stems	Vascular bundles scattered throughout stem 	Vascular bundles arranged in a ring 
Roots	Fibrous roots 	Taproot 

# Switch to Sporophyte Dominance





# Plant Ecology



# Ecology


The word first coined by E. Haeckel (1869) and H. Reiter (1885) from the Greek words **OIKOS** (habitat or home) and **LOGOS** (the study or knowledge).

**Ecology** is the study of living organisms in relation to their habits and habitats.


It seeks to explain how many different kinds of organisms can live together in the same place for many generations (share habitats).

It deals with the reciprocal relationships between organisms and their environment.

# The reciprocal relationships between organisms and their environment



**Odum (1971) and Clark (1973) have defined ecology as a science of ecosystems, i.e. study of the structure and function of nature.**



**The interplay of organisms and environment combines between objectivity of science and subjectivity of art.**

# The three fundamental ecological questions

➤ **What is there?**

➤ **How much is there?**

➤ **Why is it there?**



# Environment

Summation of all **biotic** (living) and **abiotic** (non-living) components that surround or potentially influence the organisms and their habitats.

It is a **complex of factors** acting, reacting and interacting with the organism complex, **i.e., organisms and their environment are wedded together in state of constant flux.**

- ❖ **Macro-environment** (prevailing regional climate)
- ❖ **Micro-environment** (close to an organism to be influenced by it)


# Biosphere (sphere of life)

Total portion of lithosphere, hydrosphere and atmosphere that supports the life of organisms.

مكونات المحيط الحيوي

Atmosphere

الغلاف الجوي



Hydrosphere

Lithosphere

# Levels of organization


The biological portion of an interactive organism-environment system.

In ecology, the levels of organization include:

- ❖ **Individuals** - the fundamental functional units.
- ❖ **Populations** - members of the same species co-occurring in space and time and sharing the same resources.
- ❖ **Guilds** - group of populations of different species exploit the same resources.
- ❖ **Communities** - populations of organisms living together in the same environment.
- ❖ **Ecosystems** – Dynamic systems of organisms interacting with each other and their environment.
- ❖ **Biomes** - regional ecosystem types with similar communities.

# Levels of organization

## مستويات التعضي





# Ecosystem

**All organisms and their physical environment in a single location.**

**Self-sustaining and self-regulating communities of organisms interacting with one another and with their environment.**

**The matter that cycles into and out of the ecosystem is small compared with the quantities that are internally recycled in a continuous exchange of the essentials of life.**

**Various ecosystems make up the largest life unit called biosphere.**

# Ecosystem structure

## 1. Abiotic components

- **Energy** - solar energy
- **Physical factors** - temperature, light, wind, etc.
- **Chemicals** - inorganic substances (oxygen, carbon, etc.) and organic substances (carbohydrates, proteins, etc.)

## 2. Biotic components

- **Producers** - green plants (autotrophs)
- **Consumers** - animals (heterotrophs)
  - Herbivores** (primary consumers)
  - Carnivores** (primary, secondary, tertiary, etc. consumers)
  - Omnivores** (generalists)-can feed on both plants and animals.
  - Scavengers** (top carnivores)-utilize the dead remains of animals

## 3. Decomposers (saprotrophs)

bacteria and fungi


# Food chain

**A linear energy and chemical flow through organisms**

**i.e. food from one trophic (feeding) level reaches to the other trophic level (who eats whom?).**

**In the classical food chain, plants are eaten only by primary consumers, primary consumers are eaten by secondary consumers, secondary consumers are eaten by tertiary consumers, and so forth.**

# Food Chain



# **Food web**


**The actual pattern of food consumption in a natural ecosystem.**

**A given organism may obtain nourishment from many different trophic levels and thus gives rise to a complex, interwoven (interconnected) series of energy transfers.**

**The more complex the food web, the more stable the ecosystem.**



# Food web



# Food chain and pollutant accumulation



# Types of Systems


- (1) **Closed system:** **Energy** but not matter is exchanged between the system and environment, e.g., earth.
- (2) **Open system:** Both **energy** and matter are exchanged between the system and environment, e.g., lakes or living organisms.
- (3) **Isolated system:** Neither **energy** nor matter are exchanged between the system and environment, e.g., (examples and comments given as student assignment).

# Zonobiomes (Biomes)

- |                                 |   |
|---------------------------------|---|
| 1. Equatorial diurnal climate   | المناطق المدارية اليومية                |
| 2. Tropical                     | المناطق الاستوائية                      |
| 3. Subtropical (Desert)         | المناطق تحت الاستوائية (الصحراء)        |
| 4. Mediterranean                | مناطق البحر المتوسط                     |
| 5. Warm temperate               | المناطق المعتدلة الدافئة                |
| 6. Temperate                    | المناطق المعتدلة                        |
| 7. Arid temperate (Continental) | المناطق المعتدلة الجافة (المناخ القاري) |
| 8. Cold temperate               | المناطق المعتدلة الباردة                |
| 9. Arctic (Tundra)              | المناطق القطبية (التندرا)               |



# Zonobiomes



Tundra

Coniferous forests  
(Montane and Taiga)

Deciduous forests

Grasslands

Shrublands (chaparral,  
mediterranean, and tropical)

Deserts

Tropical rain forests

Semidesert (including tropical thornwoods)  
arid grasslands

Mountains with mixed biomes

Tropical savannas



# Allelopathy

- ❖ **Allelopathy** is a **chemical interaction** between two or more populations that suppresses one population while the another remains stable through the release of metabolic by-products (**allelochemicals**) into the environment.
- ❖ The allelochemicals are selectively toxic, affecting some species but not the others.
- ❖ Many allelopathic species release **autotoxic** compounds which affect their own growth and development negatively (**Autopathy**).
- ❖ To gain a selective advantage, there must be a trade-offs between allelopathy and autopathy through which the source (donor) species inhibits the target (receptor) species more than its self-inhibition.

- ❖ **Amensalism:** An interaction between two populations in which one **is not** affected and the other is **negatively** affected.
- ❖ **Commensalism:** An interaction between two populations in which one **is not** affected and the other is **positively** affected.
- ❖ **Predation:** An interaction in which one living organism serves as a food source for another organism; one **positively** (predator) affected and the other is **negatively** (prey) affected.

- ❖ **Parasitism:** An interaction in which an organism serves as a food source (**host**) for other organisms (**parasites**) that commonly are much smaller in size, ultimately results in the death of, or detriment to, the host organism.
- ❖ **Symbiosis:** An interaction in which **both** species are **positively** influenced as a result of their **co-occurrence** (**happy and strong together**).
- ❖ **Mutualism:** An interaction in which each member derives a positive benefit and also provides a portion of the cost of the interaction.

**THE END ...**