# **CEREBRAL BLOOD CIRCULATION**



Khaleel Alyahya, PhD, MEd King Saud University School of Medicine @khaleelya

# **OBJECTIVES**

#### At the end of the lecture, students should be able to:

- List the cerebral arteries.
- Describe the cerebral arterial supply regarding the origin, distribution and branches.
- $\circ~$  Describe the arterial Circle of Willis .
- Describe the cerebral venous drainage and its termination.
- Describe arterial & venous vascular disorders and their clinical manifestations.





## **Review: THE BRAIN**

- ✓ Large mass of nervous tissue located in cranial cavity.
- ✓ Has four major regions.



# **Review: CEREBRUM**

- The largest part of the brain, and has two hemispheres.
- The surface shows elevations called gyri, separated by depressions called sulci.
- Each hemispheres divided into four lobes by deeper grooves.
- Lobs are separated by deep grooves called fissures.



## **Review: BLOOD VESSELS**

- Blood vessels are the part of the circulatory system that transports blood throughout the human body.
- There are three major types of blood vessels:
  - Arteries, which carry the blood away from the heart.
  - Capillaries, which enable the actual exchange of water and chemicals between the blood and the tissues.
  - Veins, which carry blood from the capillaries back toward the heart.
- The word vascular, meaning relating to the blood vessels, is derived from the Latin vas, meaning vessel.
  - Avascular refers to being without (blood) vessels.



# **Review: HISTOLOGY**

- The arteries and veins have three layers, but the middle layer is thicker in the arteries than it is in the veins:
  - Tunica Intima (the thinnest layer): a single layer of simple squamous endothelial cells.
  - Tunica Media (the thickest layer in arteries): is made up of smooth muscle cells and elastic tissue.
  - Tunica Adventitia (the thickest layer in veins) entirely made of connective tissue.
- Capillaries consist of little more than a layer of endothelium and occasional connective tissue.



# **Review: BLOOD**

- Blood is the actual carrier of the oxygen and nutrients into arteries.
- Blood is made mostly of plasma, which is a yellowish liquid that is 90% water.
- Plasma contains also salts, glucose and other substances.
- Most important, plasma contains proteins that carry important nutrients to the body's cells and strengthen the body's immune system.
- Blood has main 3 types of blood cells that circulate with the plasma.



## **CEREBRAL CIRCULATION**

- The movement of blood through the network of blood vessels to supply the brain.
- The arteries carry oxygenated blood and other nutrients to the brain.
- The veins carry deoxygenated blood back to the heart removing carbon dioxide and other metabolic products.
- The movement of blood in the cerebral circulation is called cerebral blood flow.



# **CEREBRAL ARTERIAL SUPPLY**

- The arterial cerebral circulation is divided into anterior and posterior cerebral circulations.
- The anterior and posterior cerebral circulations are interconnected via bilateral posterior communicating arteries.
  - Posterior communicating arteries are part of Circle of Willis.
    - $\checkmark$  Located on the base of the brain.
    - ✓ It Encircles:
      - Optic chiasma
      - Hypothalamus
      - Midbrain
- The cerebral arterial supply is provided by two systems:
  - Carotid System
    - Supply anterior portion of the brain.
  - Vertebro-Basilar System
    - Supply posterior portion of the brain.



## CIRCULUS ARTERIOSUS (CIRCLE OF WILLIS)

Named after Thomas Willis (1621–1675), an English physician

- It is Formed by:
  - Two Anterior cerebral arteries
  - Two Internal carotid arteries
  - Two Posterior cerebral arteries
  - Two Posterior communicating arteries
  - One Anterior communicating artery
- It Gives numerous small vessels that penetrate the surface of the brain
  - Perforating arteries
- They are divided into:
  - Anterior perforating arteries
  - Posterior perforating arteries



# **ANTERIOR PERFORATING ARTERIES**

- Arise from:
  - Anterior cerebral artery
  - Anterior communicating artery
  - Middle cerebral artery
- Enter brain through:
  - Anterior perforated substance
    - irregularly quadrilateral area in front of the optic tract and behind the olfactory trigone.
- Supply:
  - Large part of basal ganglia
  - Optic chiasma
  - Internal capsule
  - Hypothalamus



# **POSTERIOR PERFORATING ARTERIES**

- Arise from:
  - Posterior cerebral artery
  - Posterior communicating artery
- Enter brain through:
  - Posterior Perforated substance
- Supply:
  - Ventral portion of Midbrain
  - Parts of Subthalamus and Hypothalamus



## **ANTERIOR CEREBRAL ARTERY**

#### • Supplies: Orbital and medial surfaces of frontal and parietal lobes



# MIDDLE CEREBRAL ARTERY

- O Supplies: Entire Superolateral surface:
  - Somatosensory Cortex
  - Motor Cortex
  - Broca's Area
    - Inked to speech production.
  - Heschl's Gyrus
    - to process incoming auditory information
  - Wernicke's Area
    - It is involved in the understanding of written and spoken language





## **POSTERIOR CEREBRAL ARTERY**

- Supplies:
  - Anterior and inferior temporal lobes
  - Uncus
    - Located on the tip end of the medial surface of the parahippocampal gyrus.
    - Part of the olfactory cortex that processes information from the sense of smell.
  - Inferior temporal gyri
  - Inferior and Medial Occipital lobe



## **CEREBRAL ARTERIES**





# DISTRIBUTION OF CEREBRAL ARTERIES

Cortical vascular territories

Cortical vascular territories





# **BASILAR ARTERY**

- Supplies: Midbrain and Cerebellum.
- Branches:
  - Anterior inferior cerebellar artery
  - Pontine branches
  - Superior cerebellar artery



## **ARTERIAL DISORDER**

#### O Stroke

- Sudden occlusion
- Hemorrhage
- o Aneurysm
  - localized, blood-filled balloonlike bulge in the wall of a blood vessel.

#### o Angioma

 is benign tumors derived from cells of the vascular or lymphatic vessel walls (epithelium) or derived from cells of the tissues surrounding these vessels.



# **ACCLUSION OF ACA**

in

#### • Manifestations:

 Motor disturbance contralateral distal leg

#### Difficulty in Prefrontal lobe Functions:

- Cognitive thinking
- Judgment
- Motor initiation
- Self monitoring





# **ACCLUSION OF MCA**

#### • Manifestations:

- Contralateral weakness of:
  - face, arm, and hand more than legs
- Contralateral sensory loss of:
  - face, arm, and hand more than legs
  - visual field cut (damage to optic radiation)

#### Aphasia: language disturbances

- Broca's: production
- Wernicke's: comprehension





# **ACCLUSION OF PCA**

#### • Manifestations:

- Visual disturbances
  - Contralateral homonymous hemianopsia
  - Bilateral lesions: cortical blindness
    - ✓ patients unaware they cannot see (Anton's syndrome)
- Memory impairment
  - If temporal lobe is affected



# **HOW WE ARE DOING ..?**

- Which statement(s) of the following is NOT Wrong?
- Anterior cerebral arteries supply Broca's and Wernicke's Area..!!
- Occlusion of MCA causes difficulty in Prefrontal lobe's functions..!!
- Middle cerebral arteries are part of Willis Circle..!!
- Aneurysm is benign tumors derived from cells of the vascular or lymphatic vessel walls..!!
- Posterior cerebral arteries supply anterior and inferior temporal lobes..!!

# **CEREBRAL VENOUS DRAINAGE**

- It involves:
  - Superficial (cortical) veins:
    - Drain the cortical surface
  - Deep veins:
    - Drain the deep structures
- These veins ultimately drain into:
  - Dural Venous Sinuses
- The Veins are thin walled and are devoid of valves.



- Lie on the brain surface, in the Subarchnoid space.
- They are divided into:
  - Superior cerebral veins
  - Inferior cerebral veins
  - Superficial middle cerebral vein



#### Superior Cerebral Veins

- 6 to 12 veins
- Drain lateral surface of brain above the lateral sulcus
- Terminate mainly into the Superior Sagittal sinus, and partly into superficial middle cerebral vein.





#### Inferior Cerebral Veins

- Run below the lateral sulcus
- Drain the lateral surface of the temporal lobe
- Terminate partly into superficial middle cerebral vein & partly into Transverse sinus.





#### Superficial Middle Cerebral Vein

Ο

- Runs along the lateral sulcus
- Terminates into the Cavernous sinus
- Connected posteriorly by Superior & Inferior anastomotic veins to Superior Sagittal & Transverse sinuses respectively.





## DEEP CEREBRAL VEINS

- They drain the internal structures;
  - Basal ganglia
  - Internal capsule
  - Thalamus
- They merge to form the Internal Cerebral Veins.
- The two veins unite in the midline to form the Great Cerebral vein.
- This short vessel is continuous with the Straight Sinus.





# **DURAL VENOUS SINUSES**



#### Blood flows from transverse & sigmoid sinuses into IJV





# **VENOUS DISORDER**

#### • Infarction

- refers to tissue death (necrosis) that is caused by a local lack of oxygen due to obstruction of the tissue's blood supply
- Sinus thrombosis:
  - SSS thrombosis
    - Superior Sagittal Sinus
    - Can complicates ear infection
  - Cavernous Sinus thrombosis
    - As a complication of infection in the dangerous area of the face
  - Obstruction of venous drainage of the brain leads to Cerebral swelling (edema) and raised Intracranial Pressure.



# ALSO, HOW WE ARE DOING ..?

- Which statement(s) of the following is Wrong?
- 1. Superior Cerebral Veins terminate mainly into the Superior Sagittal sinus, and partly into superficial middle cerebral vein..!!
- 2. Infarction refers to tissue death (necrosis)..!!
- 3. Superior Cerebral Veins drain lateral surface of brain above the lateral sulcus..!!
- 4. Inferior Cerebral Veins terminate partly into superficial middle cerebral vein & partly into Transverse sinus..!!
- 5. Superficial Middle Cerebral Vein drains the lateral surface of the temporal lobe..!!

