

# Ch 04-3 Numerical Integration I

Dr. Feras Fraige

# Outline

- Introduction to Numerical Integration
- The Trapezoidal Rule
- Simpson's Rule
- Comparing the Trapezoidal Rule with Simpson's Rule

# Introduction to Numerical Integration

## Numerical Quadrature

- The need often arises for evaluating the definite integral of a function that has no explicit antiderivative or whose antiderivative is not easy to obtain.
- The basic method involved in approximating  $\int_a^b f(x) dx$  is called **numerical quadrature**. It uses a sum  $\sum_{i=0}^n a_i f(x_i)$  to approximate  $\int_a^b f(x) dx$ .

# Introduction to Numerical Integration

## Quadrature based on interpolation polynomials

- The methods of quadrature in this section are based on the interpolation polynomials.
- The basic idea is to select a set of distinct nodes  $\{x_0, \dots, x_n\}$  from the interval  $[a, b]$ .
- Then integrate the Lagrange interpolating polynomial

$$P_n(x) = \sum_{i=0}^n f(x_i)L_i(x)$$

and its truncation error term over  $[a, b]$  to obtain:

# Introduction to Numerical Integration

## Quadrature based on interpolation polynomials (Cont'd)

$$\begin{aligned}\int_a^b f(x) dx &= \int_a^b \sum_{i=0}^n f(x_i) L_i(x) dx + \int_a^b \prod_{i=0}^n (x - x_i) \frac{f^{(n+1)}(\xi(x))}{(n+1)!} dx \\ &= \sum_{i=0}^n a_i f(x_i) + \frac{1}{(n+1)!} \int_a^b \prod_{i=0}^n (x - x_i) f^{(n+1)}(\xi(x)) dx\end{aligned}$$

where  $\xi(x)$  is in  $[a, b]$  for each  $x$  and

$$a_i = \int_a^b L_i(x) dx, \quad \text{for each } i = 0, 1, \dots, n$$

# Introduction to Numerical Integration

## Quadrature based on interpolation polynomials (Cont'd)

The quadrature formula is, therefore,

$$\int_a^b f(x) dx \approx \sum_{i=0}^n a_i f(x_i)$$

where

$$a_i = \int_a^b L_i(x) dx, \quad \text{for each } i = 0, 1, \dots, n$$

and with error given by

$$E(f) = \frac{1}{(n+1)!} \int_a^b \prod_{i=0}^n (x - x_i) f^{(n+1)}(\xi(x)) dx$$

# Numerical Integration: Trapezoidal Rule

## Derivation (1/3)

To derive the Trapezoidal rule for approximating  $\int_a^b f(x) dx$ , let  $x_0 = a$ ,  $x_1 = b$ ,  $h = b - a$  and use the linear Lagrange polynomial:

$$P_1(x) = \frac{(x - x_1)}{(x_0 - x_1)} f(x_0) + \frac{(x - x_0)}{(x_1 - x_0)} f(x_1)$$

Then

$$\begin{aligned} \int_a^b f(x) dx &= \int_{x_0}^{x_1} \left[ \frac{(x - x_1)}{(x_0 - x_1)} f(x_0) + \frac{(x - x_0)}{(x_1 - x_0)} f(x_1) \right] dx \\ &\quad + \frac{1}{2} \int_{x_0}^{x_1} f''(\xi(x))(x - x_0)(x - x_1) dx. \end{aligned}$$

# Numerical Integration: Trapezoidal Rule

## Derivation (2/3)

The product  $(x - x_0)(x - x_1)$  does not change sign on  $[x_0, x_1]$ , so the Weighted Mean Value Theorem for Integrals [▶ See Theorem](#) can be applied to the error term to give, for some  $\xi$  in  $(x_0, x_1)$ ,

$$\begin{aligned} & \int_{x_0}^{x_1} f'''(\xi(x))(x - x_0)(x - x_1) dx \\ &= f'''(\xi) \int_{x_0}^{x_1} (x - x_0)(x - x_1) dx \\ &= f'''(\xi) \left[ \frac{x^3}{3} - \frac{(x_1 + x_0)}{2}x^2 + x_0x_1x \right]_{x_0}^{x_1} \\ &= -\frac{h^3}{6}f'''(\xi) \end{aligned}$$



# Numerical Integration: Trapezoidal Rule

## Derivation (3/3)

Consequently, the last equation, namely

$$\int_{x_0}^{x_1} f''(\xi(x))(x - x_0)(x - x_1) dx = -\frac{h^3}{6}f''(\xi)$$

implies that

$$\begin{aligned}\int_a^b f(x) dx &= \left[ \frac{(x - x_1)^2}{2(x_0 - x_1)} f(x_0) + \frac{(x - x_0)^2}{2(x_1 - x_0)} f(x_1) \right]_{x_0}^{x_1} - \frac{h^3}{12} f''(\xi) \\ &= \frac{(x_1 - x_0)}{2} [f(x_0) + f(x_1)] - \frac{h^3}{12} f''(\xi)\end{aligned}$$

# Numerical Integration: Trapezoidal Rule

Using the notation  $h = x_1 - x_0$  gives the following rule:


## The Trapezoidal Rule

$$\int_a^b f(x) dx = \frac{h}{2}[f(x_0) + f(x_1)] - \frac{h^3}{12}f''(\xi)$$

### Note:


- The error term for the Trapezoidal rule involves  $f''$ , so the rule gives the exact result when applied to any function whose second derivative is identically zero, that is, any polynomial of degree one or less.
- The method is called the Trapezoidal rule because, when  $f$  is a function with positive values,  $\int_a^b f(x) dx$  is approximated by the area in a trapezoid, as shown in the following diagram.

# Trapezoidal Rule: The Area in a Trapezoid



# Numerical Integration: Simpson's Rule

Simpson's rule results from integrating over  $[a, b]$  the second Lagrange polynomial with equally-spaced nodes  $x_0 = a$ ,  $x_2 = b$ , and  $x_1 = a + h$ , where  $h = (b - a)/2$ :



# Numerical Integration: Simpson's Rule

## Simpson's Rule

$$\int_{x_0}^{x_2} f(x) dx = \frac{h}{3}[f(x_0) + 4f(x_1) + f(x_2)] - \frac{h^5}{90}f^{(4)}(\xi)$$

The error term in Simpson's rule involves the fourth derivative of  $f$ , so it gives exact results when applied to any polynomial of degree three or less.

# Trapezoidal Rule .v. Simpson's Rule

## Example

Compare the Trapezoidal rule and Simpson's rule approximations to

$\int_0^2 f(x) dx$  when  $f(x)$  is

(a)  $x^2$

(b)  $x^4$

(c)  $(x + 1)^{-1}$

(d)  $\sqrt{1 + x^2}$

(e)  $\sin x$

(f)  $e^x$

## Trapezoidal Rule .v. Simpson's Rule

### Solution (1/3)

On  $[0, 2]$ , the Trapezoidal and Simpson's rule have the forms

$$\text{Trapezoidal: } \int_0^2 f(x) dx \approx f(0) + f(2)$$

$$\text{Simpson's: } \int_0^2 f(x) dx \approx \frac{1}{3}[f(0) + 4f(1) + f(2)]$$

# Trapezoidal Rule .v. Simpson's Rule

## Solution (1/3)

On  $[0, 2]$ , the Trapezoidal and Simpson's rule have the forms

$$\text{Trapezoidal: } \int_0^2 f(x) dx \approx f(0) + f(2)$$

$$\text{Simpson's: } \int_0^2 f(x) dx \approx \frac{1}{3}[f(0) + 4f(1) + f(2)]$$

When  $f(x) = x^2$  they give

$$\text{Trapezoidal: } \int_0^2 f(x) dx \approx 0^2 + 2^2 = 4$$

$$\text{Simpson's: } \int_0^2 f(x) dx \approx \frac{1}{3}[(0^2) + 4 \cdot 1^2 + 2^2] = \frac{8}{3}$$



## Trapezoidal Rule .v. Simpson's Rule

### Solution (2/3)

- The approximation from Simpson's rule is exact because its truncation error involves  $f^{(4)}$ , which is identically 0 when  $f(x) = x^2$ .
- The results to three places for the functions are summarized in the following table.

# Trapezoidal Rule .v. Simpson's Rule

## Solution (3/3): Summary Results

$f(x)$	(a) $x^2$	(b) $x^4$	(c) $(x + 1)^{-1}$	(d) $\sqrt{1 + x^2}$	(e) $\sin x$	(f) $e^x$
Exact value	2.667	6.400	1.099	2.958	1.416	6.389
Trapezoidal	4.000	16.000	1.333	3.326	0.909	8.389
Simpson's	2.667	6.667	1.111	2.964	1.425	6.421

Notice that, in each instance, Simpson's Rule is significantly superior.