

The background of the slide features a close-up, slightly blurred image of a clock face with Roman numerals. Overlaid on the clock is a dark, coiled spiral spring, likely from a mechanical watch movement. The overall color palette is warm, with shades of orange, yellow, and brown.

# 10

## **PARAMETRIC EQUATIONS AND POLAR COORDINATES**

## PARAMETRIC EQUATIONS & POLAR COORDINATES

We have seen how to represent curves by parametric equations.

Now, we apply the methods of calculus to these parametric curves.

## 10.2

### Calculus with Parametric Curves

In this section, we will:

Use parametric curves to obtain formulas for  
tangents, areas, arc lengths, and surface areas.

## TANGENTS

In Section 10.1, we saw that some curves defined by parametric equations  $x = f(t)$  and  $y = g(t)$  can also be expressed—by eliminating the parameter—in the form  $y = F(x)$ .

- See Exercise 67 for general conditions under which this is possible.

## TANGENTS

If we substitute  $x = f(t)$  and  $y = g(t)$  in the equation  $y = F(x)$ , we get:

$$g(t) = F(f(t))$$

- So, if  $g$ ,  $F$ , and  $f$  are differentiable, the Chain Rule gives:

$$g'(t) = F'(f(t))f'(t) = F'(x)f'(t)$$

If  $f'(t) \neq 0$ , we can solve  
for  $F'(x)$ :

$$F'(x) = \frac{g'(t)}{f'(t)}$$

## TANGENTS

The slope of the tangent to the curve  $y = F(x)$  at  $(x, F(x))$  is  $F'(x)$ .

Thus, Equation 1 enables us to find tangents to parametric curves without having to eliminate the parameter.

Using Leibniz notation, we can rewrite Equation 1 in an easily remembered form:

$$\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} \quad \text{if } \frac{dx}{dt} \neq 0$$



## TANGENTS

If we think of a parametric curve as being traced out by a moving particle, then

- $dy/dt$  and  $dx/dt$  are the vertical and horizontal velocities of the particle.
- Formula 2 says that the slope of the tangent is the ratio of these velocities.

## TANGENTS

From Equation 2, we can see that the curve has:

- A horizontal tangent when  $dy/dt = 0$  (provided  $dx/dt \neq 0$ ).
- A vertical tangent when  $dx/dt = 0$  (provided  $dy/dt \neq 0$ ).

## TANGENTS

This information is useful for sketching parametric curves.

## TANGENTS

As we know from Chapter 4, it is also useful to consider  $d^2y/dx^2$ .

This can be found by replacing  $y$  by  $dy/dx$  in Equation 2:

$$\frac{d^2 y}{dx^2} = \frac{d}{dx} \left( \frac{dy}{dx} \right) = \frac{\frac{d}{dt} \left( \frac{dy}{dx} \right)}{\frac{dx}{dt}}$$

## TANGENTS

### Example 1

A curve  $C$  is defined by the parametric equations  $x = t^2$ ,  $y = t^3 - 3t$ .

- Show that  $C$  has two tangents at the point  $(3, 0)$  and find their equations.
- Find the points on  $C$  where the tangent is horizontal or vertical.
- Determine where the curve is concave upward or downward.
- Sketch the curve.

## TANGENTS

### Example 1 a

Notice that  $y = t^3 - 3t = t(t^2 - 3) = 0$   
when  $t = 0$  or  $t = \pm\sqrt{3}$ .

- Thus, the point  $(3, 0)$  on  $C$  arises from two values of the parameter:  $t = \sqrt{3}$  and  $t = -\sqrt{3}$
- This indicates that  $C$  crosses itself at  $(3, 0)$ .

## TANGENTS

### Example 1 a

Since

$$\frac{dy}{dx} = \frac{dy / dt}{dx / dt} = \frac{3t^2 - 3}{2t} = \frac{3}{2} \left( t - \frac{1}{t} \right)$$

the slope of the tangent when  $t = \pm\sqrt{3}$   
is:

$$dy/dx = \pm 6/(2\sqrt{3}) = \pm \sqrt{3}$$

So, the equations of the tangents at (3, 0) are:

$$y = \sqrt{3}(x - 3)$$

and

$$y = -\sqrt{3}(x - 3)$$



## TANGENTS

### Example 1 b

$C$  has a horizontal tangent when  $dy/dx = 0$ , that is, when  $dy/dt = 0$  and  $dx/dt \neq 0$ .

- Since  $dy/dt = 3t^2 - 3$ , this happens when  $t^2 = 1$ , that is,  $t = \pm 1$ .
- The corresponding points on  $C$  are  $(1, -2)$  and  $(1, 2)$ .

$C$  has a vertical tangent when  $dx/dt = 2t = 0$ , that is,  $t = 0$ .

- Note that  $dy/dt \neq 0$  there.
- The corresponding point on  $C$  is  $(0, 0)$ .

To determine concavity, we calculate the second derivative:


$$\frac{d^2 y}{dx^2} = \frac{\frac{d}{dt} \left( \frac{dy}{dx} \right)}{\frac{dx}{dt}} = \frac{\frac{3}{2} \left( 1 + \frac{1}{t^2} \right)}{2t} = \frac{3(t^2 + 1)}{4t^3}$$

- The curve is concave upward when  $t > 0$ .
- It is concave downward when  $t < 0$ .

## TANGENTS

### Example 1 d

Using the information from (b) and (c), we sketch C.



a. Find the tangent to the cycloid

$$x = r(\theta - \sin \theta), y = r(1 - \cos \theta)$$

at the point where  $\theta = \pi/3$ .

- See Example 7 in Section 10.1

b. At what points is the tangent horizontal?  
When is it vertical?

The slope of the tangent line is:

$$\frac{dy}{dx} = \frac{dy / d\theta}{dx / d\theta} = \frac{r \sin \theta}{r(1 - \cos \theta)} = \frac{\sin \theta}{1 - \cos \theta}$$

## TANGENTS

### Example 2 a

When  $\theta = \pi/3$ , we have

$$x = r \left( \frac{\pi}{3} - \sin \frac{\pi}{3} \right) = r \left( \frac{\pi}{3} - \frac{\sqrt{3}}{2} \right)$$

$$y = r \left( 1 - \cos \frac{\pi}{3} \right) = \frac{r}{2}$$

and

$$\frac{dy}{dx} = \frac{\sin(\pi/3)}{1 - \cos(\pi/3)} = \frac{\sqrt{3}/2}{1 - \frac{1}{2}} = \sqrt{3}$$

## TANGENTS

### Example 2 a

Hence, the slope of the tangent is  $\sqrt{3}$  .

Its equation is:

$$y - \frac{r}{2} = \sqrt{3} \left( x - \frac{r\pi}{3} + \frac{r\sqrt{3}}{2} \right)$$

or


$$\sqrt{3}x - y = r \left( \frac{\pi}{\sqrt{3}} - 2 \right)$$



## TANGENTS

### Example 2 a

The tangent is sketched here.



The tangent is horizontal when  
 $dy/dx = 0$ .


- This occurs when  $\sin \theta = 0$  and  $1 - \cos \theta \neq 0$ , that is,  $\theta = (2n - 1)\pi$ ,  $n$  an integer.
- The corresponding point on the cycloid is  $((2n - 1)\pi r, 2r)$ .

## TANGENTS

### Example 2 b

When  $\theta = 2n\pi$ , both  $dx/d\theta$  and  $dy/d\theta$  are 0.

It appears from the graph that there are vertical tangents at those points.



We can verify this by using l'Hospital's Rule as follows:

$$\begin{aligned}\lim_{\theta \rightarrow 2n\pi^+} \frac{dy}{dx} &= \lim_{\theta \rightarrow 2n\pi^+} \frac{\sin \theta}{1 - \cos \theta} \\ &= \lim_{\theta \rightarrow 2n\pi^+} \frac{\cos \theta}{\sin \theta} = \infty\end{aligned}$$

A similar computation shows  
that  $dy/dx \rightarrow -\infty$  as  $\theta \rightarrow 2n\pi^-$ .

- So, indeed, there are vertical tangents when  $\theta = 2n\pi$ , that is, when  $x = 2n\pi r$ .

## AREAS

We know that the area under a curve  $y = F(x)$  from  $a$  to  $b$  is

$$A = \int_a^b F(x) dx$$

where  $F(x) \geq 0$ .

## AREAS

Suppose the curve is traced out once by the parametric equations  $x = f(t)$  and  $y = g(t)$ ,  $\alpha \leq t \leq \beta$ .

- Then, we can calculate an area formula by using the Substitution Rule for Definite Integrals.


$$A = \int_{\alpha}^{\beta} y \, dx = \int_{\alpha}^{\beta} g(t) f'(t) \, dt$$
$$\left[ \text{or } \int_{\beta}^{\alpha} g(t) f'(t) \, dt \right]$$

## AREAS

### Example 3

Find the area under one arch  
of the cycloid

$$x = r(\theta - \sin \theta) \quad y = r(1 - \cos \theta)$$





One arch of the cycloid is given by  $0 \leq \theta \leq 2\pi$ .

Using the Substitution Rule with  
 $y = r(1 - \cos \theta)$  and  $dx = r(1 - \cos \theta) d\theta$ ,  
we have the following result.

## AREAS

## Example 3

$$\begin{aligned} A &= \int_0^{2\pi r} y \, dx \\ &= \int_0^{2\pi} r(1 - \cos \theta) r(1 - \cos \theta) \, d\theta \\ &= r^2 \int_0^{2\pi} (1 - \cos \theta)^2 \, d\theta \\ &= r^2 \int_0^{2\pi} (1 - 2\cos \theta + \cos^2 \theta) \, d\theta \\ &= r^2 \int_0^{2\pi} \left[ 1 - 2\cos \theta + \frac{1}{2}(1 + \cos 2\theta) \right] \, d\theta \\ &= r^2 \left[ \frac{3}{2}\theta - 2\sin \theta + \frac{1}{4}\sin 2\theta \right]_0^{2\pi} = r^2 \left( \frac{3}{2} \cdot 2\pi \right) = 3\pi r^2 \end{aligned}$$

The result of Example 3 says that the area under one arch of the cycloid is three times the area of the rolling circle that generates the cycloid (Example 7 in Section 10.1).

- Galileo guessed this result.
- However, it was first proved by the French mathematician Roberval and the Italian mathematician Torricelli.

## ARC LENGTH

We already know how to find the length  $L$  of a curve  $C$  given in the form

$$y = F(x), a \leq x \leq b$$

Formula 3 in Section 8.1 says that, if  $F'$  is continuous, then

$$L = \int_a^b \sqrt{1 + \left( \frac{dy}{dx} \right)^2} dx$$

## ARC LENGTH

Suppose that  $C$  can also be described by the parametric equations  $x = f(t)$  and  $y = g(t)$ ,  $\alpha \leq t \leq \beta$ , where  $dx/dt = f'(t) > 0$ .

- This means that  $C$  is traversed once, from left to right, as  $t$  increases from  $\alpha$  to  $\beta$  and  $f(\alpha) = a$  and  $f(\beta) = b$ .

## ARC LENGTH

Putting Formula 2 into Formula 3 and using the Substitution Rule, we obtain:

$$\begin{aligned} L &= \int_a^b \sqrt{1 + \left( \frac{dy}{dx} \right)^2} dx \\ &= \int_{\alpha}^{\beta} \sqrt{1 + \left( \frac{dy/dt}{dx/dt} \right)^2} \frac{dx}{dt} dt \end{aligned}$$

Since  $dx/dt > 0$ , we have:

$$L = \int_{\alpha}^{\beta} \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$



## ARC LENGTH

Even if  $C$  can't be expressed in the form  $y = f(x)$ , Formula 4 is still valid.

However, we obtain it by polygonal approximations.


## ARC LENGTH

We divide the parameter interval  $[\alpha, \beta]$  into  $n$  subintervals of equal width  $\Delta t$ .

If  $t_0, t_1, t_2, \dots, t_n$  are the endpoints of these subintervals, then  $x_i = f(t_i)$  and  $y_i = g(t_i)$  are the coordinates of points  $P_i(x_i, y_i)$  that lie on  $C$ .

## ARC LENGTH

So, the polygon with vertices  $P_0, P_1, \dots, P_n$  approximates  $C$ .



## ARC LENGTH

As in Section 8.1, we define the length  $L$  of  $C$  to be the limit of the lengths of these approximating polygons as  $n \rightarrow \infty$ :

$$L = \lim_{n \rightarrow \infty} \sum_{i=1}^n |P_{i-1}P_i|$$

## ARC LENGTH

The Mean Value Theorem, when applied to  $f$  on the interval  $[t_{i-1}, t_i]$ , gives a number  $t_i^*$  in  $(t_{i-1}, t_i)$  such that:

$$f(t_i) - f(t_{i-1}) = f'(t_i^*)(t_i - t_{i-1})$$

- If we let  $\Delta x_i = x_i - x_{i-1}$  and  $\Delta y_i = y_i - y_{i-1}$ , the equation becomes:

$$\Delta x_i = f'(t_i^*) \Delta t$$

## ARC LENGTH

Similarly, when applied to  $g$ , the Mean Value Theorem gives a number  $t_i^{**}$  in  $(t_{i-1}, t_i)$  such that:

$$\Delta y_i = g'(t_i^{**}) \Delta t$$

## ARC LENGTH

Therefore,

$$\begin{aligned} |P_{i-1}P_i| &= \sqrt{(\Delta x_i)^2 + (\Delta y_i)^2} \\ &= \sqrt{\left[f'(t_i^*)\Delta t\right]^2 + \left[g'(t_i^{**})\Delta t\right]^2} \\ &= \sqrt{\left[f'(t_i^*)\right]^2 + \left[g'(t_i^{**})\right]^2} \Delta t \end{aligned}$$

Thus,

$$L = \lim_{n \rightarrow \infty} \sum_{i=1}^n \sqrt{\left[ f' \left( t_i^* \right) \right]^2 + \left[ g' \left( t_i^{**} \right) \right]^2} \Delta t$$



## ARC LENGTH

The sum in Equation 5 resembles a Riemann sum for the function  $\sqrt{[f'(t)]^2 + [g'(t)]^2}$  .

However, is not exactly a Riemann sum because  $t_i^* \neq t_i^{**}$  in general.

## ARC LENGTH

Nevertheless, if  $f'$  and  $g'$  are continuous, it can be shown that the limit in Equation 5 is the same as if  $t_i^*$  and  $t_i^{**}$  were equal, namely,

$$L = \int_{\alpha}^{\beta} \sqrt{[f'(t)]^2 + [g'(t)]^2} dt$$

## ARC LENGTH

Thus, using Leibniz notation, we have the following result—which has the same form as Formula 4.

## THEOREM

## Theorem 6

Let a curve  $C$  is described by the parametric equations  $x = f(t)$ ,  $y = g(t)$ ,  $\alpha \leq t \leq \beta$ , where:

- $f'$  and  $g'$  are continuous on  $[\alpha, \beta]$ .
- $C$  is traversed exactly once as  $t$  increases from  $\alpha$  to  $\beta$ .

Then, the length of  $C$  is:

$$L = \int_{\alpha}^{\beta} \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

## ARC LENGTH

Notice that the formula in Theorem 6 is consistent with these general formulas from Section 8.1:

$$L = \int ds \quad \text{and} \quad (ds)^2 = (dx)^2 + (dy)^2$$

Suppose we use the representation of the unit circle given in Example 2 in Section 10.1:

$$x = \cos t \quad y = \sin t \quad 0 \leq t \leq 2\pi$$

- Then,

$$dx/dt = -\sin t \quad \text{and} \quad dy/dt = \cos t$$

## ARC LENGTH

### Example 4

- So, as expected, Theorem 6 gives:

$$\begin{aligned}\int_0^{2\pi} \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt &= \int_0^{2\pi} \sqrt{\sin^2 t + \cos^2 t} dt \\ &= \int_0^{2\pi} dt = 2\pi \\ &= 2\pi\end{aligned}$$

On the other hand, suppose we use the representation given in Example 3 in Section 10.1:

$$x = \sin 2t \quad y = \cos 2t \quad 0 \leq t \leq 2\pi$$

■ Then,

$$\underline{dx/dt} = 2\cos 2t \quad \text{and} \quad dy/dt = -2\sin 2t$$



## ARC LENGTH

### Example 4

- Then, the integral in Theorem 6 gives:

$$\begin{aligned}\int_0^{2\pi} \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt &= \int_0^{2\pi} \sqrt{4\cos^2 2t + 4\sin^2 2t} dt \\ &= \int_0^{2\pi} 2 dt \\ &= 4\pi\end{aligned}$$

Notice that the integral gives twice the arc length of the circle.

- This is because, as  $t$  increases from 0 to  $2\pi$ , the point  $(\sin 2t, \cos 2t)$  traverses the circle twice.

In general, when finding the length of a curve  $C$  from a parametric representation, we have to be careful to ensure that  $C$  is traversed only once as  $t$  increases from  $\alpha$  to  $\beta$ .

Find the length of one arch  
of the cycloid

$$x = r(\theta - \sin \theta) \quad y = r(1 - \cos \theta)$$

- From Example 3, we see that one arch is described by the parameter interval  $0 \leq \theta \leq 2\pi$ .

We have:

$$\frac{dx}{d\theta} = r(1 - \cos \theta)$$

and

$$\frac{dy}{d\theta} = r \sin \theta$$

Thus,

$$\begin{aligned} L &= \int_0^{2\pi} \sqrt{\left(\frac{dx}{d\theta}\right)^2 + \left(\frac{dy}{d\theta}\right)^2} d\theta \\ &= \int_0^{2\pi} \sqrt{r^2 (1 - \cos \theta)^2 + r^2 \sin^2 \theta} d\theta \\ &= \int_0^{2\pi} \sqrt{r^2 (1 - 2\cos \theta + \cos^2 \theta + \sin^2 \theta)} d\theta \\ &= r \int_0^{2\pi} \sqrt{2(1 - \cos \theta)} d\theta \end{aligned}$$

To evaluate this integral, we use the identity  $\sin^2 x = \frac{1}{2}(1 - \cos 2x)$  with  $\theta = 2x$ .

- This gives  $1 - \cos \theta = 2\sin^2(\theta/2)$ .

## ARC LENGTH

### Example 5

Since  $0 \leq \theta \leq 2\pi$ , we have  $0 \leq \theta/2 \leq \pi$ ,  
and so  $\sin(\theta/2) \geq 0$ .

$$\begin{aligned}\text{Therefore, } \sqrt{2(1 - \cos \theta)} &= \sqrt{4 \sin^2(\theta/2)} \\ &= 2|\sin(\theta/2)| \\ &= 2\sin(\theta/2)\end{aligned}$$




Hence,

$$\begin{aligned} L &= 2r \int_0^{2\pi} \sin(\theta/2) d\theta \\ &= 2r \left[ -2 \cos(\theta/2) \right]_0^{2\pi} \\ &= 2r [2 + 2] \\ &= 8r \end{aligned}$$

## ARC LENGTH

The result of Example 5 says that the length of one arch of a cycloid is eight times the radius of the generating circle.

- This was first proved in 1658 by Sir Christopher Wren.



## SURFACE AREA

In the same way as for arc length,  
we can adapt Formula 5 in Section 8.2  
to obtain a formula for surface area.

## SURFACE AREA

## Formula 7

Let the curve given by the parametric equations  $x = f(t)$ ,  $y = g(t)$ ,  $\alpha \leq t \leq \beta$ , be rotated about the  $x$ -axis, where:

- $f'$ ,  $g'$  are continuous.
- $g(t) \geq 0$ .

Then, the area of the resulting surface is given by:

$$S = \int_{\alpha}^{\beta} 2\pi y \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

## SURFACE AREA

The general symbolic formulas

$S = \int 2\pi y \, ds$  and  $S = \int 2\pi x \, ds$  (Formulas 7 and 8 in Section 8.2) are still valid.

However, for parametric curves,

we use:

$$ds = \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

Show that the surface area of a sphere of radius  $r$  is  $4\pi r^2$ .

- The sphere is obtained by rotating the semicircle

$$x = r \cos t \quad y = r \sin t \quad 0 \leq t \leq \pi$$

about the  $x$ -axis.

## SURFACE AREA

### Example 6

- So, from Formula 7, we get:

$$\begin{aligned} S &= \int_0^{\pi} 2\pi r \sin t \sqrt{(-r \sin t)^2 + (r \cos t)^2} dt \\ &= 2\pi \int_0^{\pi} r \sin t \sqrt{r^2 (\sin^2 t + \cos^2 t)} dt \\ &= 2\pi \int_0^{\pi} r \sin t * r dt \\ &= 2\pi r^2 \int_0^{\pi} \sin t dt \\ &= 2\pi r^2 (-\cos t) \Big|_0^{\pi} = 4\pi r^2 \end{aligned}$$