

The background of the slide features a warm, orange-toned image of a clock face with Roman numerals. A pendulum with a circular weight is visible on the left side, partially overlapping the clock face. The right side of the slide is a solid white area where the title and chapter number are located.

16

MULTIPLE INTEGRALS


16.4

Double Integrals in Polar Coordinates

In this section, we will learn:
How to express double integrals
in polar coordinates.


DOUBLE INTEGRALS IN POLAR COORDINATES

Suppose that we want to evaluate a double integral $\iint_R f(x, y) dA$, where R is one of the regions shown here.



(a) $R = \{(r, \theta) \mid 0 \leq r \leq 1, 0 \leq \theta \leq 2\pi\}$


© Thomson Higher Education



(b) $R = \{(r, \theta) \mid 1 \leq r \leq 2, 0 \leq \theta \leq \pi\}$


DOUBLE INTEGRALS IN POLAR COORDINATES

In either case, the description of R in terms of rectangular coordinates is rather complicated but R is easily described by polar coordinates.



(a) $R = \{(r, \theta) \mid 0 \leq r \leq 1, 0 \leq \theta \leq 2\pi\}$

© Thomson Higher Education



(b) $R = \{(r, \theta) \mid 1 \leq r \leq 2, 0 \leq \theta \leq \pi\}$

DOUBLE INTEGRALS IN POLAR COORDINATES

Recall from this figure that the polar coordinates (r, θ) of a point are related to the rectangular coordinates (x, y) by the equations

$$r^2 = x^2 + y^2$$

$$x = r \cos \theta$$

$$y = r \sin \theta$$




Fig. 16.4.2, p. 1010

POLAR RECTANGLE

The regions in the first figure are special cases of a polar rectangle

$$R = \{(r, \theta) \mid a \leq r \leq b, \alpha \leq \theta \leq \beta\}$$

shown here.




Fig. 16.4.3, p. 1010

POLAR RECTANGLE

To compute the double integral $\iint_R f(x, y) dA$

where R is a polar rectangle, we divide:

- The interval $[a, b]$ into m subintervals $[r_{j-1}, r_j]$ of equal width $\Delta r = (b - a)/m$.
- The interval $[\alpha, \beta]$ into n subintervals $[\theta_{j-1}, \theta_j]$ of equal width $\Delta \theta = (\beta - \alpha)/n$.

POLAR RECTANGLES

Then, the circles $r = r_i$ and the rays $\theta = \theta_j$ divide the polar rectangle R into the small polar rectangles shown here.




Fig. 16.4.4, p. 1010

POLAR SUBRECTANGLE

The “center” of the polar subrectangle

$$R_{ij} = \{(r, \theta) \mid r_{i-1} \leq r \leq r_i, \theta_{j-1} \leq \theta \leq \theta_j\}$$

has polar coordinates

$$r_i^* = \frac{1}{2} (r_{i-1} + r_i)$$

$$\theta_j^* = \frac{1}{2} (\theta_{j-1} + \theta_j)$$




Fig. 16.4.4, p. 1010

POLAR SUBRECTANGLE

We compute the area of R_{ij} using the fact that the area of a sector of a circle with radius r and central angle θ is $\frac{1}{2}r^2\theta$.

POLAR SUBRECTANGLE

Subtracting the areas of two such sectors, each of which has central angle $\Delta\theta = \theta_j - \theta_{j-1}$, we find that the area of R_{ij} is:

$$\begin{aligned}\Delta A_i &= \frac{1}{2} r_i^2 \theta - \frac{1}{2} r_{i-1}^2 \Delta\theta \\ &= \frac{1}{2} (r_i^2 - r_{i-1}^2) \Delta\theta \\ &= \frac{1}{2} (r_i + r_{i-1})(r_i - r_{i-1}) \Delta\theta \\ &= r_i^* \Delta r \Delta\theta\end{aligned}$$

POLAR RECTANGLES

We have defined the double integral

in terms of ordinary rectangles. $\iint_R f(x, y) dA$

However, it can be shown that, for continuous functions f , we always obtain the same answer using polar rectangles.

The rectangular coordinates of the center of R_{ij} are $(r_i^* \cos \theta_j^*, r_i^* \sin \theta_j^*)$.

So, a typical Riemann sum is:

$$\begin{aligned} & \sum_{i=1}^m \sum_{j=1}^n f(r_i^* \cos \theta_j^*, r_i^* \sin \theta_j^*) \Delta A_i \\ &= \sum_{i=1}^m \sum_{j=1}^n f(r_i^* \cos \theta_j^*, r_i^* \sin \theta_j^*) r_i^* \Delta r \Delta \theta \end{aligned}$$

POLAR RECTANGLES

If we write $g(r, \theta) = r f(r \cos \theta, r \sin \theta)$,
the Riemann sum in Equation 1 can be
written as:

$$\sum_{i=1}^m \sum_{j=1}^n g(r_i^*, \theta_j^*) \Delta r \Delta \theta$$

- This is a Riemann sum for the double integral

$$\int_{\alpha}^{\beta} \int_a^b g(r, \theta) dr d\theta$$

POLAR RECTANGLES

Thus, we have:

$$\begin{aligned}\iint_R f(x, y) dA &= \lim_{m, n \rightarrow \infty} \sum_{i=1}^m \sum_{j=1}^n f(r_i^* \cos \theta_j^*, r_i^* \sin \theta_j^*) \Delta A_i \\&= \lim_{m, n \rightarrow \infty} \sum_{i=1}^m \sum_{j=1}^n g(r_i^*, \theta_j^*) \Delta r \Delta \theta \\&= \int_{\alpha}^{\beta} \int_a^b g(r, \theta) dr d\theta \\&= \int_{\alpha}^{\beta} \int_a^b f(r \cos \theta, r \sin \theta) r dr d\theta\end{aligned}$$

CHANGE TO POLAR COORDS.

Formula 2

If f is continuous on a polar rectangle R given by

$$0 \leq a \leq r \leq b, \alpha \leq \theta \leq \beta$$

where $0 \leq \beta - \alpha \leq 2\pi$, then

$$\iint_R f(x, y) dA = \int_{\alpha}^{\beta} \int_a^b f(r \cos \theta, r \sin \theta) r dr d\theta$$

CHANGE TO POLAR COORDS.

Formula 2 says that we convert from rectangular to polar coordinates in a double integral by:

- Writing $x = r \cos \theta$ and $y = r \sin \theta$
- Using the appropriate limits of integration for r and θ
- Replacing dA by $dr d\theta$

CHANGE TO POLAR COORDS.

Be careful not to forget
the additional factor r on
the right side of Formula 2.

CHANGE TO POLAR COORDS.

A classical method for remembering the formula is shown here.

- The “infinitesimal” polar rectangle can be thought of as an ordinary rectangle with dimensions $r d\theta$ and dr .
- So, it has “area” $dA = r dr d\theta$.




Fig. 16.4.5, p. 1012

Evaluate

$$\iint_R (3x + 4y^2) dA$$

where R is the region in the upper half-plane bounded by the circles $x^2 + y^2 = 1$ and $x^2 + y^2 = 4$.

CHANGE TO POLAR COORDS.

Example 1

The region R can be described as:

$$R = \{(x, y) \mid y \geq 0, 1 \leq x^2 + y^2 \leq 4\}$$

CHANGE TO POLAR COORDS.

Example 1

It is the half-ring shown here.

In polar coordinates,
it is given by:

$$1 \leq r \leq 2, 0 \leq \theta \leq \pi$$




Fig. 16.4.1b, p. 1010

CHANGE TO POLAR COORDS.

Example 1

Hence, by Formula 2,

$$\begin{aligned} & \iint_R (3x + 4y^2) dA \\ &= \int_0^\pi \int_1^2 (3r \cos \theta + 4r^2 \sin^2 \theta) r dr d\theta \\ &= \int_0^\pi \int_1^2 (3r^2 \cos \theta + 4r^3 \sin^2 \theta) dr d\theta \\ &= \int_0^\pi [r^3 \cos \theta + r^4 \sin^2 \theta]_{r=1}^{r=2} d\theta \end{aligned}$$

CHANGE TO POLAR COORDS.

Example 1

$$\begin{aligned} &= \int_0^{\pi} (7 \cos \theta + 15 \sin^2 \theta) d\theta \\ &= \int_0^{\pi} \left[7 \cos \theta + \frac{15}{2} (1 - \cos 2\theta) \right] d\theta \\ &= 7 \sin \theta + \frac{15\theta}{2} - \frac{15}{4} \sin 2\theta \bigg|_0^{\pi} \\ &= \frac{15\pi}{2} \end{aligned}$$

Find the volume of the solid bounded by:

- The plane $z = 0$
- The paraboloid $z = 1 - x^2 - y^2$

CHANGE TO POLAR COORDS.

Example 2

If we put $z = 0$ in the equation of the paraboloid, we get $x^2 + y^2 = 1$.

- This means that the plane intersects the paraboloid in the circle $x^2 + y^2 = 1$.

CHANGE TO POLAR COORDS.

Example 2

So, the solid lies under the paraboloid and above the circular disk D given by $x^2 + y^2 \leq 1$.




Fig. 16.4.1a, p. 1010




Fig. 16.4.6, p. 1012

CHANGE TO POLAR COORDS.

Example 2

In polar coordinates, D is given by

$$0 \leq r \leq 1, 0 \leq \theta \leq 2\pi.$$

- As $1 - x^2 - y^2 = 1 - r^2$, the volume is:

$$V = \iint_D (1 - x^2 - y^2) dA = \int_0^{2\pi} \int_0^1 (1 - r^2) r dr d\theta$$

$$= \int_0^{2\pi} d\theta \int_0^1 (r - r^3) dr$$

$$= 2\pi \left[\frac{r^2}{2} - \frac{r^4}{4} \right]_0^1 = \frac{\pi}{2}$$

CHANGE TO POLAR COORDS.

Example 2

Had we used rectangular coordinates instead, we would have obtained:

$$\begin{aligned} V &= \iint_D (1 - x^2 - y^2) dA \\ &= \int_{-1}^1 \int_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}} (1 - x^2 - y^2) dy dx \end{aligned}$$

- This is not easy to evaluate because it involves finding $\int (1 - x^2)^{3/2} dx$

CHANGE TO POLAR COORDS.

What we have done so far can be extended to the more complicated type of region shown here.

- It's similar to the type II rectangular regions considered in Section 15.3




Fig. 16.4.7, p. 1013

CHANGE TO POLAR COORDS.

In fact, by combining Formula 2 in this section with Formula 5 in Section 16.3, we obtain the following formula.

CHANGE TO POLAR COORDS.

Formula 3

If f is continuous on a polar region of the form

$$D = \{(r, \theta) \mid \alpha \leq \theta \leq \beta, h_1(\theta) \leq r \leq h_2(\theta)\}$$

then

$$\iint_D f(x, y) dA = \int_{\alpha}^{\beta} \int_{h_1(\theta)}^{h_2(\theta)} f(r \cos \theta, r \sin \theta) r dr d\theta$$

CHANGE TO POLAR COORDS.

In particular, taking $f(x, y) = 1$, $h_1(\theta) = 0$, and $h_2(\theta) = h(\theta)$ in the formula, we see that the area of the region D bounded by $\theta = \alpha$, $\theta = \beta$, and $r = h(\theta)$ is:

$$\begin{aligned} A(D) &= \iint_D 1 \, dA = \int_{\alpha}^{\beta} \int_0^{h(\theta)} r \, dr \, d\theta = \int_{\alpha}^{\beta} \left[\frac{r^2}{2} \right]_0^{h(\theta)} d\theta \\ &= \int_{\alpha}^{\beta} \frac{1}{2} [h(\theta)]^2 d\theta \end{aligned}$$

- This agrees with Formula 3 in Section 10.4

CHANGE TO POLAR COORDS.

Example 3

Use a double integral to find the area enclosed by one loop of the four-leaved rose $r = \cos 2\theta$.

CHANGE TO POLAR COORDS.

Example 3

From this sketch of the curve, we see that a loop is given by the region

$$D = \{(r, \theta) \mid -\pi/4 \leq \theta \leq \pi/4, 0 \leq r \leq \cos 2\theta\}$$




Fig. 16.4.8, p. 1013

CHANGE TO POLAR COORDS.

Example 3

So, the area is:

$$\begin{aligned} A(D) &= \iint_D dA = \int_{-\pi/4}^{\pi/4} \int_0^{\cos 2\theta} r \, dr \, d\theta \\ &= \int_{-\pi/4}^{\pi/4} \left[\frac{1}{2} r^2 \right]_0^{\cos 2\theta} d\theta \\ &= \frac{1}{2} \int_{-\pi/4}^{\pi/4} \cos^2 2\theta \, d\theta \\ &= \frac{1}{4} \int_{-\pi/4}^{\pi/4} (1 + \cos 4\theta) \, d\theta \\ &= \frac{1}{4} \left[\theta + \frac{1}{4} \sin 4\theta \right]_{-\pi/4}^{\pi/4} = \frac{\pi}{8} \end{aligned}$$

Find the volume of the solid that lies:

- Under the paraboloid $z = x^2 + y^2$
- Above the xy -plane
- Inside the cylinder $x^2 + y^2 = 2x$

CHANGE TO POLAR COORDS.

Example 4

The solid lies above the disk D whose boundary circle has equation $x^2 + y^2 = 2x$.

- After completing the square, that is: $(x - 1)^2 + y^2 = 1$




Fig. 16.4.9, p. 1014




Fig. 16.4.10, p. 1014

CHANGE TO POLAR COORDS.

Example 4

In polar coordinates, we have:

$$x^2 + y^2 = r^2 \text{ and } x = r \cos \theta$$

So, the boundary circle becomes:

$$r^2 = 2r \cos \theta$$

or

$$r = 2 \cos \theta$$

CHANGE TO POLAR COORDS.

Example 4

Thus, the disk D is given by:

$$D =$$

$$\{(r, \theta) \mid -\pi/2 \leq \theta \leq \pi/2, 0 \leq r \leq 2 \cos \theta\}$$

CHANGE TO POLAR COORDS.

Example 4

So, by Formula 3, we have:

$$V$$

$$= \iint_D (x^2 + y^2) dA$$

$$= \int_{-\pi/2}^{\pi/2} \int_0^{2\cos\theta} r^2 r dr d\theta$$

$$= \int_{-\pi/2}^{\pi/2} \left[\frac{r^4}{4} \right]_0^{2\cos\theta} d\theta$$

$$= 4 \int_{-\pi/2}^{\pi/2} \cos^4 \theta d\theta$$

CHANGE TO POLAR COORDS.

Example 4

$$= 8 \int_0^{\pi/2} \cos^4 \theta \, d\theta$$

$$= 8 \int_0^{\pi/2} \left(\frac{1 + \cos 2\theta}{2} \right)^2 d\theta$$

$$= 2 \int_0^{\pi/2} \left[1 + 2 \cos 2\theta + \frac{1}{2} (1 + \cos 4\theta) \right] d\theta$$

$$= 2 \left[\frac{3}{2} \theta + \sin 2\theta + \frac{1}{8} \sin 4\theta \right]_0^{\pi/2}$$

$$= 2 \left(\frac{3}{2} \right) \left(\frac{\pi}{2} \right) = \frac{3\pi}{2}$$