

Chapter 5

Numerical Differentiation and Integration

Dr. Mohamed Abdelwahed
King Saud University

Introduction

Engineers are frequently confronted with the problem of differentiating functions which are defined in tabular or graphical form rather than as explicit functions. The interpretation of experimentally obtained data is a good example of this. A similar situation involves the integration of functions which have explicit forms that are difficult or impossible to integrate in terms of elementary functions.

Important Points of the Chapter 5

- I.** In this chapter we shall find the approximate solutions of derivative (first- and second-order) and antiderivative (definite integral only).
- II.** Given data points should be equally spaced only (length of each subinterval should be same). Smaller the length of the interval better the approximation.
- III.** Numerical methods for differentiation and integration can be derived using Lagrange interpolating polynomial at equally-spaced data points.
- IV.** Error term for each numerical method will be discuss

Numerical Differentiation

When a function is represented by a table of values, the most obvious approach is to differentiate the Lagrange interpolation formula

$$f(x) = p_n(x) + \frac{f^{(n+1)}(\eta(x))}{(n+1)!} \prod_{i=0}^n (x - x_i), \quad (5.3)$$

First Derivative Numerical Formulas

To obtain general formula for approximation of the first derivative of a function $f(x)$, we consider that $\{x_0, x_1, \dots, x_n\}$ are $(n+1)$ distinct equally spaced points in some interval I and function $f(x)$ is continuous and its $(n+1)$ th derivatives exist in the given interval, that is, $f \in C^{n+1}(I)$. Then by differentiating (5.3) with respect to x and at $x = x_k$, we have

$$f'(x_k) = \sum_{i=0}^n f(x_i) L'_i(x_k) + \frac{f^{(n+1)}(\eta(x_k))}{(n+1)!} \prod_{\substack{i=0 \\ i \neq k}}^n (x_k - x_i). \quad (5.4)$$

The formula (5.4) is called the $(n+1)$ -point formula to approximate $f'(x_k)$.

Two-point Formula

Consider two distinct points x_0 and x_1 , $x_1 = x_0 + h$ for some $h \neq 0$

$$f(x) \approx p_1(x) = \left(\frac{x - x_1}{x_0 - x_1} \right) f(x_0) + \left(\frac{x - x_0}{x_1 - x_0} \right) f(x_1).$$

By taking derivative with respect to x and at $x = x_0$, we obtain

$$f'(x)|_{x=x_0} \approx p_1'(x)|_{x=x_0} = \frac{f(x_0)}{x_0 - x_1} + \frac{f(x_1)}{x_1 - x_0}.$$

$$f'(x_0) \approx -\frac{f(x_0)}{h} + \frac{f(x_0 + h)}{h}, \quad \longleftrightarrow \quad f'(x_0) \approx \frac{f(x_0 + h) - f(x_0)}{h} = D_h f(x_0). \quad (5.6)$$

It is called the *two-point formula* for smaller values of h .

If $h < 0$, then the formula (5.6) is also called the *two-point backward-difference formula*, which can be written as

$$f'(x_0) \approx \frac{f(x_0) - f(x_0 - h)}{h}. \quad (5.7)$$

Example 5.1 Let $f(x) = e^x$. Then use the two-point formula to approximate $f'(2)$, when $h = 0.1$ and $h = 0.01$.

Solution. Using the formula (5.6), with $x_0 = 2$, we have

$$f'(2) \approx \frac{f(2+h) - f(2)}{h}.$$

Then for $h = 0.1$, we get

$$f'(2) \approx \frac{f(2.1) - f(2)}{0.1} \approx \frac{e^{2.1} - e^2}{0.1} = 7.7712.$$

Similarly, by using $h = 0.01$, we obtain

$$f'(2) \approx \frac{(e^{2.01} - e^2)}{0.01} = 7.4262.$$

Since the exact solution of $f'(2) = e^2$ is, 7.3891, so the corresponding actual errors with $h = 0.1$ and $h = 0.01$ are, -0.3821 and -0.0371 respectively. This shows that the approximation obtained with $h = 0.01$ is better than the approximation with $h = 0.1$. •

Similarly, by using the formula (5.7), with $x_0 = 2$, we have $f'(2) \approx \frac{f(2) - f(2-h)}{h}$,

then for $h = 0.1$, we have $f'(2) \approx \frac{f(2) - f(1.9)}{0.1} = \frac{e^2 - e^{1.9}}{0.1} = 7.0316$.

For $h = 0.01$, we have $f'(2) \approx \frac{e^2 - e^{1.99}}{0.01} = 7.3522$.

Error Term of Two-point Formula

$$f(x) - p_1(x) = \frac{f''(\eta(x))}{2!} \prod_{i=0}^1 (x - x_i), \quad \eta(x) \in (x_0, x_1).$$

$$\begin{aligned} f'(x_0) - p_1'(x_0) &= \left(\frac{d}{dx} f''(\eta(x)) \Big|_{x=x_0} \right) \frac{(x - x_0)(x - x_1)}{2} \\ &\quad + \frac{f''(\eta(x_0))}{2} \left(\frac{d}{dx} (x^2 - x(x_0 + h) - xx_0 + x_0(x_0 + h)) \Big|_{x=x_0} \right). \end{aligned}$$

Since $\frac{d}{dx} f''(\eta(x)) = 0$ only if $x = x_0$, so error in the forward-difference formula (5.6) is

$$E_F(f, h) = f'(x_0) - D_h f(x_0) = -\frac{h}{2} f''(\eta(x)), \quad \text{where } \eta(x) \in (x_0, x_1), \quad (5.8)$$

which is called the *error formula* of the two-point formula (5.6). Hence the formula (5.6) can be written as

$$f'(x_0) = \frac{f(x_0 + h) - f(x_0)}{h} - \frac{h}{2} f''(\eta(x)), \quad \text{where } \eta \in (x_0, x_1). \quad (5.9)$$

Example 5.3 Let $f(x) = x^2 \cos x$ and $h = 0.1$. Then

- (a) Compute the approximate value of $f'(1)$ using forward difference two-point formula (5.6).
- (b) Compute the error bound for your approximation using the formula (5.8).
- (c) Compute the absolute error.
- (d) What best maximum value of stepsize h required to obtain the approximate value of $f'(1)$ correct to 10^{-2} .

Solution. (a) Given $x_0 = 1, h = 0.1$, then by using the formula (5.6), we have

$$f'(1) \approx \frac{f(1+0.1) - f(1)}{0.1} = \frac{f(1.1) - f(1)}{0.1} = D_h f(1).$$

Thus

$$f'(1) \approx \frac{(1.1)^2 \cos(1.1) - (1)^2 \cos(1)}{0.1} \approx \frac{0.5489 - 0.5403}{0.1} = 0.0860,$$

which is the required approximation of $f'(x)$ at $x = 1$.

(b) To find the error bound, we use the formula (5.8), which gives

$$E_F(f, h) = -\frac{0.1}{2} f''(\eta(x)), \quad \text{where } \eta(x) \in (1, 1.1),$$

or

$$|E_F(f, h)| = \left| -\frac{0.1}{2} \right| |f''(\eta(x))|, \quad \text{for } \eta \in (1, 1.1).$$

The second derivative $f''(x)$ of the function can be found as

$$f(x) = x^2 \cos x, \quad \text{gives } f''(x) = (2 - x^2) \cos x - 4x \sin x.$$

The value of the second derivative $f''(\eta(x))$ cannot be computed exactly because $\eta(x)$ is not known. But one can bound the error by computing the largest possible value for $|f''(\eta(x))|$. So bound $|f''|$ on $[1, 1.1]$ can be obtain

$$M = \max_{1 \leq x \leq 1.1} |(2 - x^2) \cos x - 4x \sin x| = 3.5630,$$

at $x = 1.1$. Since $|f''(\eta(x))| \leq M$, therefore, for $h = 0.1$, we have

$$|E_F(f, h)| \leq \frac{0.1}{2} M = 0.05(3.5630) = 0.1782,$$

which is the possible maximum error in our approximation.

(c) Since the exact value of the derivative $f'(1)$ is 0.2392, therefore the absolute error $|E|$ can be computed as follows:

$$|E| = |f'(1) - D_h f(1)| = |0.2391 - 0.0860| = 0.1531.$$

(d) Since the given accuracy required is 10^{-2} , so

$$|E_F(f, h)| = \left| -\frac{h}{2} f''(\eta(x)) \right| \leq 10^{-2},$$

for $\eta(x) \in (1, 1.1)$. This gives

$$\frac{h}{2} M \leq 10^{-2}, \quad \text{or} \quad h \leq \frac{(2 \times 10^{-2})}{M}.$$

Using $M = 3.5630$, we obtain

$$h \leq \frac{2}{356.3000} = 0.0056,$$

which is the best maximum value of h to get the required accuracy. ●

Example 5.1 Let $f(x) = x^3$ be defined in the interval $[0.2, 0.3]$. Use the error formula (5.8) of two-point formula for the approximation of $f'(0.2)$ to compute a value of unknown point η .

Solution. Since the exact value of the first derivative of the function at $x_0 = 0.2$ is

$$f'(x) = 3x^2 \quad \text{and} \quad f'(0.2) = 3(0.2)^2 = 0.12,$$

and the approximate value of $f'(0.2)$ using two point formula is

$$f'(0.2) \approx \frac{f(0.3) - f(0.2)}{0.1} = \frac{(0.3)^3 - (0.2)^3}{0.1} = 0.19,$$

so error E can be calculated as

$$E = 0.12 - 0.19 = -0.07.$$

Using the formula (5.6) and $f''(\eta) = 6\eta$, we have, $-0.07 = -\frac{0.1}{2}6\eta$, and solving for η , we get the value of the unknown point $\eta = 0.233$ in the interval $(0.2, 0.3)$. •

Three-point Central Difference Formula

Consider the quadratic Lagrange interpolating polynomial $p_2(x)$ to the three distinct equally spaced points x_0, x_1 , and x_2 , with $x_1 = x_0 + h$ and $x_2 = x_0 + 2h$, for smaller value h , we have

$$f(x) \approx p_2(x) = \frac{(x - x_1)(x - x_2)}{(x_0 - x_1)(x_0 - x_2)}f(x_0) + \frac{(x - x_0)(x - x_2)}{(x_1 - x_0)(x_1 - x_2)}f(x_1) + \frac{(x - x_0)(x - x_1)}{(x_2 - x_0)(x_2 - x_1)}f(x_2). \quad (5.11)$$

Now taking the derivative of the above expression with respect to x and then take $x = x_k$, for $k = 0, 1, 2$, we have

$$f'(x_k) \approx \frac{(2x_k - x_1 - x_2)}{(x_0 - x_1)(x_0 - x_2)}f(x_0) + \frac{(2x_k - x_0 - x_2)}{(x_1 - x_0)(x_1 - x_2)}f(x_1) + \frac{(2x_k - x_0 - x_1)}{(x_2 - x_0)(x_2 - x_1)}f(x_2).$$

Firstly, we take $x_k = x_1$,

$$\begin{aligned} f'(x_1) &\approx \frac{(2x_1 - x_1 - x_2)}{(x_0 - x_1)(x_0 - x_2)}f(x_0) + \frac{(2x_1 - x_0 - x_2)}{(x_1 - x_0)(x_1 - x_2)}f(x_1) + \frac{(2x_1 - x_0 - x_1)}{(x_2 - x_0)(x_2 - x_1)}f(x_2). \\ &\approx \frac{f(x_1 + h) - f(x_1 - h)}{2h} = D_h f(x_1). \end{aligned}$$

It is called the *three-point central-difference formula*

Error Formula of Central Difference Formula

$$E_C(f, h) = f'(x_1) - D_h f(x_1) = -\frac{h^2}{6} f'''(\eta(x_1)), \quad \text{where } \eta(x_1) \in (x_1 - h, x_1 + h).$$

Three-point Forward Difference Formula

by taking $x_k = x_0$ in the formula (5.11),

$$f'(x_0) \approx \frac{-3f(x_0) + 4f(x_0 + h) - f(x_0 + 2h)}{2h} = D_h f(x_0),$$

which is called the *three-point forward-difference formula*

Error Formula

$$E_F(f, h) = \frac{h^2}{3} f'''(\eta(x_0)), \quad \text{where } \eta(x_0) \in (x_0, x_0 + 2h).$$

Three-point Backward Difference Formula:

Similarly, taking $x_k = x_2$ in the formula (5.11),

$$f'(x_2) \approx \frac{f(x_2 - 2h) - 4f(x_2 - h) + 3f(x_2)}{2h} = D_h f(x_2),$$

which is called the *three-point backward-difference formula*

Error Formula

$$E_B(f, h) = \frac{h^2}{3} f'''(\eta(x_2)), \quad \text{where } \eta(x_2) \in (x_2 - 2h, x_2).$$

Example 5.5 Let $f(x) = x^2 + \cos x$ and $h = 0.1$. Then

- (a) Compute the approximate value of $f'(1)$ by using three-point central difference formula (5.13).
- (b) Compute the error bound for your approximation using (5.14).
- (c) Compute the absolute error.
- (d) What is the best maximum value of stepsize h required to obtain the approximate value of $f'(1)$ correct to 10^{-2} .

Solution. (a) Given $x_1 = 1, h = 0.1$, then using the formula (5.13), we have

$$f'(1) \approx \frac{f(1+0.1) - f(1-0.1)}{2(0.1)} = \frac{f(1.1) - f(0.9)}{0.2} = D_h f(1).$$

Then

$$f'(1) \approx \frac{[(1.1)^2 + \cos(1.1)] - [(0.9)^2 + \cos(0.9)]}{0.2} \approx \frac{1.6636 - 1.4316}{0.2} = 1.1600.$$

(b) By using the error formula (5.14), we have

$$|E_C(f, h)| = \left| -\frac{(0.1)^2}{6} |f'''(\eta(x_1))| \right|, \quad \text{for } \eta(x_1) \in (0.9, 1.1).$$

Since $f'''(\eta(x_1)) = \sin(\eta(x_1))$. The above formula cannot be computed exactly because $\eta(x_1)$ is not known. But one can bound the error by computing the largest possible value for $|f'''(\eta(x_1))|$. So bound $|f'''|$ on $[0.9, 1.1]$ is

$$M = \max_{0.9 \leq x \leq 1.1} |\sin x| = 0.8912,$$

at $x = 1.1$. Thus, for $|f'''(\eta(x_1))| \leq M$ and $h = 0.1$, gives

$$|E_C(f, h)| \leq \frac{0.01}{6} M = \frac{0.01}{6} (0.8912) = 0.0015,$$

which is the possible maximum error in our approximation.

(c) Since the exact value of the derivative $f'(1)$ is, $(2(1) - \sin 1) = 1.1585$, therefore, the absolute error $|E|$ can be computed as follows

$$|E| = |f'(1) - D_h f(1)| = |(2 - \sin 1) - 1.1600| = |1.1585 - 1.1600| = 0.0015.$$

(d) Since the given accuracy required is 10^{-2} , so

$$|E_C(f, h)| = \left| -\frac{h^2}{6} f'''(\eta(x_1)) \right| \leq 10^{-2},$$

for $\eta(x_1) \in (0.9, 1.1)$. Then

$$\frac{h^2}{6} M \leq 10^{-2}, \quad h \leq \sqrt{\frac{6 \times 10^{-2}}{M}}, \quad h \leq \sqrt{\frac{6 \times 10^{-2}}{0.8912}} = 0.2312,$$

the best maximum value of h .

•

Example 5.6 Consider the following table for set of data points

x	1	1.6	2	2.3	2.8	3	3.9	4	4.8	5
$f(x)$	0.00	0.47	0.69	0.83	1.03	1.10	1.36	1.39	1.57	1.61

- (a) Use three-point formula for smaller value of h to find approximation of $f'(3)$.
(b) The function tabulated is $\ln x$, find error bound and absolute error for the approximation of $f'(3)$.
(c) What is the best maximum value of stepsize h required to obtain the approximate value of $f'(3)$ within the accuracy 10^{-4} .

Solution. (a) For the given table of data points, we can use all three-points formulas as for the central difference we can take

$$x_0 = x_1 - h = 2, \quad x_1 = 3, \quad x_2 = x_1 + h = 4, \quad \text{gives } h = 1,$$

for the forward difference formula we can take

$$x_0 = 3, \quad x_1 = x_0 + h = 3.9, \quad x_2 = x_0 + 2h = 4.8, \quad \text{gives } h = 0.9,$$

and for the backward difference formula we can take

$$x_0 = x_2 - 2h = 1.6, \quad x_1 = x_2 - h = 2.3, \quad x_2 = 3, \quad \text{gives } h = 0.7.$$

Since we know that smaller the value of h better the approximation of the derivative of the function, therefore, for the given problem, backward difference is the best formula to find approximation of $f'(3)$ as

$$f'(3) \approx \frac{f(1.6) - 4f(2.3) + 3f(3)}{2(0.7)} \approx \frac{[0.47 - 4(0.83) + 3(1.10)]}{1.4} = 0.3214.$$

(b) Using error term of backward difference formula, we have

$$E_B(f, h) = \frac{h^2}{3} f'''(\eta), \quad \text{or} \quad |E_B(f, h)| \leq \frac{h^2}{3} |f'''(\eta)|.$$

Taking $|f'''(\eta(x_2))| \leq M = \max_{1.6 \leq x \leq 3} |f'''(x)| = \max_{1.6 \leq x \leq 3} |2/x^3| = 0.4883$. Thus using $h = 0.7$, we obtain

$$|E_B(f, h)| \leq \frac{(0.7)^2}{3} (0.4883) = 0.0798,$$

the required error bounds for the approximations. To compute the absolute error we do as

$$|E| = |f'(3) - 0.3214| = |0.3333 - 0.3214| = 0.0119.$$

(c) Since the given accuracy required is 10^{-4} , so

$$|E_B(f, h)| = \left| \frac{h^2}{3} f'''(\eta) \right| \leq 10^{-4},$$

for $\eta \in (1.6, 3)$. Then

$$\frac{h^2}{3} M \leq 10^{-4}.$$

Solving for h by taking $M = 0.4883$, we obtain

$$h^2 \leq \frac{3 \times 10^{-4}}{0.4883} = 0.0248,$$

and so $h = 0.025$ the best maximum value of h .

•

Example 5.7 Use the best three-point formula to find approximation of $f'(1.5)$ using the following table for set of data points

x	1	1.6	2	2.3	2.8	3	3.9	4	4.8	5
$f(x)$	0.00	0.47	0.69	0.83	1.03	1.10	1.36	1.39	1.57	1.61

Solution. The best three-point formula for this problem is the central difference because this formula does not need the value of $f(1.5)$ which is not given in the table while the other two need this value. Since $x_1 = 1.5$, so by taking $x_1 + h = 2$ and $x_1 - h = 1$, gives $h = 0.5$, we have

$$f'(1.5) \approx \frac{f(2) - f(1)}{2(0.5)} = \frac{0.69 - 0.0}{1} = 0.69,$$

the required approximation of $f'(1.5)$. •

Example 5.10 Consider following set of data points

x	0.0	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.9	1.0	1.1	1.2
$f(x)$	1.00	1.10	1.18	1.26	1.32	1.38	1.43	1.47	1.52	1.54	1.55	1.56

Use the table, find the best approximation of $f'(0.75)$ and the worst approximations of $f'(0.1)$ by using three-point formulas.

Solution. For the best approximation of $f'(0.75)$, we have to take small value of $h = 0.15$, so using the central difference three-point formula (5.13), we get

$$f'(0.75) \approx \frac{f(0.9) - f(0.6)}{2(0.15)} \approx \frac{1.52 - 1.43}{0.3} = 0.3,$$

while the exact value of $f'(0.75)$ is 0.3184. For the worst approximation of $f'(0.1)$, we have to take big value of $h = 0.5$, so using the forward difference three-point formula (5.16), we get

$$f'(0.1) \approx \frac{-3f(0.1) + 4f(0.6) - f(1.1)}{2(0.5)} \approx \frac{-3(1.1) + 4(1.43) - 1.55}{1} \approx 0.8700,$$

the required worst approximation. •

Second Derivative Numerical Formula

Three-point Central Difference Formula

$$f''(x_1) \approx \frac{f(x_1 + h) - 2f(x_1) + f(x_1 - h)}{h^2}$$

is called the *three-point central-difference formula* for the approximation of the second derivative of a function $f(x)$ at the given point $x = x_1$.

Error Formula

$$E_C(f, h) = -\frac{h^2}{12}f^{(4)}(\eta(x_1)), \quad \text{where } \eta(x_1) \in (x_1 - h, x_1 + h).$$

Example 5.16 Find the approximation of $f''(0.8)$ by using the following set of data points using numerical rule:

x	0.0	0.11	0.24	0.3	0.4	0.5	0.6	0.72	0.8	0.9	1.05	1.11	1.2
$f(x)$	1.00	1.10	1.2	1.26	1.32	1.38	1.43	1.47	1.50	1.52	1.55	1.55	1.56

If the function is $f(x) = x + \cos x$, then compute an error bound for your approximation. How many subintervals approximate the given derivative to within accuracy of 10^{-6} using this differentiation rule ?

Solution. Given $x_1 = 0.8, h = 0.4$, then the formula (5.20) becomes

$$f''(0.8) \approx \frac{f(0.8 + 0.4) - 2f(0.8) + f(0.8 - 0.4)}{(0.4)^2} = D_h^2 f(1),$$

or

$$f''(0.8) \approx \frac{f(1.2) - 2f(0.8) + f(0.4)}{0.16} = -0.75 = D_h^2 f(1).$$

To compute the error bound for our approximation in part (a), we use the formula as

$$|E_C(f, h)| = \left| -\frac{h^2}{12} |f^{(4)}(\eta(x_1))| \right|, \quad \text{for } \eta(x_1) \in (0.9, 1.1).$$

The fourth derivative of the given function at $\eta(x_1)$ is $f^{(4)}(\eta(x_1)) = \cos \eta(x_1)$, and it cannot be computed exactly because $\eta(x_1)$ is not known. But one can bound the error by computing the largest possible value for $|f^{(4)}(\eta(x_1))|$. So bound $|f^{(4)}|$ on the interval $(0.4, 1.2)$ is

$$M = \max_{0.4 \leq x \leq 1.2} |\cos x| = 0.9211,$$

at $x = 0.4$, Thus, for $|f^{(4)}(\eta(x))| \leq M$, we have the possible maximum error as

$$|E_C(f, h)| \leq \frac{h^2}{12} M \leq \frac{(0.4)^2}{12} (0.9211) = 0.01228.$$

Since the given accuracy required is 10^{-6} , so

$$|E_C(f, h)| = \left| -\frac{h^2}{12} f^{(4)}(\eta(x_1)) \right| \leq 10^{-6},$$

for $\eta(x_1) \in (0.4, 1.2)$. Then for $|f^{(4)}(\eta(x_1))| \leq M$, we have

$$\frac{h^2}{12} M \leq 10^{-6}, \quad \frac{(1.2 - 0.4)^2}{n^2} \leq \frac{(12 \times 10^{-6})}{M},$$

solving for n , we get

$$n \geq \sqrt{(0.64 \times 0.9211 \times 10^6)/12} \geq 221.6424, \quad \text{gives } n = 222,$$

the required subintervals for approximating the given derivative to the given accuracy . •

Numerical Integration

we wish to find an approximation to the *definite integral*

$$I(f) = \int_a^b f(x)dx,$$

An obvious approach is to replace a function $f(x)$ in the integral (5.21) by

$$I(f) = \int_a^b f(x)dx \approx \int_a^b p(x)dx.$$

Simple Trapezoidal Rule


$x_0 = a, x_1 = b$ and $h = x_1 - x_0$,

$$f(x) \approx p_1(x) = \left(\frac{x - x_1}{x_0 - x_1}\right) f(x_0) + \left(\frac{x - x_0}{x_1 - x_0}\right) f(x_1).$$

$$\begin{aligned} \int_{x_0}^{x_1} f(x)dx &\approx \frac{f(x_0)}{x_0 - x_1} \int_{x_0}^{x_1} (x - x_1)dx + \frac{f(x_1)}{x_1 - x_0} \int_{x_0}^{x_1} (x - x_0)dx, \\ &\approx \frac{f(x_0)}{x_0 - x_1} \left[\frac{(x - x_1)^2}{2} \Big|_{x_0}^{x_1} \right] + \frac{f(x_1)}{x_1 - x_0} \left[\frac{(x - x_0)^2}{2} \Big|_{x_0}^{x_1} \right] \approx \frac{(x_1 - x_0)}{2} [f(x_0) + f(x_1)], \end{aligned}$$

and by taking $h = x_1 - x_0$, we get

$$\int_a^b f(x)dx \approx T_1(f) = \frac{h}{2} [f(x_0) + f(x_1)].$$



Example 5.16 Approximate the following integral

$$\int_1^2 \frac{1}{x+1} dx,$$

using the Trapezoidal rule and compute the absolute error.

Solution. Given $f(x) = \frac{1}{x+1}$ and $h = 1$, so using Trapezoidal rule (5.28), gives

$$T_1(f) = \frac{1}{2}[f(1) + f(2)] = 0.4167.$$

The exact solution of the given integral is

$$I(f) = \ln(3/2) = 0.4055, \quad \text{so} \quad |E_{T_1}(f)| = |I(f) - T_1(f)| = |0.4055 - 0.4167| = 0.0112,$$

is the required absolute error. •

Composite Trapezoidal Rule

The interval $[a, b]$ is partitioned into n subintervals (x_{i-1}, x_i) , $i = 1, 2, \dots, n$ with $a = x_0$ and $b = x_n$ of equal width $h = (b - a)/n$

Theorem 5.2 (Composite Trapezoidal Rule)

Let $f \in C^2[a, b]$, n may be odd or even, $h = (b - a)/n$, and $x_i = a + ih$ for each $i = 0, 1, 2, \dots, n$. Then the composite Trapezoidal rule for n subintervals can be written as

$$\int_{a=x_0}^{b=x_n} f(x)dx \approx T_n(f) = \frac{h}{2} \left[f(a) + 2 \sum_{i=1}^{n-1} f(x_i) + f(b) \right]. \quad (5.29)$$

Proof. Since for the composite form of the Trapezoidal rule, the interval is divided into n equal subintervals of width h so that $h = \frac{b-a}{n}$, and $(n+1)$ distinct points $a = x_0 < x_1 < x_2 \dots < x_n = b$, then we have

$$\int_a^b f(x)dx = \int_{x_0}^{x_1} f(x)dx + \int_{x_1}^{x_2} f(x)dx + \dots + \int_{x_{n-1}}^{x_n} f(x)dx.$$

Applying the Trapezoidal rule (5.28) for one strip to each of these integral, we have

$$\int_a^b f(x)dx \approx \frac{h}{2}[f(x_0) + f(x_1)] + \frac{h}{2}[f(x_1) + f(x_2)] + \dots + \frac{h}{2}[f(x_{n-1}) + f(x_n)].$$

Note that each of the interior point is counted twice and therefore has a coefficient of two whereas the endpoints are counted once and therefore has a coefficient one. •

Example 5.17 Evaluate the integral $\int_0^1 e^{4x} dx$ by using the Trapezoidal rule with $n = 1, 2, 4, 8$. Also compute the corresponding actual errors.

Solution. For $n = 1$, we use the formula (5.28) for $h = 1$, as follows

$$T_1(f) = \frac{1}{2} [f(0) + f(1)] = 27.7991.$$

For $n = 2$, using the formula (5.29) and $h = 0.5$, we have

$$T_2(f) = \frac{0.5}{2} [f(0) + 2f(0.5) + f(1)] = 17.5941.$$

For $n = 4$, using the formula (5.29) and $h = 0.25$, we have

$$T_4(f) = \frac{0.25}{2} [f(0) + 2[f(0.25) + f(0.5) + f(0.75)] + f(1)] = 14.4980.$$

Finally, for $n = 8$, using (5.29) and $h = 0.125$, we have

$$\begin{aligned} T_8(f) &= \frac{0.125}{2} [f(0) + 2[f(0.125) + f(0.25) + f(0.375) + f(0.5) \\ &\quad + f(0.625) + f(0.75) + f(0.875)] + f(1)] = 13.6776. \end{aligned}$$

Since the exact value of the given integral is

$$I(f) = \frac{1}{4} [e^4 - 1] = 13.4000.$$

So the corresponding actual errors are, -14.3991 , -4.1941 , -1.0980 and -0.2776 , respectively, which decrease by a factor of about *four* at each stage. •

Error Terms for Trapezoidal Rule

Theorem 5.3 (Error term for Simple Trapezoidal Rule)

Let $f \in C^2[a, b]$, and $h = (b - a)$. The local error that the simple Trapezoidal rule (5.28) makes in estimating the definite integral (5.21) is

$$E_{T_1}(f) = -\frac{h^3}{12}f''(\eta(x)), \quad (5.30)$$

where $\eta(x) \in (a, b)$.

Error Term for Composite Trapezoidal Rule

The *global error* of the Trapezoidal rule (5.29) equals the sum of n local errors of the Trapezoidal rule (5.28), that is

$$\begin{aligned} E_{T_n}(f) &= -\frac{h^3}{12}f''(\eta_1(x)) - \frac{h^3}{12}f''(\eta_2(x)) - \cdots - \frac{h^3}{12}f''(\eta_n(x)), \\ &= -\frac{h^3}{12} \sum_{i=1}^n f''(\eta_i(x)), \quad \text{for } \eta_i(x) \in (x_{i-1}, x_i), \\ &= -\frac{h^3}{12} n f''(\eta(x)), \\ &= -\frac{h^2}{12} (b - a) f''(\eta(x)), \quad \eta(x) \in (a, b). \end{aligned}$$

Example 5.20 (a) Find approximation of $\int_1^2 f(x) dx$ taking $h = 0.2$ by using the following set of data points

x	1.0	1.1	1.2	1.3	1.4	1.5	1.6	1.7	1.8	1.9	2.0
$f(x)$	0.368	0.366	0.361	0.354	0.355	0.335	0.323	0.311	0.298	0.284	0.271

The function tabulated is xe^{-x} , compute error bound and the absolute error for the approximation using Trapezoidal rule.

(b) How many subintervals approximate the given integral to an accuracy of at least 10^{-6} ?

Solution. (a) Given $h = 0.2$, so we have the select following set of data points for Trapezoidal rule as

x	1.0	1.2	1.4	1.6	1.8	2.0
$f(x)$	0.368	0.361	0.355	0.323	0.298	0.271

so the composite Trapezoidal rule (5.29) for six points can be written as

$$\int_1^2 f(x) dx \approx T_5(f) = \frac{h}{2} [f(x_0) + 2(f(x_1) + f(x_2) + f(x_3) + f(x_4)) + f(x_5)],$$

and by using the given values, we get

$$\int_1^2 f(x) dx \approx 0.1 [0.368 + 2(0.361 + 0.355 + 0.323 + 0.298) + 0.271] = 0.3313.$$

The second derivative of the function $f(x) = xe^{-x}$ can be obtain as

$$f'(x) = (1 - x)e^{-x} \quad \text{and} \quad f''(x) = (x - 2)e^{-x}.$$

Since $\eta(x)$ is unknown point in $(1, 2)$, therefore, the bound $|f''|$ on $[1, 2]$ is

$$M = \max_{1 \leq x \leq 2} |f''(x)| = \max_{1 \leq x \leq 2} |(x - 2)e^{-x}| = 0.3679,$$

at $x = 1$. Thus the error formula (5.35) becomes

$$|E_{T_5}(f)| \leq \frac{(0.2)^2(1)}{12}(0.3679) = 0.0012,$$

which is the possible maximum error in our approximation.

We can easily compute the exact value of the given integral as

$$\int_1^2 xe^{-x} dx = (-xe^{-x} - e^{-x}) \Big|_1^2 = 0.3298.$$

Thus the absolute error $|E|$ in our approximation is given as

$$|E| = |0.3298 - T_5(f)| = |0.3298 - 0.3313| = 0.0015.$$

(b) To find the minimum subintervals for the given accuracy, we use the formula (5.35) such that


$$|E_{T_n}(f)| \leq \frac{|-(b-a)^3|}{12n^2} M \leq 10^{-6},$$

where $h = (b-a)/n$. Since $M = 0.3679$, then solving for n^2 , we obtain

$$n^2 \geq 30658.3333, \quad \text{gives } n \geq 175.0952.$$

Hence to get the required accuracy, we need 176 subintervals or 177 points. •

Simple Simpson's Rule



Let us consider the second-degree Lagrange interpolating polynomial, with equally spaced base points, that is, $x_0 = a$, $x_1 = a + h$ and $x_2 = a + 2h$, with $h = (b - a)/2$, then

$$f(x) \approx p_2(x) = \frac{(x - x_1)(x - x_2)}{(x_0 - x_1)(x_0 - x_2)}f(x_0) + \frac{(x - x_0)(x - x_2)}{(x_1 - x_0)(x_1 - x_2)}f(x_1) + \frac{(x - x_0)(x - x_1)}{(x_2 - x_0)(x_2 - x_1)}f(x_2).$$

Taking integral on both sides of the above equation with respect to x between the limits x_0 and x_2 , we have

$$\begin{aligned} \int_{x_0}^{x_2} f(x)dx &\approx \frac{f(x_0)}{(x_0 - x_1)(x_0 - x_2)} \int_{x_0}^{x_2} (x - x_1)(x - x_2)dx + \frac{f(x_1)}{(x_1 - x_0)(x_1 - x_2)} \int_{x_0}^{x_2} (x - x_0)(x - x_2)dx \\ &+ \frac{f(x_2)}{(x_2 - x_0)(x_2 - x_1)} \int_{x_0}^{x_2} (x - x_0)(x - x_1)dx, \approx \frac{f(x_0)}{2h^2} I_1 + \frac{f(x_1)}{-h^2} I_2 + \frac{f(x_2)}{2h^2} I_3, \end{aligned}$$

where

$$I_1 = \int_{x_0}^{x_2} (x - x_1)(x - x_2)dx = \frac{2h^3}{3},$$

$$I_2 = \int_{x_0}^{x_2} (x - x_0)(x - x_2)dx; = -\frac{4h^3}{3},$$

$$I_3 = \int_{x_0}^{x_2} (x - x_0)(x - x_1)dx = \frac{2h^3}{3}.$$

By using these values, we have

$$\int_a^b f(x)dx \approx \frac{f(x_0)}{2h^2} \left(\frac{2h^3}{3} \right) + \frac{f(x_1)}{-h^2} \left(\frac{-4h^3}{3} \right) + \frac{f(x_2)}{2h^2} \left(\frac{2h^3}{3} \right).$$

Simplifying, gives

$$\int_a^b f(x)dx \approx S_2(f) = \frac{h}{3}[f(x_0) + 4f(x_1) + f(x_2)].$$

which is called the *simple Simpson's rule* or Simpson's rule for two strips (or 3 points).

Example 5.22 Approximate the following integral

$$\int_1^2 \frac{1}{x+1} dx,$$

using simple Simpson's rule. Compute the actual error.

Solution. Since $f(x) = \frac{1}{x+1}$ and $h = (2 - 1)/2 = 0.5$, then by using Simpson's rule (5.37), we have

$$S_2(f) = \frac{0.5}{3} [f(1) + 4f(1.5) + f(2)] = (0.1667)[0.5 + 1.6 + 0.3333] = 0.4056.$$

Hence

$$\int_1^2 \frac{1}{x+1} dx \approx S_2(f) = 0.4056.$$

Since the exact solution of the given integral is, 0.4055, therefore, the actual error is

$$E_{S_2} = I(f) - S_2(f) = -0.0001.$$

To compare this error with the error got by using the simple Trapezoidal rule, the error in Simpson's rule is much smaller than for the Trapezoidal rule by a factor of about 123, a significant increase in accuracy. ●

Theorem 5.4 (Composite Simpson's Rule)

Let $f \in C^4[a, b]$, n be even, $h = (b - a)/n$, and $x_i = a + ih$ for each $i = 0, 1, 2, \dots, n$. Then the composite Simpson's rule for n subintervals can be written as

$$\int_a^b f(x)dx \approx S_n(f) = \frac{h}{3} \left[f(a) + 2 \sum_{i=1}^{n/2-1} f(x_{2i}) + 4 \sum_{i=1}^{n/2} f(x_{2i-1}) + f(b) \right]. \quad (5.38)$$

Proof. Since for the composite form of the Simpson's rule, the interval is divided into n equal subintervals of width h so that $h = \frac{b-a}{n}$. For this rule to work, n must be even number and the total number of $(n+1)$ distinct points $a = x_0 < x_1 < x_2 \dots < x_n = b$ should be odd. The total integral can be represented as

$$\int_{x_0}^{x_n} f(x)dx = \int_{x_0}^{x_2} f(x)dx + \int_{x_2}^{x_4} f(x)dx + \dots + \int_{x_{n-2}}^{x_n} f(x)dx.$$

Substitute the simple Simpson's rule (5.37) for the individual integral yields

$$\begin{aligned} \int_{x_0}^{x_n} f(x)dx &\approx \frac{h}{3}[f(x_0) + 4f(x_1) + f(x_2)] + \frac{h}{3}[f(x_2) + 4f(x_3) + f(x_4)] \\ &+ \dots + \frac{h}{3}[f(x_{n-2}) + 4f(x_{n-1}) + f(x_n)]. \end{aligned}$$

To avoid repetition of terms, we summed them. Note that each of the odd interior point is counted four and so has a coefficient of four whereas each of the even interior point is counted two and so has a coefficient of two. Endpoints are counted once and so has a coefficient one. •

Example 5.27 Evaluate the integral $\int_0^1 e^{4x} dx$ by using the Simpson's rule with $n = 2, 4, 8$. Also, compute the corresponding actual errors.

Solution. For $n = 2$, using the formula (5.37) and $h = 0.5$, we have

$$S_2(f) = \frac{0.5}{3} [f(0) + 4f(0.5) + f(1)] = 14.1924.$$

For $n = 4$, using the formula (5.38) and $h = 0.25$, we have

$$S_4(f) = \frac{0.25}{3} [f(0) + 4[f(0.25) + f(0.75)] + 2f(0.5) + f(1)] = 13.4659.$$

For $n = 8$, using the formula (5.38) and $h = 0.125$, we have

$$S_8(f) = \frac{0.125}{3} [f(0) + 4[f(0.125) + f(0.375) + f(0.625) + f(0.875)] + 2[f(0.25) + f(0.5) + f(0.75)] + f(1)] = 13.4041.$$

Note that the exact value of the given integral is 13.39995, and so the corresponding errors are, 0.79245, 0.06595, and 0.00411 respectively, which decrease by a factor of about 16 at each stage. •

Error Terms for Simpson's Rule

Now we discuss the local error and the global error formulas for Simpson's rule.

Theorem 5.5 (Error Term for Simple Simpson's Rule)

Let $f \in C^4[a, b]$, and $h = (b - a)/2$. The local error that the Simpson's rule makes in estimating the definite integral (5.21) is

$$E_{S_2}(f) = -\frac{h^5}{90}f^{(4)}(\eta(x)), \quad \text{where } \eta(x) \in (a, b).$$

Example 5.28 Compute the local error for the Simpson's rule using the following integral

$$\int_1^2 \frac{1}{x+1} dx.$$

Solution. Given $f(x) = \frac{1}{x+1}$, and $[a, b] = [1, 2]$, then the fourth derivative of the function can be obtain as

$$f' = \frac{-1}{(x+1)^2}, \quad f'' = \frac{2}{(x+1)^3}, \quad f''' = \frac{-6}{(x+1)^4}, \quad f^{(4)} = \frac{24}{(x+1)^5}.$$

$$|E_{S_2}(f)| = \left| -\frac{h^5}{90} \left| f^{(4)}(\eta(x)) \right| \right|, \quad \text{for } \eta(x) \in (1, 2).$$

$$M = \max_{1 \leq x \leq 2} \left| \frac{24}{(x+1)^5} \right| = 0.75. \quad \text{we get} \quad |E_{S_2}(f)| \leq \frac{(0.03125)}{90}(0.75) = 0.0003.$$

Comparing this with the actual error -0.0001 , this bound is about 3 times the actual error.

Error Term for Composite Simpson's Rule

$$E_{S_n}(f) = -\frac{h^5}{90}f^{(4)}(\eta_1(x)) - \frac{h^5}{90}f^{(4)}(\eta_2(x)) - \cdots - \frac{h^5}{90}f^{(4)}(\eta_{n/2}(x)),$$

$$E_{S_n}(f) = -\frac{(b-a)}{180}h^4f^{(4)}(\eta(x)), \quad \text{for } \eta(x) \in (a, b) \text{ and } nh = b - a.$$

is known as the *global error* of the Simpson's rule.

Example 5.30 Consider the integral $I(f) = \int_1^2 \ln(x+1)dx$; $n = 6$.

- Find the approximation of the give integral using the composite Simpson's rule.
- Compute the error bound for the approximation using the formula (5.57).
- Compute the absolute error.
- How many subintervals approximate the given integral to an accuracy of at least 10^{-4} using the composite Simpson's rule ?

Solution. (a) Given $f(x) = \ln(x+1)$, $n = 6$, and so $h = \frac{2-1}{6} = \frac{1}{6}$, then the composite Simpson's rule (5.38) for $n = 6$, can be written as

$$\begin{aligned} \int_1^2 \ln(x+1)dx \approx S_6(f) &= \frac{1/6}{3} \left[\ln(1+1) + 4\left(\ln\left(\frac{7}{6}+1\right) + \ln\left(\frac{9}{6}+1\right) + \ln\left(\frac{11}{6}+1\right)\right) \right] \\ &+ \left[2\left(\ln\left(\frac{8}{6}+1\right) + \ln\left(\frac{10}{6}+1\right)\right) + \ln(2+1) \right]. \\ &= \frac{1}{18} \left[0.6932 + 4(2.7309) + 2(1.8281) + 1.0986 \right] = 0.9095. \end{aligned}$$

(b) Since the fourth derivative of the function is

$$f^{(4)}(x) = \frac{-6}{(x+1)^4}.$$

Since $\eta(x)$ is unknown point in $(1, 2)$, therefore, the bound $|f^{(4)}|$ on $[1, 2]$ is

$$M = \max_{1 \leq x \leq 2} |f^{(4)}(x)| = \left| \frac{-6}{(x+1)^4} \right| = 6/16 = 0.375.$$

Thus the error formula (5.57) becomes

$$|E_{T_6}(f)| \leq \frac{(1/6)^4}{180}(0.375) = 0.000002,$$

which is the possible maximum error in our approximation in part (a).

(c) The absolute error $|E|$ in our approximation is given as

$$|E| = |3 \ln 3 - 2 \ln 2 - 1 - S_6(f)| = 0.0000003.$$

(d) To find the minimum subintervals for the given accuracy, we use the error formula (5.57) which is

$$|E_{S_n}(f)| \leq \frac{(b-a)^5}{180n^4} M \leq 10^{-4}.$$

Since we know $M = 0.375$, then we have

$$n^4 \geq 20.83333, \quad \text{gives } n \geq 2.136435032.$$

Hence to get the required accuracy, we need 4 subintervals (because n should be even) that ensures the stipulated accuracy. •