

Chapter(11)

Chi-Square Tests

(Examples)

The Chi-Square distribution

1. Chi-Square Test for the difference between two proportions.
2. Chi-Square Test for the difference among more than two proportions.
3. Chi-Square Test of independent.

The properties of the chi square distribution.

- χ^2_v Is continuous distribution.
- χ^2_v Positive skewed curve (skewed to the right curve).
- χ^2_v It is not symmetric curve.

1- Chi-Square Test for the difference between two proportions.

H_0 : The two proportions should be the same

H_1 : The two proportions should not be the same

Example (1):

Suppose we examine a sample of 300 children Left-Handed vs. Gend. Sample results organized in a contingency table:

Gender	Hand Preference		
	Left	Right	Total
Female	12	108	120
Male	24	156	180
Total	36	264	300

Test where the Proportion of females who are left handed is equal to the proportion of males who are left handed ($\alpha=0.05$).


Solution

Step (1): $H_0: \pi_1 = \pi_2$ (Proportion of females who are left handed is equal to the proportion of males who are left handed)

$H_1: \pi_1 \neq \pi_2$ (The two proportions are not the same)

If H_0 is true, then the proportion of left-handed females should be the same as the proportion of left-handed males

Step (2):



$$\chi^2_{(r-1)(c-1),\alpha} = \chi^2_{(2-1)(2-1),0.05} = \chi^2_{1 \times 1,0.05} = \chi^2_{1,0.05} = 3.841$$

Degree of freedom = $(2-1)(2-1) = 1$

The shaded area is equal to α for $\chi^2 = \chi^2_{\alpha}$.

df	$\chi^2_{.995}$	$\chi^2_{.990}$	$\chi^2_{.975}$	$\chi^2_{.950}$	$\chi^2_{.900}$	$\chi^2_{.100}$	$\chi^2_{.050}$	$\chi^2_{.025}$	$\chi^2_{.010}$	$\chi^2_{.005}$
①	0.000	0.000	0.001	0.004	0.016	2.706	③.841	5.024	6.635	7.879
2	0.010	0.020	0.051	0.103	0.211	4.605	5.991	7.378	9.210	10.597

Step (3):

Decision Rule:

If

$$\chi^2_{stat} > 3.841$$

Reject H_0 , otherwise, do not reject H_0

Step (4):

	Hand Preference		
Gender	Left	Right	
Female	12 14.4	108 105.6	120
Male	24 21.6	156 158.4	180
	36	264	300

where:

f_o = observed frequency in a particular cell

f_e = expected frequency in a particular cell if H_0 is true

$$f_e = \frac{\text{row total} \times \text{column total}}{n}$$

For example: $E_{11} = \frac{120 \times 36}{300} = 14.4$, and so on.

$$\begin{aligned} \chi_{STAT}^2 &= \sum_{\text{all cells}} \frac{(f_o - f_e)^2}{f_e} \\ &= \frac{(12 - 14.4)^2}{14.4} + \frac{(108 - 105.6)^2}{105.6} + \frac{(24 - 21.6)^2}{21.6} + \frac{(156 - 158.4)^2}{158.4} = 0.7576 \end{aligned}$$

Step (5):

So we do not reject H_0 and conclude that there is not sufficient evidence that the two proportions are different at $\alpha = 0.05$

Example (2):

The following table is the contingency table for the hotel guest satisfaction study. The contingency table has two rows, indicating whether the guest would return to the hotel or would not return to the hotel, and two columns, one for each hotel. The cells in the table indicate the frequency of each row-and-column combination. The row totals indicate the number of guests who would return to the hotel and the number of guests who would not return to the hotel. The column totals are the sample sizes for each hotel location.

Choose Hotel Again	Hotel		
	Beachcomber	Windsurfer	Total
Yes	163	154	317
No	64	108	172
Total	227	262	489


Test where the Proportion of guests who would return Beachcomber, π_1 , is equal to the population proportion of guests who would return to the Windsurfer, π_2 , you can use the Chi-Square test for the difference between two proportion. ($\alpha=0.05$)

Solution**Step (1):**

$H_0: \pi_1 = \pi_2$ (There is no difference between the two population Proportion)

$H_1: \pi_1 \neq \pi_2$ (The population Proportion are not the same)

If H_0 is true, there is no difference between the proportions of guests who are likely to choose either of these hotels again

Step (2):

$$\chi^2_{(r-1)(c-1),\alpha} = \chi^2_{(2-1)(2-1),0.05} = \chi^2_{1 \times 1,0.05} = \chi^2_{1,0.05} = 3.841$$

$$\text{Degree of freedom} = (2-1)(2-1) = 1$$

The shaded area is equal to α for $\chi^2 = \chi^2_{\alpha}$.

df	$\chi^2_{.995}$	$\chi^2_{.990}$	$\chi^2_{.975}$	$\chi^2_{.950}$	$\chi^2_{.900}$	$\chi^2_{.100}$	$\chi^2_{.050}$	$\chi^2_{.025}$	$\chi^2_{.010}$	$\chi^2_{.005}$
①	0.000	0.000	0.001	0.004	0.016	2.706	3.841	5.024	6.635	7.879
2	0.010	0.020	0.051	0.103	0.211	4.605	5.991	7.378	9.210	10.597

Step (3):

Decision Rule:

If

$$\chi^2_{\text{stat}} > 3.841$$

Reject H_0 , otherwise, do not reject H_0

Step (4):

Choose Hotel Again	Hotel		
	Beachcomber	Windsurfer	Total
Yes	163 147.16	154 169.84	317
No	64 79.84	108 92.16	172
Total	227	262	489

where:

f_o = observed frequency in a particular cell

f_e = expected frequency in a particular cell if H_0 is true

$$f_e = \frac{\text{row total} \times \text{column total}}{n}$$

For example: $E_{11} = \frac{317 \times 227}{489} = 147.16$, and so on.

$$\begin{aligned} \chi^2_{\text{STAT}} &= \sum_{\text{all cells}} \frac{(f_o - f_e)^2}{f_e} \\ &= \frac{(163 - 147.16)^2}{147.16} + \frac{(154 - 169.84)^2}{169.84} + \frac{(64 - 79.84)^2}{79.84} + \frac{(108 - 92.16)^2}{92.16} = 9.048 \end{aligned}$$

Step (5):

We reject H_0 and conclude that proportion of guests who would return to Beachcomber is different from the proportion of guests who would return to the Windsurfer

2- Chi-Square Test for the difference among more than two proportions.

H0: The proportions should be the same

H1: The proportions should not be the same

Example (3):

Most companies consider big data analytics critical to success .However, is there a difference among small (<100 employees), mid-size (100-999 employees), and large (1000+ employees) companies in the proportion of companies that have already deployed big data project? A study showed the results for the different company size. (Data extracted from 2014 big data outlook:big data is transformative –where is your company?)

Have already deployed big data projects	Company Size			
	Small	Mid-sized	large	Total
Ye	18	74	52	144
No	182	126	148	456
Total	200	200	200	600

Assume that 200 decision makers involved in big data purchases within each company size were surveyed.

At the 0.05 level of significance, is there evidence of a difference among companies of different sizes with respect to the proportion of companies that have already deployed big data projects?

Solution

Step (1):

H₀: $\pi_1 = \pi_2 = \pi_3$


H₁: At least one proportion differs where

π_1 = small, π_2 = medium, π_3 = large

(There is no difference between the three populations Proportion)

If H0 is true, there is no difference between the three proportions.

Step (2):



$$\chi^2_{(r-1)(c-1),\alpha} = \chi^2_{(2-1)(3-1),0.05} = \chi^2_{2 \times 1,0.05} = \chi^2_{2,0.05} = 5.991$$

Degree of freedom = (2-1)(3-1)=2

The shaded area is equal to α for $\chi^2 = \chi^2_{\alpha}$.

df	$\chi^2_{.995}$	$\chi^2_{.990}$	$\chi^2_{.975}$	$\chi^2_{.950}$	$\chi^2_{.900}$	$\chi^2_{.100}$	$\chi^2_{.050}$	$\chi^2_{.025}$	$\chi^2_{.010}$	$\chi^2_{.005}$
1	0.000	0.000	0.001	0.004	0.016	2.706	3.841	5.024	6.635	7.879
2	0.010	0.020	0.051	0.103	0.211	4.605	5.991	7.378	9.210	10.597
3	0.072	0.115	0.216	0.352	0.584	6.251	7.815	9.348	11.345	12.838
4	0.207	0.297	0.484	0.711	1.064	7.779	9.488	11.143	13.277	14.860
5	0.412	0.554	0.831	1.145	1.610	9.236	11.070	12.833	15.086	16.750

Step (3):

Decision Rule:

If

$$\chi^2_{stat} > 5.991$$

Reject H0, otherwise, do not reject H0

Step (4):

Have already deployed big data projects	Company Size			
	Small	Mid-sized	large	Total
Ye	18	74	52	144
No	182	126	148	456
Total	200	200	200	600

where:

f_o = observed frequency in a particular cell

f_e = expected frequency in a particular cell if H0 is true

$$f_e = \frac{\text{row total} \times \text{column total}}{n}$$

For example: $E_{11} = \frac{144 \times 200}{600} = 48$, and so on.

$$\begin{aligned} \chi^2_{STAT} &= \sum_{\text{all cells}} \frac{(f_o - f_e)^2}{f_e} \\ &= \frac{(18-48)^2}{48} + \frac{(74-48)^2}{48} + \frac{(52-48)^2}{48} + \frac{(182-152)^2}{152} + \frac{(126-152)^2}{152} + \frac{(148-152)^2}{152} = 43.64035 \end{aligned}$$

Step (5):

Since $\chi^2_{stat} = 43.64035$ is greater than the upper critical value of 5.991, reject H0
 .There is evidence among the groups with respect to the proportion of companies that have already deployed big data projects

3- Chi-Square Test of independent.

The test is applied when you have two categorical variables from a single population, and it is used to determine whether there is a significant association between the two variables.

H0: The two categorical variables are independent
(i.e., there is no relationship between them)

H1: The two categorical variables are dependent
(i.e., there is a relationship between them)

Example (4):

The meal plan selected by 200 students is shown below:

Class Standing	Number of meals per week			Total
	20/week	10/week	none	
Fresh.	24	32	14	70
Soph.	22	26	12	60
Junior	10	14	6	30
Senior	14	16	10	40
Total	70	88	42	200

At the 0.05 level of significance, is there evidence that meal plan and class standing are independent(i.e., there is no relationship between them)


Solution

Step (1):

H0: Meal plan and class standing are independent
(i.e., there is no relationship between them)

H1: Meal plan and class standing are dependent
(i.e., there is a relationship between them)

Step (2):



$$\chi^2_{(r-1)(c-1),\alpha} = \chi^2_{(4-1)(3-1),0.05} = \chi^2_{3 \times 2,0.05} = \chi^2_{6,0.05} = 12.592$$

Degree of freedom= (4-1)(3-1)=6

The shaded area is equal to α for $\chi^2 = \chi^2_{\alpha}$.

df	$\chi^2_{.995}$	$\chi^2_{.990}$	$\chi^2_{.975}$	$\chi^2_{.950}$	$\chi^2_{.900}$	$\chi^2_{.100}$	$\chi^2_{.050}$	$\chi^2_{.025}$	$\chi^2_{.010}$	$\chi^2_{.005}$
1	0.000	0.000	0.001	0.004	0.016	2.706	3.841	5.024	6.635	7.879
2	0.010	0.020	0.051	0.103	0.211	4.605	5.991	7.378	9.210	10.597
3	0.072	0.115	0.216	0.352	0.584	6.251	7.815	9.348	11.345	12.838
4	0.207	0.297	0.484	0.711	1.064	7.779	9.488	11.143	13.277	14.860
5	0.412	0.554	0.831	1.145	1.610	9.236	11.070	12.833	15.086	16.750
6	0.676	0.872	1.237	1.635	2.204	10.645	12.592	14.449	16.812	18.548
7	0.989	1.239	1.690	2.167	2.833	12.017	14.067	16.013	18.475	20.278

Step (3):

Decision Rule: If

$$\chi^2_{\text{stat}} > 12.592$$

Reject H_0 , otherwise, do not reject H_0

Step (4):

Class Standing	Number of meals per week			Total
	20/week	10/week	none	
Fresh.	24 24 24.5	32 32 30.8	14 14 14.8	70
Soph.	22 22 21	26 26 26.4	12 12 12.6	60
Junior	10 10 10.5	14 14 13.2	6 6 6.3	30
Senior	14 14 14	16 16 17.6	10 10 8.4	40
Total	70	88	42	200

where:

f_o = observed frequency in a particular cell

f_e = expected frequency in a particular cell if H_0 is true

$$f_e = \frac{\text{row total} \times \text{column total}}{n}$$

For example: $E_{11} = \frac{70 \times 70}{200} = 24.5$, and so on.

$$\begin{aligned} \chi^2_{\text{STAT}} &= \sum_{\text{all cells}} \frac{(f_o - f_e)^2}{f_e} \\ &= \frac{(24 - 24.5)^2}{24.5} + \frac{(32 - 30.8)^2}{30.8} + \frac{(14 - 14.8)^2}{14.8} + \frac{(22 - 21)^2}{21} + \frac{(26 - 26.4)^2}{26.4} + \frac{(12 - 12.6)^2}{12.6} \\ &\quad + \frac{(10 - 10.5)^2}{10.5} + \frac{(14 - 13.2)^2}{13.2} + \frac{(6 - 6.3)^2}{6.3} + \frac{(14 - 14)^2}{14} + \frac{(16 - 17.6)^2}{17.6} + \frac{(10 - 8.4)^2}{8.4} = 0.709 \end{aligned}$$

Step (5):

Here,

$$\chi^2_{\text{stat}} = 0.709 < \chi^2_{0.05} = 12.592$$

so do not reject H_0

Conclusion: there is not sufficient evidence that meal plan and class standing are related at $\alpha = 0.05$

Example (5):


Suppose you conducted a drug trial on a group of animals and you hypothesized that the animals receiving the drug would show increased heart rates compared to those that did not receive the drug. You conduct the study and collect the following data($\alpha=0.05$):

	Heart Rate Increased	No Heart Rate Increase	Total
Treated	36	14	50
Not treated	30	25	55
Total	66	39	105

Solution:**Step (1):**

H0: The proportion of animals whose heart rate increased is independent of drug treatment.

H1: The proportion of animals whose heart rate increased is associated with drug treatment.

Step (2):

$$\chi^2_{(r-1)(c-1),\alpha} = \chi^2_{(2-1)(2-1),0.05} = \chi^2_{1 \times 1,0.05} = \chi^2_{1,0.05} = 3.841$$

Degree of freedom= (2-1)(2-1)=1

The shaded area is equal to α for $\chi^2 = \chi^2_{\alpha}$.

df	$\chi^2_{.995}$	$\chi^2_{.990}$	$\chi^2_{.975}$	$\chi^2_{.950}$	$\chi^2_{.900}$	$\chi^2_{.100}$	$\chi^2_{.050}$	$\chi^2_{.025}$	$\chi^2_{.010}$	$\chi^2_{.005}$
①	0.000	0.000	0.001	0.004	0.016	2.706	③.841	5.024	6.635	7.879
2	0.010	0.020	0.051	0.103	0.211	4.605	5.991	7.378	9.210	10.597
...

Step (3):

Decision Rule:

If

$$\chi^2_{\text{stat}} > 3.841$$

Reject H_0 , otherwise, do not reject H_0

Step (4):

	Heart Rate Increased	No Heart Rate Increase	Total
Treated	36 31.4	14 18.6	50
Not treated	30 35.6	25 20.4	55
Total	66	39	105

where:

f_o = observed frequency in a particular cell

f_e = expected frequency in a particular cell if H_0 is true

$$f_e = \frac{\text{row total} \times \text{column total}}{n}$$

For example: $E_{11} = \frac{50 \times 66}{105} = 31.4$, and so on.

$$\begin{aligned}\chi^2_{\text{STAT}} &= \sum_{\text{all cells}} \frac{(f_o - f_e)^2}{f_e} \\ &= \frac{(36 - 31.4)^2}{31.4} + \frac{(14 - 18.6)^2}{18.6} + \frac{(30 - 35.6)^2}{35.6} + \frac{(25 - 20.4)^2}{20.4} = 3.42\end{aligned}$$

Step (5):

Our calculated value (3.42) is less than the tabular value so we cannot reject the null hypothesis and we conclude that; the proportion of animals whose heart rate increased is independent of drug treatment.