

Characterization and gene mapping of a chlorophyll-deficient mutant *clm1* of *Triticum monococcum* L.

M.J. ANSARI^{1,6*}, A. AL-GHAMDI¹, R. KUMAR², S. USMANI³, Y. AL-ATTAL¹, A. NURU¹, A.A. MOHAMED¹, K. SINGH⁴, and H.S. DHALIWAL⁵

Department of Plant Protection, College of Food and Agriculture Sciences, King Saud University, PO Box 2460, Riyadh 11451, King Saud Arabia¹

Department of Biotechnology, ICFAI University, Dehradun-248197, India²

D.K.M. College for Women, Thiruvalluvar University, Vellore, Tamilnadu-632004, India³

School of Agricultural Biotechnology, Punjab Agricultural University, Ludhiana, Punjab-141004, India⁴

Akal School of Biotechnology, Eternal University, Baru Sahib, Himachal Pradesh-173101, India⁵

Department of Biotechnology, Indian Institute of Technology, Roorkee, Uttarakhand-247667, India⁶

Abstract

Diploid wheat *Triticum monococcum* L. is a model plant for wheat functional genomics. Chlorophyll-deficient mutant (*clm1*) was identified during manual screening of the ethylmethane sulphonate (EMS)-treated M₂ progenies of *T. monococcum* accession paul4087 in the field. The *clm1* mutant, due to significantly decreased chlorophyll content compared with the wild-type (WT), exhibited pale yellow leaves which slowly recovered to green before flowering. The *clm1* mutant showed early flowering, reduced number of tillers, trichome length and density, and different shape as compared with the WT. At the same time, *clm1* mutant culm had more chlorophyll-containing parenchymatous tissues compared to WT, presumably to absorb more sunlight for photosynthesis. Genetic analysis indicated that the *clm1* mutation was monogenic recessive. The *clm1* mutant was mapped between Xgwm473 and Xwmc96 SSR markers, with genetic distances of 2.1 and 2.6 cM, respectively, on the 7A^mL chromosome.

Additional key words: bulk segregant analysis, diploid wheat, ethylmethane sulfonate, SSR marker.

Introduction

Chlorophyll (Chl) plays significant role in absorption of sunlight by photosynthetic reaction centers (Liu *et al.* 2007) and approximately 100 terawatts of energy is captured by plants (Neelson and Conrad 1999). Mutations in the genes of Chl biosynthesis or related pathway result in Chl-deficient mutants or leaf-color mutants. These mutants are ideal materials for fundamental research in photosynthesis, photomorphogenesis, hormone physiology, resistance mechanism, and identification of gene functions (Parks and Quail 1991, Mochizuki *et al.* 2001, Stern *et al.* 2004, Beale 2005). The chemical mutagen, ethylmethane sulfonate (EMS), has been successfully applied in wheat and rice (Wang *et al.* 2009, Ansari *et al.* 2012, Tian *et al.* 2012).

The diploid wheat (*T. monococcum*, A^mA^m) is an ideal

material for induced mutations which can be easily characterized and transferred to polyploid wheat. Certain characteristics of *T. monococcum* make it an attractive diploid model for gene discovery in wheat (Wicker *et al.* 2001) and application of several functional genomics approaches. Firstly, the diploid *T. monococcum* has a small genome size (5 700 Mb) compared with hexaploid wheat (17 300 Mb). Secondly, the existence of a very high level of polymorphism for DNA based markers, conservation of colinearity and synteny with other cereal crops, and high resistance against various wheat diseases are the other attractive features. Finally, a large bacterial artificial chromosome (BAC) library is available in *T. monococcum*.

Several Chl-deficient mutants have been identified

Received 11 October 2011, accepted 24 August 2012.

Abbreviations: BSA - bulk segregant analysis; Chl - chlorophyll; *clm1* - chlorophyll-deficient mutant; EMS - ethylmethane sulfonate; M₂ - second generation after mutagenesis; PCR - polymerase chain reaction; RILs - recombinant inbred lines; SEM - scanning electron microscopy; T.S. - transverse section; WT - wild-type.

Acknowledgments: The authors are grateful to Deanship of Scientific Research and College of Food & Agriculture Science Research, King Saud University, Riyadh as well as to Council of Scientific and Industrial Research, New Delhi, Government of India, for providing research support.

* Author for correspondence; fax: (+966) 1 4679065, e-mail: mjavedansari@gmail.com

in wheat (Freeman *et al.* 1982, Falbel and Staehelin 1994, Guo *et al.* 1996), rice (Kaul and Bhan 1977, Wu *et al.* 2007, Chen *et al.* 2009), barley (Dunford and Walden 1991), soybean (Zhang *et al.* 2011), *Arabidopsis* (Long *et al.* 1993, Hsieh and Goodman 2005, Kim *et al.* 2006), and tobacco (Chang *et al.* 2001). In the hexaploid wheat, *T. aestivum*, some Chl mutants like Driscoll's *chlorina* (Pettigrew *et al.* 1969) and *chlorina-1* (Sears and Sears 1968) have been mapped on the A genome, and *Chlorina-214* (Washington and Sears 1970) on the B genome. These wheat mutants exhibit reduced Chl *b* content, a significant reduction in light harvesting complex, and formation of grana-deficient thylakoid membranes (Falbel *et al.* 1996). *CD3*, a non-lethal Chl-deficient mutant, was also identified in hexaploid wheat (Freeman *et al.* 1982). Some orthologous *chlorina* mutants similar to *CD3* have been mapped to chromosome 7 of the A, B, and D parental diploid wheat

genomes in both hexaploid (*T. aestivum*) and tetraploid (*T. turgidum*) wheat (Falbel and Staehelin 1994).

The rapid development of new simple sequence repeat (SSR) molecular markers has been increasingly exploited to develop high-density genetic maps in diploid wheat, *T. monococcum* (Dubcovsky *et al.* 1996, Singh *et al.* 2007). The first sequence coverage of the wheat genome (*T. aestivum*, Chinese Spring line 42) has been released and is available via EMBL/GenBank and CerealsDB for genomic analysis and application (<http://www.cerealsdb.uk.net>).

This article deals with the characterization and mapping of a novel, EMS induced, Chl deficient mutant (*clm1*) in diploid wheat, *T. monococcum*. The results will not only promote map-based cloning of *clm1* gene but also could have far reaching implications and applications in the investigation of photosynthesis and chloroplast development pathway in wheat and other cereals.

Materials and methods

The *clm1* mutant used in the present study was isolated from diploid wheat, *T. monococcum* accession pau14087, at the Punjab Agricultural University, Ludhiana, after seed treatment with 0.25 % EMS. This mutant was identified during manual screening of the M₂ EMS-treated population in the field. The seeds of *clm1*, the WT parent, and an accession pau5088 of *T. boeoticum* (the wild and tall progenitor of *T. monococcum*) were planted at the Indian Institute of Technology (IIT), Roorkee, in November 2005. A recombinant inbred lines (RILs) population of *T. boeoticum* pau5088 × *T. monococcum* pau14087 was also planted in the field (Singh *et al.* 2007) and some RILs were chosen at random to develop a negative bulk without the mutation. The *clm1* mutant was crossed with both the WT parent, *T. monococcum*, and its wild progenitor, *T. boeoticum*, for developing F₂ populations for inheritance and mapping studies, respectively. The F₂ populations were planted at IIT, Roorkee, in 2007 and 2008 in 2 m rows with row-to-row distance of 30 cm and plant to plant distance of 10 cm following the standard package of agronomic practices for wheat cultivation.

The total Chl *a* and *b* content of *clm1* mutant and *T. monococcum* were estimated according to Arnon (1949) and Koski (1950). Fresh leaf tissue was extracted with 85 % (v/v) acetone and absorbance was recorded on a spectrophotometer (*Lambda 25 UV/VIS*, Perkin Elmer, Shelton, USA) at 663 and 644 nm.

The samples for scanning electron microscopy (SEM) were prepared according to Mou *et al.* (2000). The tissue samples were immersed in 2.5 % (v/v) glutaraldehyde at room temperature for 2 h and dehydrated in 50 % (v/v) ethanol for 5 min, in 70 % for 30 min (twice), 90 % for 30 min (twice), and 100 % for 30 min (twice). Finally, in mixture of ethanol and amyl acetate 3:1 for 30 min, 2:2 for 30 min, 1:3 for 30 min, and in pure amyl acetate for 30 min. The samples were kept for critical

point drying for 40 min and mounted onto metal stubs with double-sided carbon tape. For sputter coating, a thin layer of gold was applied over the samples using an automated sputter coater. These samples were then analyzed using scanning electron microscope (*Leo 435*, Cambridge, USA) and the surface images were taken at 150-fold magnification.

For histological examination, the second internode of stems were excised, fixed in formalin + acetic acid + ethanol (FAA), and dehydrated in a graded ethanol series and finally in xylene. For sectioning, the tissues were embedded in paraffin wax (*Sd fine*, Mumbai, India) at 60 °C, and sectioned to 10 µm thickness on a rotary microtome. The tissues were stained with toluidine blue O (0.06 %, m/v) as suggested by Johansen (1940). Transverse sections of the stems were observed under a light microscope (*Axiostar plus 1169-151*, Carl Zeiss, Oberkochen, Germany) at different magnifications.

The genomic DNA from parents and the F₂ population was extracted following the cetyltrimethyl ammonium bromide (CTAB) method as described by Saghai-Marooof *et al.* (1984). A number of high density molecular maps have been developed in wheat (Somers *et al.* 2004) including a RIL population between *T. monococcum* acc. pau14087, used for *clm1* mutant isolation, and *T. boeoticum* acc. pau5088, used for making populations, with the *clm1* mutant (Singh *et al.* 2007). The primers for anchored SSR markers at about 10 cM from each of the diploid wheat chromosomes and polymerase chain reaction (PCR) protocols were carried out in a thermo cycler (*Applied Biosystems*, Singapore) according to methods described by Singh *et al.* (2007). The PCR products were separated on 8 % (m/v) polyacrylamide gels according to the length of the amplified fragments and stained with ethidium bromide.

To study the inheritance of *clm1* mutant, it was crossed with its WT parent *T. monococcum* and

T. boeoticum for molecular mapping. The F₁ plants were advanced to F₂ generations. The phenotypic data on leaf color (chlorotic vs. green) was recorded on each of the plant of the F₂ populations. χ^2 -test was applied to test the goodness of fit to the segregation ratio.

For bulk segregant analysis (BSA), positive bulk of 12 *clm1* mutant plants was made from homozygous mutant F₂ *chlorina* plants from the respective F₂ mapping population of *T. monococcum* × *T. boeoticum*. The negative bulk was prepared from 15 RILs of *T. monococcum* pau5088 × *T. boeoticum* pau14087 cross without any mutant in the parents. An equal amount of DNA from each plant of both bulks was pooled along

with parents. These two sets of bulks along with the parents were used to identify putative SSR markers linked to the *clm1* mutant. Genotyping of the debulked F₂ plants was done with the putatively linked marker (Xgwm473) and two other closely linked markers (Xwmc96 and Xcfd68) as per the diploid wheat map (Singh *et al.* 2007). These three SSR markers were used to genotype individual plants of the F₂ mapping population. A linkage map was constructed with MAPMAKER/EXP v. 3.0 (Lander *et al.* 1987, Lincoln *et al.* 1993) according to the linkage data of the *clm1* loci and polymorphic SSR markers in the F₂ mapping population.

Results

The Chl-deficient mutants (*clm1*) were erect with shorter stems and roots, small flag leaf, lower tiller numbers, less spikelets per spike, and slower growth rate as compared to WT parent *T. monococcum* (Fig. 1A,B). The change in color was observed in the *clm1* mutant in comparison to WT at 14 d after sowing. The *clm1* mutant exhibited yellow-green leaves (14 to 56 d after sowing)

which gradually changed from pale yellow to yellowish green (approximately 70 d after sowing) and became pale green just before flowering (approximately 84 d after sowing). The *clm1* mutant was more sensitive to a herbicide *Topik* as compared to WT.

Chlorophyll content was measured in the *clm1* and WT at 14-d intervals up to 84 d (Table 1).




Fig. 1. Morphological and histological comparison of *T. monococcum* wild type (WT) and its *clm1* mutant: A - field view; B - 25-d-old seedling; C and D - scanning electron microscopy of trichomes on upper leaf surface of WT and *clm1* mutant (45-d-old seedlings (150×)); E and F - transverse section of second internode (toluidine blue staining) of WT and *clm1* mutant (Sc - sclerenchyma, P - parenchyma; OV - outer or peripheral vascular bundles; IV - inner vascular bundle; T - trichome).

Table 1. Chlorophyll content [mg g^{-1} (f.m.)] in the leaves during ontogeny of *T. monococcum* WT and *clm1* mutant (mean \pm SE, $n = 3$).

Ontogeny	WT Chl <i>a</i>	Chl <i>b</i>	Chl <i>a+b</i>	<i>clm1</i> Chl <i>a</i>	Chl <i>b</i>	Chl <i>a+b</i>
Day 14	1.10 \pm 0.02	0.26 \pm 0.016	1.36 \pm 0.02	0.21 \pm 0.01	0.07 \pm 0.009	0.28 \pm 0.01
Day 28	1.82 \pm 0.02	0.38 \pm 0.02	2.20 \pm 0.03	0.31 \pm 0.01	0.11 \pm 0.01	0.42 \pm 0.02
Day 56	1.98 \pm 0.03	0.41 \pm 0.02	2.39 \pm 0.02	0.64 \pm 0.02	0.21 \pm 0.01	0.85 \pm 0.02
Day 84	2.16 \pm 0.03	0.47 \pm 0.02	2.63 \pm 0.03	1.08 \pm 0.02	0.41 \pm 0.01	1.49 \pm 0.02

The *clm1* mutant showed reduced content of total Chl (43.3% to 84.5 % reduction) and an increased Chl *a/b* ratio. In comparison to the WT, the content of Chl *a* and Chl *b* remained significantly reduced throughout the whole *clm1* development with the lowest values at 14 d seedling stage. Increased Chl *a/b* ratio occurred probably because of the relatively lower synthesis of Chl *b* than Chl *a*. However, *clm1* mutant accumulated significant quantities of Chl in the upper leaves to become slightly green at flowering stage.

The morphology and distribution of trichomes in WT and *clm1* mutant varied in length, density, and type. In *clm1* mutant, the number of trichomes was low whereas in the WT they were packed densely. In the leaves of WT, three types of trichomes were found. One type had a round globular base without a pointed tip which occurred near the stomata and over the veins, the second type had a globular base with a short pointed tip and the third type had a globular base with a long pointed tip. All three types were observed over the upper leaf surface although the third type was rare. In the *clm1* mutant, only one type of elongated trichomes with a pointed tip and without a globular base were observed. All the trichomes in the WT and *clm1* were unbranched. A higher density of glandular and globular trichomes was observed over the leaf veins of the WT, whereas no trichomes were observed over the leaf veins in *clm1*. It

indicated the pleiotrophic effect of *clm1* locus which not only affects the Chl content but also plays a significant role in controlling the shape and types of trichomes (Fig. 1C,D).

To determine whether the less Chl accumulation was associated with a change in the histology of *clm1* mutant, transverse sections of the culm of the WT and mutant were stained with toluidine blue. The transverse section revealed that the diameter of the second internode in the WT was larger than that in the *clm1* mutant. Cell density (number of cells per unit area) of the internode was similar in both the *clm1* mutant and WT. The amount of sclerenchymatous tissues in the *clm1* mutant was significantly reduced and only small strips of sclerenchyma were observed attached to the epidermis on the outer side and to vascular bundles towards the inner side in the *clm1* mutant. However, in WT, sclerenchyma formed a thick and dense layer towards the periphery and around the vascular bundles. Large chloroplast-containing parenchyma, which was similar to mesophyll cells of the leaves, was present in the *clm1* mutant probably to absorb more sunlight and to increase the photosynthetic rate. The increased size of the parenchyma was negatively correlated with the reduced amount of Chl in the chloroplast presumably in an effort to compensate for the reduced photosynthetic rate in *clm1*. In WT, few Chl-containing parenchyma tissues were present. In *clm1*,




Fig. 2. *A* - Identification of putative SSR marker Xgwm473 linked with *clm1* mutant through bulk segregant analysis. *B* - PCR amplification of debulks of positive bulks (1 - 12) of F_2 plants using putatively linked marker Xgwm473 identified in BSA and its closely linked marker Xwmc96 (Singh *et al.* 2007).




Fig. 3. The segregation of SSR marker *Xgwm473* in representative 35 F_2 plants of *clm1* \times *T. boeoticum* cross.

peripheral vascular bundles were surrounded by parenchyma from the outer side, whereas in WT, peripheral vascular bundles were completely surrounded by sclerenchyma. (Fig. 1E,F).

The F_1 obtained from the cross of *T. monococcum* \times *clm1* mutant was like the WT parent indicating that the *clm1* mutant was recessive. The segregation for Chl-deficiency was found in the F_2 population. Out of 160 F_2 plants in total, 117 plants were WT and 43 were found Chl-deficient (chlorina) which gave a good fit to 3:1 ratio indicating that the *clm1* mutant was monogenic recessive.

Anchored SSR markers (133) polymorphic between *T. monococcum* and *T. boeoticum* (Singh *et al.* 2007) were used for BSA. The marker *Xgwm473* and *Xwmc96* (not shown in BSA) located on chromosome 7A^{ML} showed polymorphism between the positive and negative bulk for *clm1* (Fig. 2A). Debulking of positive bulk plants also showed close linkage with markers *xgwm473* and *Xwmc96* in the *clm1* mutant. Genotyping of the debulked F_2 plants with the putatively linked markers from BSA confirmed their close linkage with *clm1* mutant (Fig. 2B). To map the *clm1* mutant, F_2 mapping population developed by crossing the *clm1* mutant with *T. boeoticum* acc. pau 5088, a non-chlorina and tall wild progenitor of *T. monococcum*, was used. A total of 160 individual F_2 plants were used for genotyping of *clm1* F_2 populations (Fig. 3). Co-segregation analysis of individual markers, *Xgwm473*, *Xwmc96*, and *Xcfd68* located on chromosome 7A^{ML} and *clm1* using the marker and mutant genotypes of F_2 plants, was carried out with the help of recombination frequency among markers at each locus. Based on the data on recombination frequency, the marker *Xgwm473*, *Xwmc96*, and *Xcfd68* were mapped at distances of 2.1, 2.6, and 4.6 cM, respectively, from the *clm1* gene on chromosome 7A^{ML} (Fig. 4).




Fig. 4. Mapping of *clm1* mutant on chromosome 7AL of *T. monococcum* (Singh *et al.* 2007).

Discussion

The EMS-induced Chl-deficient mutant *clm1* was used for phenotypic and molecular characterization in the present study. The *clm1* mutant was found to be controlled by a single recessive gene. Several Chl-deficient mutants have been identified in tetraploid and hexaploid wheat and mapped on A, B, and D genomes (Sears and Sears 1968, Pettigrew *et al.* 1969, Washington and Sears 1970, Freeman *et al.* 1987, Falbel and Staehelin 1994, Guo *et al.* 1996). However, no Chl-defi-

cient mutant has been reported in diploid wheat. The *clm1* mutant exhibits a yellow green leaf phenotype and reduced content of Chl *a* and *b* at the seedling stage. The Chl content reached maximum of 67 % of the WT, *T. monococcum*, just before flowering and the upper leaves of the *clm1* mutant ultimately turned green. This suggests that the *clm1* mutant exhibited delayed greening during photomorphogenesis because of a slower rate of Chl accumulation. The *clm1* mutation affected Chl

synthesis most drastically in the early developmental stages and recovered in the later stage during flowering suggesting that it may be a development stage specific or a conditional mutant. These results were consistent with those of Wu *et al.* (2007), Du *et al.* (2009) and Zhang *et al.* (2011).

Three types of trichomes were observed in the WT. However, elongated trichomes with pointed tip and without a globular base clearly different from the trichomes present in the *T. monococcum* were present in the *clm1* mutant. This indicated that *clm1* mutation not only affected the Chl content but also had a pleiotropic effect on leaf trichome morphology, type, and distribution. The transverse section of second internode revealed an increase in the quantity of Chl-containing parenchyma in *clm1* mutant. Several Chl-deficient mutants have been reported in tetraploid and hexaploid wheats. All these mutants exhibited reduced Chl *b* content, a significant reduction in light harvesting CP complex, and formation of grana-deficient thylakoid membranes (Falbel *et al.* 1996). Some of these mutants have been mapped to chromosome 7 A, B, or D genomes in both hexaploid (*T. aestivum*) and tetraploid (*T. turgidum*) wheat (Falbel and Staehelin 1994).

In the F₂ population between the *clm1* mutant and its WT, *T. monococcum*, WT phenotype and the mutant showed a segregation ratio of 3:1 (green : yellow plants) indicating that the *clm1* mutant was controlled by a single recessive nuclear gene. Such mutants probably could not have been recovered in polyploid wheat because of their

orthologous loci on other genomes unless multiple mutants were induced at all the loci or certain loci were silenced during evolution. The *clm1* mutant has been mapped to chromosome 7A^m of diploid wheat, *T. monococcum*, which is similar to the findings previously reported for Chl-deficient mutants, such as CD3 in hexaploid wheat on the homoeologous chromosome 7 (Falbel and Staehelin 1996). Similar mutants were obtained in rice (Huang *et al.* 2008), barley (Liu *et al.* 2008), and soybean (Zhang *et al.* 2011). The photosynthetic metabolic pathway is very complex, and many genes have remained undiscovered. Several QTLs and genes related to Chl metabolic pathway have been mapped in rice (Chen *et al.* 2007, 2009, Shen *et al.* 2007, Huang *et al.* 2008, Jiang *et al.* 2008, Wang *et al.* 2009, Li *et al.* 2010, Sang *et al.* 2010), and soybean (Zou *et al.* 2003, Zhang *et al.* 2011). Map-based cloning was done for yellow green leaf 1 (*yg1l*), a Chl-deficient recessive mutant reported in rice with yellow green leaves, decreased Chl content, and delayed chloroplast development in young plants (Wu *et al.* 2007). The *clm1* mutant also had reduced Chl content and inheritance similar to *yg1l*.

Chl deficiency is one of the most important agronomic traits that affect not only grain production but also the usefulness of cereal straws as animal forage. As an important locus regulating wheat Chl synthesis, knowledge of *clm1* (and its orthologs in other cereals) could make a significant contribution to the future improvement of wheat for enhanced Chl content and photosynthetic rate.

References

- Ansari, M.J., Kumar, R., Singh, K., Dhaliwal, H.S.: Characterization and molecular mapping of EMS-induced brittle culm mutants of diploid wheat (*Triticum monococcum* L.). - *Euphytica* **186**: 165-176, 2012.
- Arnon, D.I.: Copper enzymes in isolated chloroplast: polyphenols-oxidase in *Beta vulgaris*. - *Plant Physiol.* **24**: 1-15, 1949.
- Beale, S.I.: Green genes gleaned. - *Trends Plant Sci.* **10**: 309-312, 2005.
- Chang, H.B., Abe, T., Matsuyama, T., Fukunishi, N., Nagata, N., Nakano, T., Kaneko, Y., Miyoshi, K., Matsushima, H., Yoshida, S.: Regulation of chloroplast gene expression is affected in *ali*, a novel tobacco albino mutant. - *Ann. Bot.* **88**: 545-553, 2001.
- Chen, T., Zhang, Y.D., Zhao, L., Zhu, Z., Lin, J., Zhang, S.B., Wang, C.L.: Physiological character and gene mapping in a new green-revertible albino mutant in rice. - *J. Genet. Genomics* **34**: 331-338, 2007.
- Chen, T., Zhang, Y.D., Zhao, L., Zhu, Z., Lin, J., Zhang, S.B., Wang, C.L.: Fine mapping and candidate gene analysis of a green-revertible albino gene *gra(t)* in rice. - *J. Genet. Genomics* **36**: 117-123, 2009.
- Du, P., Ling, Y.H., Sang, X.C., Zhao, F.M., Xie, R., Yang, Z.L., He, G.H.: Gene mapping related to yellow green leaf in a mutant line in rice. - *Genes Genomics* **31**: 165-171, 2009.
- Dubcovsky, J., Luo, M.C., Zhang, G.Y., Bainsteitter, R., Desai, A., Kilian, A., Kleinhofs, A., Dvorak, J.: Genetic map of diploid wheat *T. monococcum* L. and its comparison with maps of *H. vulgare* L. - *Genetics* **143**: 983-999, 1996.
- Dunford, R., Walden, R.M.: Plastid genome structure and plastid-related transcript levels in albino barley plants derived from anther culture. - *Curr. Genet.* **20**: 339-347, 1991.
- Falbel, T.G., Meehl, J.B., Staehelin, A.: Severity of mutant phenotype in a series of chlorophyll deficient wheat mutants depends on light intensity and the severity of the block in chlorophyll synthesis. - *Plant Physiol.* **112**: 821-832, 1996.
- Falbel, T.G., Staehelin, L.A.: Characterization of a family of chlorophyll-deficient wheat and barley mutants with defects in the Mg-insertion step of chlorophyll biosynthesis. - *Plant Physiol.* **104**: 639-648, 1994.
- Falbel, T.G., Staehelin, L.A.: Partial block in the early steps of the chlorophyll synthesis pathway: a common feature of chlorophyll *b*-deficient mutants. - *Physiol. Plant.* **97**: 311-320, 1996.
- Freeman, T.P., Duysen, M.E., Olson, N.H., Williams, N.D.: Electron transport and chloroplast ultrastructure of a chlorophyll-deficient mutant of wheat. - *Photosyn. Res.* **3**: 179-189, 1982.
- Freeman, T.P., Duysen, M.E., Williams, N.D.: Effects of gene dosage on light harvesting chlorophyll accumulation, chloroplast development, and photosynthesis in wheat. - *Can. J. Bot.* **65**: 2118-2123, 1987.
- Guo, A.G., Feng, X.Z., Zhao, L.L., Wang, P.H.: [Studies on the activity of translation and transcription in the leaves of wheat albescent line.] - *Acta agr. boreali-occidentalis sin.*

- 24: 1-4, 1996. [In Chin., ab: E]
- Hsieh, M.H., Goodman, H.M.: The *Arabidopsis* IspH homolog is involved in the plastid non-mevalonate pathway of isoprenoid biosynthesis. - *Plant Physiol.* **138**: 641-653, 2005.
- Huang, X., Wang, P., Zhao, H., Deng, X.: Genetic analysis and molecular mapping of a novel chlorophyll-deficient mutant gene in rice. - *Rice Sci.* **15**: 7-12, 2008.
- Jiang, L., Guo, L., Jiang, H., Zeng, D., Hu, J., Wu, L., Liu, J., Gao, Z., Qian, Q.: Genetic analysis and fine-mapping of a *dwarfing with withered leaf-tip* mutant in rice. - *J. Genet. Genomics* **35**: 715-721, 2008.
- Johansen, D.A.: *Plant Micro-Technique*. - McGraw-Hill Book Company, New York 1940.
- Kaul, M.L.H., Bhan, A.K.: Mutagenic effectiveness and efficiency of EMS, DES and gamma rays in rice. - *Theor. appl. Genet.* **50**: 241-246, 1977.
- Kim, D.Y., Bovet, L., Kushnir, S., Noh, E.W., Martinoia, E., Lee, Y.: AtATM3 is involved in heavy metal resistance in *Arabidopsis*. - *Plant Physiol.* **140**: 922-932, 2006.
- Koski, V.: Chlorophyll formation in seedlings of *Zea mays* L. - *Arch. Biochem. Biophys.* **29**: 339-343, 1950.
- Lander, E.S., Green, P., Abrahamson, J., Barlow, A., Daly, M.J., Lincoln, S.E., Newburg, L.: *MAPMAKER*: an interactive computer package for constructing primary genetic linkage maps of experimental and natural populations. - *Genomics* **1**: 174-181, 1987.
- Li, C., Lin, D.-Z., Dong, Y.-J., Ye, S.-H., Zhang, X.-M.: Genetic analysis and mapping of a thermo-sensitive white stripe-leaf mutant at seedling stage in rice (*Oryza sativa*). - *Rice Sci.* **17**: 276-281, 2010.
- Lincoln, S.E., Daly, M.J., Lander, E.S.: Constructing genetic maps with *MAPMAKER/EXP version 3.0*: a tutorial and reference manual. - Whitehead Institute for Biomedical Research, Cambridge 1993.
- Liu, W.Z., Fu, Y.P., Hu, G.C., Si, H.M., Zhu, L., Wu, C., Sun, Z.X.: Identification and fine mapping of a thermo-sensitive chlorophyll deficient mutant in rice (*Oryza sativa* L.). - *Planta* **226**: 785-795, 2007.
- Liu, Z.L., Yang, Z., Liu, W.J., Du, J.B., Tian, W.J., Luo, M.H., Lin, H.H.: Mutation mechanism of chlorophyll-less barley mutant NYB. - *Photosynthetica* **46**: 73-78, 2008.
- Long, D., Martin, M., Sundberg, E., Swinburne, J., Puangsomlee, Coupland, G.: The maize transposable element system *Ac/Ds* as a mutagen in *Arabidopsis*: Identification of an albino mutation induced by Ds insertion. - *Proc. nat. Acad. Sci. USA* **90**: 10370-10374, 1993.
- Mochizuki, N., Brusslan, J.A., Larkin, R., Nagatani, A., Chory, J.: *Arabidopsis* genome's uncoupled 5 (GUN5) mutant reveals the involvement of Mg-chelatase H subunit in plastid-to-nucleus signal transduction. - *Proc. nat. Acad. Sci. USA* **98**: 2053-2058, 2001.
- Mou, Z.L., He, Y.K., Dai, Y., Liu, X.F., Li, J.: Deficiency in fatty acid synthase leads to premature cell death and dramatic alteration in plant morphology. - *Plant Cell* **12**: 405-417, 2000.
- Nealson, K.H., Conrad, P.G.: Life: past, present and future. - *Phil Trans. roy. Soc. London* **B 354**: 1923-1939, 1999.
- Parks, B.M., Quail, P.H.: Phytochrome-deficient *hy1* and *hy2* long hypocotyls mutant of *Arabidopsis* are defective in phytochrome chromophore biosynthesis. - *Plant Cell* **3**: 1177-1186, 1991.
- Pettigrew, R., Driscoll, C.J., Rienits, K.G.: A spontaneous chlorophyll mutant in hexaploid wheat. - *Heredity* **24**: 481-487, 1969.
- Saghai-Maroo, M.A., Soliman, K.M., Jorgensen, A.R., Allard, R.W.: Ribosomal DNA spacer length polymorphism in barley: Mendelian inheritance, chromosomal location and population dynamics. - *Proc. nat. Acad. Sci.* **81**: 8014-8018, 1984.
- Sang, X., Fang, L., Vanichpakorn, Y., Ling, Y., Du, P., Zhao, F., Yang, Z., He, G.: Physiological character and molecular mapping of leaf-color mutant *wyv1* in rice (*Oryza sativa* L.). - *Genes Genomics* **32**: 123-128, 2010.
- Sears, L.M.S., Sears, E.R.: The mutants chlorina-1 and Hermesen's virescent. - In: Finlay, K.W., Shepherd, K.W. (ed.): *Proceedings of the Third International Wheat Genetics Symposium*. Pp. 299-305. Plenum Press, New York 1968.
- Shen, B., Zhuang, J., Zhang, K., Dai, W., Lu, Y., Fu, L., Ding, J., Zheng, K.: QTL mapping of chlorophyll contents in rice. - *Agr. Sci. China* **6**: 17-24, 2007.
- Singh, K., Ghai, M., Garg, M., Chhuneja, P., Kaur, P., Schnurbusch, T., Keller, B., Dhaliwal, H.S.: An integrated molecular linkage map of diploid wheat based on a *Triticum boeoticum* × *T. monococcum* RIL population. - *Theor. appl. Genet.* **115**: 301-312, 2007.
- Somers, D.J., Isaac, P., Edwards, K.: A high-density microsatellite consensus map for bread wheat (*Triticum aestivum* L.). - *Theor. appl. Genet.* **109**: 1105-1114, 2004.
- Stern, D.B., Hanson, M.R., Barkan, A.: Genetics and genomics of chloroplast biogenesis: maize as a model system. - *Trends Plant Sci.* **9**: 293-301, 2004.
- Tian, F.X., Gong, J.F., Wang, G.P., Wang, G.K., Fan, Z.Y., Wang, W.: Improved drought resistance in a wheat stay-green mutant *tasg1* under field conditions. - *Biol. Plant.* **56**: 509-515, 2012.
- Wang, Q.S., Sang, X.C., Ling, Y.H., Zhao, F.M., Yang, Z.L., Li, Y., He, G.H.: Genetic analysis and molecular mapping of a novel gene for zebra mutation in rice (*Oryza sativa* L.). - *J. Genet. Genomics* **36**: 679-684, 2009.
- Washington, W.J., Sears, E.R.: Ethyl methane sulfonate-induced chlorophyll mutations in *Triticum aestivum*. - *Can. J. Genet. Cytol.* **12**: 851-859, 1970.
- Wicker, T., Stein, N., Albar, L., Feuillet, C., Schlagenhauf, E., Keller, B.: Analysis of a contiguous 211 kb sequence in diploid wheat (*Triticum monococcum* L.) reveals multiple mechanisms of genome evolution. - *Plant J.* **26**: 307-316, 2001.
- Wu, Z.M., Zhang, X., He, B., Diao, L.P., Sheng, S.L., Wang, J.L., Guo, X.P., Su, N., Wang, L.F., Jiang, L., Wang, C.M., Zhai, H.Q., Wan, J.M.: A chlorophyll-deficient rice mutant with impaired chlorophyllide esterification in chlorophyll biosynthesis. - *Plant Physiol.* **145**: 29-40, 2007.
- Zhang, H., Zhang, D., Han, S., Zhang, X., Yu, D.: Identification and gene mapping of a soybean chlorophyll-deficient mutant. - *Plant Breed.* **130**: 133-138, 2011.
- Zou, J.J., Singh, R.G., Hymowitz, T.: Association of the yellow leaf (y10) mutant to soybean chromosome 3. - *J. Hered.* **94**: 352-355, 2003.