

Gorlin Syndrome in a type IV-skin person with a novel PTCH1 mutation: Case report and literature review

S. AlSalem^a, Y. Binamer^{b,c,*}^a King Khalid University Hospital, College of Medicine, King Saud University, Riyadh, Saudi Arabia^b King Faisal Specialist Hospital and Research Centre, Riyadh, Saudi Arabia^c Alfaisal University, Riyadh, Saudi Arabia

Received 10 May 2015; accepted 23 August 2015

Abstract

Nevoid basal cell carcinoma syndrome (NBCCS), or Gorlin Syndrome, is an autosomal dominant multisystem disorder, characterized by multiple developmental abnormalities and inactivation germline mutations in the human homolog of the patched (PTCH) gene. We are presenting a case of NBCCS in a skin type 4 Saudi male with a novel PTCH1 gene mutation. To the best of our knowledge, this is the third case reported in Saudi Arabia but the first in adult population. Moreover, our patient harbors a novel heterozygosity mutation in patch1 gene.

© 2015 The Authors. Production and hosting by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Keywords: Basal cell carcinoma; Gorlin Syndrome

1. Introduction

Nevoid basal cell carcinoma syndrome (NBCCS), basal cell nevus syndrome, or Gorlin Syndrome [OMIM, 109400] refer to the same rare genetic mulitsystemic disorder. Inheritance is autosomal dominant with near-complete penetrance and variable expression. It is characterized by multiple developmental abnormalities and inactivation germline mutations in the human homolog of the patched (PTCH) gene. NBCCS can involve skin, bone and the nervous system. Major features include multiple basal cell car-

cinomas (BCC); odontogenic keratocysts; palm and/or sole pits and calcification of the falx cerebri. Other features include as well cleft lip or palate, microphthalmos, a bifid or fused rib, scoliosis, tall stature, frontal bossing and medulloblastoma.

It is a rare condition in the dark skin population. To the best of our knowledge, herein, we report the third case of NBCCS in Saudi population and the first one in the adulthood with a novel PTCH1 gene mutation.

2. Case report

A 38 year-old male soldier from AlJouf, North of Saudi Arabia, presented with a history of multiple biopsy-proven pigmented BCCs overlying the abdomen, scalp, and right shoulder one year ago. Since then, he developed more BCCs over the same areas. He spent significant time outside without sun protection. There was no trauma, ionizing radiation, or exposure to chemicals. He is a heavy smoker. Family history is positive for non-consanguinity and a

* Corresponding author at: Alfaisal University, P.O. Box 3354, MBC 104, Riyadh 11211, Saudi Arabia.

E-mail address: Dr.binamer@gmail.com (Y. Binamer).

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier

deceased mother with numerous skin tumors that remained undiagnosed.

Upon skin examination, multiple, >20 pigmented (black) papulo-nodular lesions were found on the scalp with variability in size, Fig. 1. Skin tags were present. There was frontal bossing, with confluent eyebrows (Fig. inset b) and tall stature (184 cm). There were numerous bilateral plantar (Fig. inset c) pits.

NBCCS was suspected based on the clinical presentation. X-rays of the face revealed well-defined lytic bony lesions involving the mandible. CT of the head suggested odontogenic bony keratocysts of the mandible and maxillary bones as well as extensive calcification of the falx cerebri. There was no evidence of cardiac fibromas on echocardiogram. Confirmation with a genetic study showed positivity for a haploinsufficiency with novel partial deletion (EXON 13) of the PTCH1 gene. Therefore, nevoid basal cell carcinoma syndrome was confirmed. BCCs were treated with surgical excision and curettage.

3. Discussion

NBCCS is a rare inherited multisystem condition caused by germline mutations in the PTCH gene (Cohen, 1999). Jarisch and White first described it in 1894, which was later followed by greater establishment of the syndrome by Gorlin and Goltz (1960). It has the estimated prevalence of 1:57000 to 1:164000 individuals. It affects both genders equally, with greater incidence in Caucasians and less in blacks and Asians (Goldstein et al., 1994; Ahn et al., 2004). Extensive literature review identified two pediatric cases in the Saudi population while our patient is the first case diagnosed in adulthood.

Interestingly, more than one hundred clinical findings have been reported in the English literature. The diagnosis is clinical, and is supported by the presence of two major, or one major and two minor criteria (Table 1) Evans et al. (1993). Major manifestations are found in 75–80% of cases, and include early development of basal cell carcinomas;

odontogenic keratocysts; palm and/or plantar pitting; ectopic calcification of the falx cerebri; and family history of NBCCS (Evans et al., 1993). Minor criteria include bifid ribs; craniofacial anomalies such as coarse facial features, macrocephaly, frontal bossing, confluent brows, and hypertelorism; early onset midline medulloblastoma; ovarian or cardiac fibromas; lymphomesenteric cysts; and congenital malformations like cleft lip or palate, polydactyly, eye abnormalities, cataracts, glaucoma, and coloboma (Evans et al., 1993). Diagnosis can be difficult in young patients, and atypical cases in persons with dark complexion.

Skin manifestations include BCCs, palmoplantar pits, milia, hemangiomas, and skin tags (Walling et al., 2004). BCC occurs in approximately 75% of NBCCS Caucasian patients and are multiple (Kimonis et al., 1997). Interestingly, one-third of all sporadic basal cell carcinomas have mutations in the PTCH gene (Gailani et al., 1996). In NBCCS, the age at first diagnosis ranges from three to 53 years with the mean average of 20–21 (Kimonis et al., 1997). The histopathological examination of gorlins' BCCs is indistinct from sporadic cases (Gorlin, 2004). In this syndrome, BCCs are largely evident in Caucasians (80%) than in black (30–40%) or Asian individuals (Goldstein et al., 1994; Ahn et al., 2004; Kulkarni et al., 2003). This finding is principally expected because darker skinned populations are intrinsically resistant to UV light.

Palmoplantar pits are a common feature. They are secondary to partial or complete absence of the horny cell layer of the skin (Howell and Mehregan, 1970). Odontogenic keratocysts occur mainly in the mandible, and unlike BCCs, share no racial predilection (Goldstein et al., 1994). Ectopic calcification occurs in the falx cerebri in 65% of patients with rates increasing to 90% in those over 40 years old (Kimonis et al., 1997). Our patient had all major and some minor criteria for diagnosis in addition to tall stature, frontal bossing, and coarse facial features were present.

NBCCS must be distinguished from other rare genodermatosis with malignant potential e.g., Bazex, Rombo, Muir-torre syndrome, Xeroderma pigmentosum and Basaloid follicular hamartomas (BFH). Bazex syndrome is a X-linked condition characterized by multiple BCCs, hypotrichosis, hypohidrosis and acral follicular atrophoderma (Hauck and Manders, 1994). Rombo syndrome is characterized by multiple BCCs, vermiculate atrophoderma, milia, hypotrichosis, trichoepitheliomas and peripheral cyanosis (Hauck and Manders, 1994). Muir-Torre syndrome is an autosomal dominant condition with various skin sebaceous neoplasms and visceral malignancies (Hauck and Manders, 1994).

NBCCS results from mutations in the PTCH gene, the human homolog of *Drosophila*. It exhibits autosomal dominance with near-complete penetrance but is variably expressed (Anderson et al., 1967). The patched gene is located in the long arm of chromosome 9 (Farndon et al., 1992). The transcribed PTCH protein is part of a receptor complex (with Smoothed, SMO) for the hedgehog protein, part of the sonic hedgehog-signaling pathway (Stone



Fig. 1. Multiple BCCs over the scalp. Inset b: frontal bossing. Inset c: plantar pits.

Table 1
Major and minor clinical features (Evans et al., 1993).

Major features	Minor features
<ul style="list-style-type: none"> • Multiple basal cell carcinomas or Basal cell carcinoma before age 20 • Odontogenic keratocysts confirmed by histology • Palmar or plantar pits (>3) • Ectopic bilamellar calcification of the falx cerebri • Positive family history 	<ul style="list-style-type: none"> • Craniofacial anomalies: macrocephaly, frontal bossing, hypertelorism • Bifid, or splayed ribs • Early onset medulloblastoma • Fibroma: ovarian, or cardiac • Lymphomesenteric cysts • Congenital malformations: cleft lip/palate, eye abnormalities, hand and finger abnormalities

Table 2
Comparison of characteristics in NBCCS reports in Saudi Arabia (Alghamdi, 2008; Hellani et al., 2009).

Characteristic	Our patient	Hellani et al.	Alghamdi
Age and gender	38 years, male	15 years, male	10, 8, 6 years old, female siblings
BCC	+, >20	–	+
Ectopic Calcification	+	+	+
Jaw keratocyst	+	+	–
Palmar/plantar pits	+	+	+
Family history	+	–	–
Minor findings	Frontal bossing, skin tags, coarse facies, tall stature, confluent eyebrows	Frontal bossing, confluent eyebrows, pectus excavatum, bifid ribs, spina bifida	Frontal bossing, skin tags, coarse facies, hydrocephalus
Gene	+PTCH1, exon 13 deletion	+PTCH1, de novo mutation (c.1291delC)	Not reported

et al., 1996; Grachtchouk et al., 2003). This is supported by the fact that PTCH1 mRNA is abundant in odontogenic keratocysts and BCCs from Gorlin Syndrome patients but not in other skin neoplasms (Nagano et al., 1999). Gorlin Syndrome patients are abnormally sensitive to therapeutic doses of ionizing radiation, with development of numerous BCCs in the irradiated site shortly after exposure (Evans et al., 1993). Our patient had a novel PTCH1 gene mutation result by array-based comparative genomic hybridization (aCGH) in the form of deletion of exon 13.

NBCCS patients should be managed by a multidisciplinary team (Lo Muzio et al., 1999). BCC should be treated with surgery if possible. However, other destructive modalities or topical immunomodulators can be used. Vismodegib is a specific oral SHH pathway component inhibitor that can be used in difficult to treat or metastatic cases (Tang et al., 2012). Sun protection and close surveillance are essentials (Frentz et al., 1987).

In Saudi Arabia, two reports of pediatric NBCCS were identified. Alghamdi reported a family with three sisters with Gorlin Syndrome (Alghamdi, 2008). Another report by Hellani et al. identified a 15-year-old male with NBCCS (Hellani et al., 2009) (Table 2). To the best of our knowledge, our patient is considered the first reported case of Adult NBCCS in the kingdom of Saudi Arabia and the establishment of a novel genetic mutation of the patched1 gene.

Funding

None.

Conflicts of interest

None.

References

- Ahn, S.G., Lim, Y.S., Kim, D.K., et al., 2004. Nevoid basal cell carcinoma syndrome: a retrospective analysis of 33 affected Korean individuals. *Int. J. Oral. Maxillofac. Surg.* 33, 458.
- Alghamdi, Y., 2008. Skin tags as a presenting sign of basal cell nevus syndrome in three sisters of the same family. *Ann. Saudi Med.* 28 (2), 132–134.
- Anderson, D.E., Taylor, W.B., Falls, H.F., Davidson, R.T., 1967. The nevoid basal cell carcinoma syndrome. *Am. J. Hum. Genet.* 19, 12.
- Cohen Jr., M.M., 1999. Nevoid basal cell carcinoma syndrome: molecular biology and new hypotheses. *Int. J. Oral Maxillofac. Surg.* 28, 216.
- Evans, D.G., Ladusans, E.J., Rimmer, S., et al., 1993. Complications of the naevoid basal cell carcinoma syndrome: results of a population based study. *J. Med. Genet.* 30, 460.
- Fardon, P.A., Del Mastro, R.G., Evans, D.G., Kilpatrick, M.W., 1992. Location of gene for Gorlin syndrome. *Lancet* 339, 581.
- Frentz, G., Munch-Petersen, B., Wulf, H.C., et al., 1987. The nevoid basal cell carcinoma syndrome: sensitivity to ultraviolet and X-ray irradiation. *J. Am. Acad. Dermatol.* 17, 637.
- Gailani, M.R., Stähle-Bäckdahl, M., Leffell, D.J., et al., 1996. The role of the human homologue of *Drosophila* patched in sporadic basal cell carcinomas. *Nat. Genet.* 14, 78.
- Goldstein, A.M., Pastakia, B., DiGiovanna, J.J., et al., 1994. Clinical findings in two African-American families with the nevoid basal cell carcinoma syndrome (NBCC). *Am. J. Med. Genet.* 50, 272.
- Gorlin, R.J., 2004. Nevoid basal cell carcinoma (Gorlin) syndrome. *Genet. Med.* 6, 530–539.
- Gorlin, R.J., Goltz, R.W., 1960. Multiple nevoid basal-cell epithelioma, jaw cysts and bifid rib. A syndrome. *N. Engl. J. Med.* 262, 908–912.

- Grachtchouk, V., Grachtchouk, M., Lowe, L., et al., 2003. The magnitude of hedgehog signaling activity defines skin tumor phenotype. *EMBO J.* 22, 2741.
- Hauck, R.M., Manders, E.K., 1994. Familial syndromes with skin tumor markers. *Ann. Plast. Surg.* 33, 102–111.
- Hellani, A., Baghdadi, H., Dabbour, N., Almassri, N., Abu-Amero, K.K., 2009. A novel PTCH1 germline mutation distinguishes basal cell carcinoma from basaloid follicular hamartoma: a case report. *J. Med. Case Rep.* 3, 52.
- Howell, J.B., Mehregan, A.H., 1970. Pursuit of the pits in the nevoid basal cell carcinoma syndrome. *Arch. Dermatol.* 102, 586.
- Kimonis, V.E., Goldstein, A.M., Pastakia, B., et al., 1997. Clinical manifestations in 105 persons with nevoid basal cell carcinoma syndrome. *Am. J. Med. Genet.* 69, 299.
- Kulkarni, P., Brashear, R., Chuang, T.Y., 2003. Nevoid basal cell carcinoma syndrome in a person with dark skin. *J. Am. Acad. Dermatol.* 49, 332.
- Lo Muzio, L., Nocini, P., Bucci, P., Pannone, G., Consolo, U., Procaccini, M., 1999. Early diagnosis of nevoid basal cell carcinoma syndrome. *J. Am. Dent. Assoc.* 130, 669–674.
- Nagano, T., Bito, T., Kallassy, M., et al., 1999. Overexpression of the human homologue of *Drosophila* patched (PTCH) in skin tumours: specificity for basal cell carcinoma. *Br. J. Dermatol.* 140, 287.
- Stone, D.M., Hynes, M., Armanini, M., Swanson, T.A., Gu, Q., Johnson, R.L., Scott, M.P., Pennica, D., Goddard, A., Phillips, H., Noll, M., Hooper, J.E., de Sauvage, F., Rosenthal, A., 1996. The tumour-suppressor gene patched encodes a candidate receptor for Sonic hedgehog. *Nature* 384 (6605), 129–134.
- Tang, J.Y., Mackay-Wiggan, J.M., Aszterbaum, M., et al., 2012. Inhibiting the hedgehog pathway in patients with the basal-cell nevus syndrome. *N. Engl. J. Med.* 366, 2180.
- Walling, H.W., Fosko, S.W., Geraminejad, P.A., Whitaker, D.C., Arpey, C.J., 2004. Aggressive basal cell carcinoma: presentation, pathogenesis, and management. *Cancer Metastasis Rev.* 23 (3–4), 389–402.