

Applications of Differential Geometry:  
Cartography (Map-making)  
Math 473  
Introduction to Differential Geometry  
Lecture 21

Dr. Nasser Bin Turki

King Saud University  
Department of Mathematics

December 5, 2017

# Applications of Differential Geometry: Cartography (Map-making)

We can think of a geographic map of the Earth as a mapping from a domain in the plane to a domain on the surface of the Earth, i.e. as a surface patch that parametrises a part of a sphere.

# Applications of Differential Geometry: Cartography (Map-making)

We can think of a geographic map of the Earth as a mapping from a domain in the plane to a domain on the surface of the Earth, i.e. as a surface patch that parametrises a part of a sphere.

For the geographic map to be useful, we would like this mapping to be bijective, continuous, smooth. We would also like the mapping to preserve some important metric information, for example to preserve distances, lengths, angles, areas.

### Denition (1):

We say that a surface patch  $X : U \rightarrow \mathbb{R}^3$  is a **true map** if it preserves the lengths of all curves,

### Denition (1):

We say that a surface patch  $X : U \rightarrow \mathbb{R}^3$  is a **true map** if it preserves the lengths of all curves, i.e. if the length of a curve  $[a, b] \rightarrow U$ ,  $t \mapsto (u(t), v(t))$  in a map is the same as the length of the corresponding curve  $[a, b] \rightarrow \mathbb{R}^3$ ,  $t \mapsto X(u(t), v(t))$  on the surface.

### Denition (1):

We say that a surface patch  $X : U \rightarrow \mathbb{R}^3$  is a **true map** if it preserves the lengths of all curves, i.e. if the length of a curve  $[a, b] \rightarrow U$ ,  $t \mapsto (u(t), v(t))$  in a map is the same as the length of the corresponding curve  $[a, b] \rightarrow \mathbb{R}^3$ ,  $t \mapsto X(u(t), v(t))$  on the surface.

### Proposition (1):

A surface patch  $X : U \rightarrow \mathbb{R}^3$  is a true map, i.e. it preserves the lengths of all curves if and only if it has the first fundamental form

$$I(u, v) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \text{ for all } (u, v) \in U.$$

**Remark:** Examples of surface patches with the first fundamental form  $I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$  are the cylinder and the cone.

**Remark:** Examples of surface patches with the first fundamental form  $I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$  are the cylinder and the cone. We know that we can make (a part of) the cylinder and the cone from a flat piece of paper without stretching or contracting, hence we have true maps of the surface of the cylinder and the cone respectively.

**Proposition (2):**

There is no true map of the Earth, i.e. there is no mapping from (a part of) the plane to (an open part of) a sphere which preserves the lengths of all curves.



**Remark:** A surface patch  $X : U \rightarrow \mathbb{R}^3$  changes all lengths of curves by the same constant factor  $\lambda > 0$  if and only if it has the first fundamental form  $I(u, v) = \begin{pmatrix} \lambda & 0 \\ 0 & \lambda \end{pmatrix} = \lambda \cdot \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$  for all  $(u, v) \in U$ .

**Remark:** Even though there is no true map of the Earth, there are many maps with various useful properties, for example preserving angles or areas.

**Definition (2):** We say that a surface patch  $X : U \rightarrow \mathbb{R}^3$  is **conformal** if it preserves the angles between all curves,

**Definition (2):** We say that a surface patch  $X : U \rightarrow \mathbb{R}^3$  is **conformal** if it preserves the angles between all curves, i.e. if the angle between two curves in  $U$  is the same as the angle between the corresponding curves on the surface.

**Remark:** A surface patch  $X : U \rightarrow \mathbb{R}^3$  is conformal if and only if the coefficients of the first fundamental form satisfy the equations  $F(u, v) = 0$  and  $E(u, v) = G(u, v)$  for all  $(u, v) \in U$ . For example the stereographic projection is a conformal surface patch.

*Thanks for listening.*