

Second Fundamental Form
Math 473
Introduction to Differential Geometry
Lecture 24

Dr. Nasser Bin Turki

King Saud University
Department of Mathematics

December 10, 2017

Example (Orthogonal Sections):

Let Π be a plane containing the surface normal N . Let γ be the curve that we obtain in the intersection of the surface with the plane Π .

Example (Orthogonal Sections):

Let Π be a plane containing the surface normal N . Let γ be the curve that we obtain in the intersection of the surface with the plane Π . Then the curve γ is contained in the plane Π , hence the plane Π is the osculating plane of γ . The plane Π is spanned by the unit tangent T and the surface normal N , while the tangent plane of the surface is orthogonal to the surface normal N , hence the tangent plane of the surface and the osculating plane of the curve γ are orthogonal, i.e. the angle $\varphi = \angle(B, N) = \pm\pi/2$.

Example (Orthogonal Sections):

Let Π be a plane containing the surface normal N . Let γ be the curve that we obtain in the intersection of the surface with the plane Π . Then the curve γ is contained in the plane Π , hence the plane Π is the osculating plane of γ . The plane Π is spanned by the unit tangent T and the surface normal N , while the tangent plane of the surface is orthogonal to the surface normal N , hence the tangent plane of the surface and the osculating plane of the curve γ are orthogonal, i.e. the angle $\varphi = \angle(B, N) = \pm\pi/2$. Thus the orthogonal section γ is a geodesic, $\kappa_g = 0$, and the normal curvature is $\kappa_n = \pm\kappa$ (where κ is the curvature of the curve γ as a plane curve).

Let $X : U \rightarrow \mathbb{R}^3$ be a regular injective surface patch.

Let $N : U \rightarrow \mathbb{R}^3$ be a unit normal on the surface X . Let

$\gamma(t) = X(u(t), v(t))$ be a curve on the surface X .

Let $X : U \rightarrow \mathbb{R}^3$ be a regular injective surface patch.
Let $N : U \rightarrow \mathbb{R}^3$ be a unit normal on the surface X . Let
 $\gamma(t) = X(u(t), v(t))$ be a curve on the surface X .

Recall:

Remember how we computed $\gamma' = u' \cdot X_u + v' \cdot X_v$ and

$$\begin{aligned} |\gamma'|^2 &= \gamma' \bullet \gamma' = (u' \cdot X_u + v' \cdot X_v) \bullet (u' \cdot X_u + v' \cdot X_v) \\ &= (u')^2 \cdot (X_u \bullet X_u) + 2u'v' \cdot (X_u \bullet X_v) + (v')^2 \cdot (X_v \bullet X_v) \end{aligned}$$

and therefore defined the coefficients of the first fundamental form
as

$$E = X_u \bullet X_u, \quad F = X_u \bullet X_v, \quad G = X_v \bullet X_v.$$

The first fundamental form tell us information about the distances, angles and areas on the surface. We are going to define the second fundamental form which tell us information about the curvature of the surface.

The first fundamental form tell us information about the distances, angles and areas on the surface. We are going to define the second fundamental form which tell us information about the curvature of the surface.

Notation: $X_u = \frac{\partial X}{\partial u}$, $X_v = \frac{\partial X}{\partial v}$, $X_{uu} = \frac{\partial}{\partial u} \frac{\partial}{\partial u} X$, $X_{uv} = \frac{\partial}{\partial u} \frac{\partial}{\partial v} X$,
 $X_{vu} = \frac{\partial}{\partial v} \frac{\partial}{\partial u} X$, $X_{vv} = \frac{\partial}{\partial v} \frac{\partial}{\partial v} X$.

Note that $X_{uv} = \frac{\partial}{\partial u} \frac{\partial}{\partial v} X = \frac{\partial}{\partial v} \frac{\partial}{\partial u} X = X_{vu}$.

How to compute the normal curvature?

How to compute the normal curvature?

We know that

$$\kappa_n = \frac{\tilde{T}' \bullet \tilde{N}}{|\gamma'| \cdot |\tilde{T}| \cdot |\tilde{N}|}.$$

How to compute the normal curvature?

We know that

$$\kappa_n = \frac{\tilde{T}' \bullet \tilde{N}}{|\gamma'| \cdot |\tilde{T}| \cdot |\tilde{N}|}.$$

We take $\tilde{T} = \gamma'$ and $\tilde{N} = N$. Using the chain rule we compute $\tilde{T}' = \gamma'' = u'X_u + v'X_v$. Using the chain rule again we compute

How to compute the normal curvature?

We know that

$$\kappa_n = \frac{\tilde{T}' \bullet \tilde{N}}{|\gamma'| \cdot |\tilde{T}| \cdot |\tilde{N}|}.$$

We take $\tilde{T} = \gamma'$ and $\tilde{N} = N$. Using the chain rule we compute $\tilde{T} = \gamma' = u'X_u + v'X_v$. Using the chain rule again we compute

$$\begin{aligned}\tilde{T}' &= \gamma'' = (u'X_u + v'X_v)' \\ &= u''X_u + u'(u'(X_u)_u + v'(X_u)_v) + v''X_v + v'(u'(X_v)_u + v'(X_v)_v) \\ &= u''X_u + v''X_v + u'u'X_{uu} + u'v'X_{uv} + v'u'X_{vu} + v'v'X_{vv} \\ &= u''X_u + v''X_v + (u')^2X_{uu} + 2u'v'X_{uv} + (v')^2X_{vv}\end{aligned}$$

How to compute the normal curvature?

We know that

$$\kappa_n = \frac{\tilde{T}' \bullet \tilde{N}}{|\gamma'| \cdot |\tilde{T}| \cdot |\tilde{N}|}.$$

We take $\tilde{T} = \gamma'$ and $\tilde{N} = N$. Using the chain rule we compute $\tilde{T} = \gamma' = u'X_u + v'X_v$. Using the chain rule again we compute

$$\begin{aligned}\tilde{T}' = \gamma'' &= (u'X_u + v'X_v)' \\ &= u''X_u + u'(u'(X_u)_u + v'(X_u)_v) + v''X_v + v'(u'(X_v)_u + v'(X_v)_v) \\ &= u''X_u + v''X_v + u'u'X_{uu} + u'v'X_{uv} + v'u'X_{vu} + v'v'X_{vv} \\ &= u''X_u + v''X_v + (u')^2X_{uu} + 2u'v'X_{uv} + (v')^2X_{vv}\end{aligned}$$

and

$$\begin{aligned}\tilde{T}' \bullet \tilde{N} &= (u''X_u + v''X_v + (u')^2X_{uu} + 2u'v'X_{uv} + (v')^2X_{vv}) \bullet N \\ &= u'' \cdot (X_u \bullet N) + v'' \cdot (X_v \bullet N) \\ &\quad + (u')^2 \cdot (X_{uu} \bullet N) + 2u'v' \cdot (X_{uv} \bullet N) + (v')^2 \cdot (X_{vv} \bullet N).\end{aligned}$$

Using the fact that the normal N is perpendicular to the tangent plane (or using the fact that N is a multiple of $X_u \times X_v$), we see that $X_u \bullet N = X_v \bullet N = 0$, hence

Using the fact that the normal N is perpendicular to the tangent plane (or using the fact that N is a multiple of $X_u \times X_v$), we see that $X_u \bullet N = X_v \bullet N = 0$, hence

$$\tilde{T}' \bullet \tilde{N} = (u')^2 \cdot (X_{uu} \bullet N) + 2u'v' \cdot (X_{uv} \bullet N) + (v')^2 \cdot (X_{vv} \bullet N).$$

Definition (1):

The **coefficients of the second fundamental form** of the surface patch $X : U \rightarrow \mathbb{R}^3$ are

$$e = X_{uu} \bullet N, \quad f = X_{uv} \bullet N = X_{vu} \bullet N, \quad g = X_{vv} \bullet N.$$


The **second fundamental form** of X is

$$\text{II} = \begin{pmatrix} e & f \\ f & g \end{pmatrix}.$$

Examples

Example (1):


Let $X : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be given by $X(u, v) = (u, v, 5)$. Compute the second fundamental form of the surface X .



Examples

Example (2):

Let $X : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be given by $X(u, v) = (a \cos u, a \sin u, bv)$, where a, b are constants. Compute the coefficients of the second fundamental form of the surface X .



Proposition (1):

Let $\gamma(t) = X(u(t), v(t))$ be a curve on the surface X . Then the normal curvature of γ is

$$\kappa_n = \frac{(u')^2 \cdot e + 2u'v' \cdot f + (v')^2 \cdot g}{(u')^2 \cdot E + 2u'v' \cdot F + (v')^2 \cdot G} = \frac{(u' \ v') \cdot \begin{pmatrix} e & f \\ f & g \end{pmatrix} \cdot \begin{pmatrix} u' \\ v' \end{pmatrix}}{(u' \ v') \cdot \begin{pmatrix} E & F \\ F & G \end{pmatrix} \cdot \begin{pmatrix} u' \\ v' \end{pmatrix}}.$$

Proposition (1):

Let $\gamma(t) = X(u(t), v(t))$ be a curve on the surface X . Then the normal curvature of γ is

$$\kappa_n = \frac{(u')^2 \cdot e + 2u'v' \cdot f + (v')^2 \cdot g}{(u')^2 \cdot E + 2u'v' \cdot F + (v')^2 \cdot G} = \frac{(u' \ v') \cdot \begin{pmatrix} e & f \\ f & g \end{pmatrix} \cdot \begin{pmatrix} u' \\ v' \end{pmatrix}}{(u' \ v') \cdot \begin{pmatrix} E & F \\ F & G \end{pmatrix} \cdot \begin{pmatrix} u' \\ v' \end{pmatrix}}.$$

Proof:

Proposition (2):

Let $X : U \rightarrow \mathbb{R}^3$ be a regular surface. If the second fundamental form of the surface X vanishes i.e. $II = 0$, then the surface X is part of a plane.

Proposition (2):

Let $X : U \rightarrow \mathbb{R}^3$ be a regular surface. If the second fundamental form of the surface X vanishes i.e. $II = 0$, then the surface X is part of a plane.

Proof:

Thanks for listening.