

Serret-Frenet Basis for Space Curves
Math 473
Introduction to Differential Geometry
Lecture 5

Dr. Nasser Bin Turki

King Saud University
Department of Mathematics

October 8, 2017

Principal Normal and Binormal

Let $\alpha : I \mapsto \mathbb{R}^3$ be a regular space curve. Let $t \in I$ be such that $\kappa(t) \neq 0$.

Principal Normal and Binormal

Let $\alpha : I \mapsto \mathbb{R}^3$ be a regular space curve. Let $t \in I$ be such that $\kappa(t) \neq 0$.

Defnition (1):

We know that the vector $T'(t)$ is either orthogonal to $T(t)$ or $T'(t) = 0$. But $T'(t) \neq 0$ since $|T'(t)| = \kappa(t) \cdot |\alpha'(t)| \neq 0$. We define the **principal normal** of α as

Principal Normal and Binormal

Let $\alpha : I \mapsto \mathbb{R}^3$ be a regular space curve. Let $t \in I$ be such that $\kappa(t) \neq 0$.

Defnition (1):

We know that the vector $T'(t)$ is either orthogonal to $T(t)$ or $T'(t) = 0$. But $T'(t) \neq 0$ since $|T'(t)| = \kappa(t) \cdot |\alpha'(t)| \neq 0$. We define the **principal normal** of α as

$$N = \frac{T'}{\kappa}.$$

Principal Normal and Binormal

Let $\alpha : I \mapsto \mathbb{R}^3$ be a regular space curve. Let $t \in I$ be such that $\kappa(t) \neq 0$.

Defnition (1):

We know that the vector $T'(t)$ is either orthogonal to $T(t)$ or $T'(t) = 0$. But $T'(t) \neq 0$ since $|T'(t)| = \kappa(t) \cdot |\alpha'(t)| \neq 0$. We define the **principal normal** of α as

$$N = \frac{T'}{\kappa}.$$

We define the **binormal** of α as

Principal Normal and Binormal

Let $\alpha : I \mapsto \mathbb{R}^3$ be a regular space curve. Let $t \in I$ be such that $\kappa(t) \neq 0$.

Defnition (1):

We know that the vector $T'(t)$ is either orthogonal to $T(t)$ or $T'(t) = 0$. But $T'(t) \neq 0$ since $|T'(t)| = \kappa(t) \cdot |\alpha'(t)| \neq 0$. We define the **principal normal** of α as

$$N = \frac{T'}{\kappa}.$$

We define the **binormal** of α as

$$B = T \times N.$$

Remark:

The principal normal N is a unit vector orthogonal to T .

Remark:

The principal normal N is a unit vector orthogonal to T . The binormal B is orthogonal to both T and N and is unit length:
 $|B| = |T \times N| = |T| \cdot |N| = 1$ since $T \perp N$.

Theorem 1:

Let $\alpha : I \mapsto \mathbb{R}^3$ be a regular space curve.

Theorem 1:

Let $\alpha : I \mapsto \mathbb{R}^3$ be a regular space curve.

For each t in I with $\kappa(t) \neq 0$, the vectors $T(t)$, $N(t)$, $B(t)$ form an orthonormal basis of \mathbb{R}^3 ,

Theorem 1:

Let $\alpha : I \mapsto \mathbb{R}^3$ be a regular space curve.

For each t in I with $\kappa(t) \neq 0$, the vectors $T(t)$, $N(t)$, $B(t)$ form an orthonormal basis of \mathbb{R}^3 ,

$$\{T(t), N(t), B(t)\}$$

Theorem 1:

Let $\alpha : I \mapsto \mathbb{R}^3$ be a regular space curve.

For each t in I with $\kappa(t) \neq 0$, the vectors $T(t)$, $N(t)$, $B(t)$ form an orthonormal basis of \mathbb{R}^3 ,

$$\{T(t), N(t), B(t)\}$$

i.e. they are of length one

$$|T(t)| = 1, \quad |N(t)| = 1, \quad |B(t)| = 1$$

Theorem 1:

Let $\alpha : I \mapsto \mathbb{R}^3$ be a regular space curve.

For each t in I with $\kappa(t) \neq 0$, the vectors $T(t)$, $N(t)$, $B(t)$ form an orthonormal basis of \mathbb{R}^3 ,

$$\{T(t), N(t), B(t)\}$$

i.e. they are of length one

$$|T(t)| = 1, \quad |N(t)| = 1, \quad |B(t)| = 1$$

and orthogonal to each other:

$$T(t) \perp N(t), \quad N(t) \perp B(t), \quad B(t) \perp T(t).$$

Theorem 1:

Let $\alpha : I \mapsto \mathbb{R}^3$ be a regular space curve.

For each t in I with $\kappa(t) \neq 0$, the vectors $T(t)$, $N(t)$, $B(t)$ form an orthonormal basis of \mathbb{R}^3 ,

$$\{T(t), N(t), B(t)\}$$

i.e. they are of length one

$$|T(t)| = 1, \quad |N(t)| = 1, \quad |B(t)| = 1$$

and orthogonal to each other:

$$T(t) \perp N(t), \quad N(t) \perp B(t), \quad B(t) \perp T(t).$$

They satisfy the equations

$$T \bullet T = N \bullet N = B \bullet B = 1,$$

$$T \bullet N = N \bullet T = N \bullet B = B \bullet N = B \bullet T = T \bullet B = 0,$$

$$T \times N = B, \quad N \times B = T, \quad B \times T = N,$$

$$N \times T = -B, \quad B \times N = -T, \quad T \times B = -N.$$

Proof of Theorem 1

Defnition (2):Serret-Frenet Frame

The **Serret-Frenet frame** or the **Serret-Frenet basis** of the curve α is the orthonormal basis

$$\{T(t), N(t), B(t)\}$$

Thanks for listening.