

Effect of Infestation of *Pyxinia firma* on the Total Haemocyte Counts (THC) and Larval Growth of the *Dermestes vulpinus* (Dermestidae: Coleoptera)

M.I. Siddiqui and M.S. Al-Khalifa

Department of Zoology, College of Science, P.O. Box 2455,
King Saud University, Riyadh, Kingdom of Saudi Arabia

Abstract: *Pyxinia firma* has not previously been recorded in the mid gut of the larvae of scavenger beetle *Dermestes vulpinus*. Infected larvae of the scavenger beetles were investigated in order to determine the effects of their infection by the eugregarine, *Pyxinia firma*. Infected larvae were found to have higher total haemocyte counts and greater weight gain than uninfected ones. Infected larvae that were starved, however, lost weight much faster than uninfected larvae. The impact of infection on the physiology of the insect is also discussed and it is observed that the variation in the pattern of the total haemocyte counts between infected and uninfected larvae is indicative of the effect of the infection on the immune system of *D. vulpinus*.

Key words: *Pyxinia firma*, *Desrmestes vulpinus*, haemocytes, total haemocyte counts, THC, larval growth

INTRODUCTION

Eugregarines are known entomophilic parasites of the invertebrates (Brook, 1974; Kudo, 1966). Except for two species which are reported to be in commensal association unless under nutritional stress (Harry, 1967; Dunkel and Boush, 1969; Al-Khalifa, 1984) relatively few of these have had species their host-parasite relationships described. Among the family Dermestidae, only *Attagenus megatoma* has had its relationship with its enzootic parasite *Pyxinia frenzeli* fully reported (Dunkel *et al.*, 1982).

This study presents an original record of the species, *P. firma* in *Dermestes vulpinus* and attempts to evaluate the association between the two species. *D. vulpinus*, in its gut *P. firma* was encountered where it remains attached to the epithelial layer. During the survey of gut contents of the larvae in Saudi Arabia an identified and apparently parasitic infection was observed and advice of Natural History Museum was sought. A colony of *Pyxinia firma* and infected *D. vulpinus* separately was maintained in the laboratory for 12 months to study the host parasite relationship and transmission is carried through cannibalistic means.

MATERIALS AND METHODS

Infected and uninfected, adults and larval stages of *D. vulpinus* were collected from the skin and carcasses of sacrificed sheep and goats on the outskirts of the

Al-Kharj district, 100 km south of Riyadh. A colony was maintained in the laboratory at $28\pm1^{\circ}\text{C}$ and 70% relative humidity. During the experiment, individuals were examined regularly through dissections for signs of and gut contents were smeared and stained with Giemsa stain each day to identify the occurrence of sporonts stage of the *Pyxinia firma* in the gut and their haemolymph was collected through piercing the abdomen of the fourth instar larvae by a micro pipette and Neubauer counting chamber was used to calculate their total haemocyte counts. The location of the parasites was determined by the histological and surgical study of the whole body. Control groups were obtained by adding 8% w/w sorbic acid into the diets of a selection of the colony.

A statistical analysis was undertaken of the THC among the various larval and adult stages of multiple generations within the colony with the counts calculated through the Neubauer chamber being analysed using the SPSS Software Package Version 12.0 to perform a Least Significant Difference (LSD) test and to obtain significant values where ($p<0.05$). In addition, the degree of weight loss was recorded for larvae between the fourth and fifth instar in the five replicates of 120 infected and uninfected each. This weight loss data were calculated with standard deviations.

RESULTS

The average Total Counts (THC) between the fourth and fifth instars of treated larvae of the *D. vulpinus*




Fig. 1: The pattern of variation in the THC at different larval and adult stages of infected and gregarine free *D. vulpinus*




Fig. 3: The weight gain in fed larvae and the weight loss in starved fourth instar larvae of *D. vulpinus*




Fig. 2: Photomicrographs showing different forms of *P. firma*; E = Epimerite; P = Protomerite and D = Deutomerite

ranged between 4500-5000/mm³ with a mean of 4750±90. An approximately two fold higher Total Haemocyte Counts (THC) was recorded in infected (untreated) larvae between the fourth and fifth instar than in the treated larvae at a similar developmental stage with a range between 9000 and 11 000/mm³. The first progeny of treated larvae exhibited a higher THC than that in the treated larvae with the second progeny exhibiting higher THC still (although still lower than untreated larvae (Fig. 1 and 2).

These differences in the THC levels are reflective of the organism's active role on the immune system of the




Fig. 4: Photographs of fourth instar larvae of *D. vulpinus*; a) Starved and infected; b) fourth, starved and infection-free

beetle. Starvation of infected larvae had the effect of reducing their weight three times faster than starved eugregarine free larvae after only 3 weeks from the commencement of a starvation diet (Fig. 3). This rate of weight loss slowed down, however in the next two generations of treated specimens (Fig. 3 and 4). Strikingly when one considers these dramatic differences in weight between different categories of starved specimens, no significant difference was observed in the degree of weight gain between infected and uninfected groups of fed specimens. Histological examination, meanwhile, revealed on lesions in the epithelial layer of the mid gut.

DISCUSSION

P. firma (Leger, 1892) is known to be an inhabitant of *Dermestes frischli* while *D. vulpinus* is known as a host of *Pyxinia crystalligera*. Identification of the species in this instance was provided by The Natural History Museum, London. The results suggest that the impact of infection is manifested most obviously when larvae of *D. vulpinus* were starved. Starved larvae that were infected exhibited a much faster rate of weight loss than those that were gregarine free. These observations are compatible with those of Dunkel and Boush (1969) in respect to the effect of *P. frenzli* on the carpet beetle, *Attagenus megatoma*.

The results also indicate a higher THC in infected larvae than in uninfected larvae which conforms with the findings of Weiser and Beard (1959) in respect to the effect of infection by *Seruces* this on scarab beetle larvae.

The elevation in the THC and the rate of weight loss in infected compared to gregarine free larvae becomes less significant if larvae are treated with sorbic acid. The toxic effect of the sorbic acid, however, appears to subside with successive generations since both the THC and the rate of weight loss begin to rise again the following progeny. This tends to support the argument of Trehan and Pajni (1961) that poison causes a reduction in the total haemocyte counts in infected insects. The THC in *Locusta migratoria* (Webley, 1951) and in *P. americana* (Wigglesworth, 1955; Jones, 1964) for example increased at each mout during the nymphal period to justify their role in the immune system (Ratcliffe *et al.*, 1976; Dean *et al.*, 2004; Qamar and Jamal, 2009). In fact, insects and other arthropods present different physico-chemical methods to combat and, check the challenges posed by their biological enemies including viruses, bacteria, protozoans, fungi and cestods (Ratcliffe *et al.*, 1976; Wood and Jacinto, 2007).

Since, no injuries were caused to the site of inhabitation and on account of the fact that normally fed larvae showed little signs of infection other than elevated THC compared to uninfected larvae, the results would tend to suggest that *P. firma* in *D. vulpinus* should be considered to be more a commensal than a parasite.

CONCLUSION

Pyxinia firma is a new record in the midgut of the scavenger beetle, *Dermestes vulpinus*. It shows a negative effect on the growth of the larvae of the beetle however only when these are starved. Infected specimens exhibit a significant increase in the total haemocyte counts and these counts also rise in the progeny of treated larvae, presumably as an indication of immunological response.

ACKNOWLEDGEMENTS

Researchers are indebted to Dr. Warren of the protozoa section of Natural History Museum London, for his strenuous efforts to help find an expert on gregarines and also to Professor Theodris for his kind help in identifying the specimen. Researchers extend their appreciation to the Deanship of Scientific Research at King Saud University for funding the research through the research group project No. RGP-VPP.

REFERENCES

- Al-Khalifa, M.S., 1984. The transovarial transmission of symbionts in the grain weevil, *Sitophilus granarius*. J. Invertebr. Pathol., 44: 106-108.
- Brook, M.W., 1974. Protozoan Infections. In: Insect Diseases, Cantwell, G.E. (Ed.). Marcel Dekker Inc., New York, pp: 219-241.
- Dean, P., E.H. Richards, J.P. Edwards, S.E. Reynolds and K. Chamley, 2004. Microbial infection causes the appearance of hemocytes with extreme spreading ability in monolayers of the tobacco hornworm *Manduca sexta*. Dev. Comp. Immunol., 28: 689-700.
- Dunkel, F., P.Z. Lung, L. Chuan and H.F. Yin, 1982. Insect and fungal response to sorbic acid-treated wheat during storage in south China. J. Econ. Entomol., 75: 1083-1088.
- Dunkel, F.V. and G.M. Boush, 1969. Effect of starvation on the black carpet beetle, *Attagenus megatoma*, infected with the eugregarine *Pyxinia frenzeli*. J. Invertebr. Pathol., 14: 49-52.
- Harry, O.G., 1967. The effect of a eugregarine *Gregarina polymorpha* (Hammerschmidt) on the mealworm larva of *Tenebrio molitor* (L.). J. Protozool., 14: 539-547.
- Jones, J.C., 1964. Differential haemocyte counts from unfixed last stage *Galleria mellonella*. Am. Zool., 4: 337-346.
- Kudo, R.R., 1966. Protozoology. 5th Edn., Charles C. Thomas, Illinois, USA., Pages: 1174.
- Qamar, A. and K. Jamal, 2009. Differential haemocyte counts of 5th instar nymphs and adults of *Dysdercus cingulatus* Fabr (Hemiptera: Pyrrhocoridae) treated with acephate, an organophosphorus insecticide. Biol. Med., 1: 116-121.
- Ratcliffe, N.A., S.J. Gagen, A.F. Rowley and A. Schmit, 1976. The role of granular hemocytes in the cellular defense reactions of the waxmoth, *Galleria mellonella*. Proceedings of the 6th European Congress on Electron Microscopy, September 14-20, 1976, Jerusalem, Israel, pp: 295-297.

- Trehan, K.N. and H.R. Pajni, 1961. Mode of action of insecticides. 1. Observations of the effect of endrin in blood cell count of some insects. *Beitraege Entomologie Berlin*, 11: 1-11.
- Webley, D.P., 1951. Blood cell counts in the American migratory locust, *Locusta migratoria migratorioides* reich and fairmaire. *Proc. R. Ent. Soc. London*, 26: 25-37.
- Weiser, J. and R.L. Beard, 1959. *Adelina sericesthis* n. sp., a new coccidian parasite of scarabaeid larvae. *J. Insect Pathol.*, 1: 99-106.
- Wigglesworth, V.B., 1955. The role of the haemocytes in the growth and moulting of an insect, *Rhodnius prolixus* (Hemiptera). *J. Exp. Biol.*, 32: 649-663.
- Wood, W. and A. Jacinto, 2007. *Drosophila melanogaster* embryonic haemocytes: masters of multitasking. *Nat. Rev. Mol. Cell Biol.*, 8: 542-551.