

1. Recognized feature of Sheehan syndrome.

- A. Menorrhagia
- B. Galactorrhoea
- C. Insulin resistance
- D. Hypothyroidism
- E. Dwarfism

2. Which of the following items may be associated with a mid trimester abortion.

- A. Recurrent vaginal infections
- B. Maternal smoking
- C. Cervical incompetence
- D. Sickle cell disease
- E. Hyperemesis gravidarum

3. In complete hydatidiform mole

- A. Rarely contain maternal genetic material
- B. Can present with hypothyroidism
- C. It is an indication for prophylactic cytotoxic therapy
- D. Diagnosis is confirm with a very low blood level of human beta chorionic gonadotrophin
- E. Pre-eclampsia is the most common symptom and occur in 97% of patients.

4. Feature of polycystic as ovarian syndrome

- A. Elevated level of serum FSH
- B. Endometrial hyperplasia
- C. Breast atrophy
- D. Polymenorrhoea
- E. Regular ovulation

5. The following disorders are correctly linked with a recognized association.

- A. Klinefelter syndrome 47 XYY
- B. Polycystic ovarian syndrome high progesterone level
- C. Androgen insensitivity syndrome Hirsutism
- D. Turner syndrome streak ovaries
- E. Ovarian failure low FSH and LH

6. Serum alpha fetoprotein concentration is raised in

- A. Trisomy 21
- B. Women taking oral contraceptive pills
- C. After threatened abortion
- D. In presence of closed spina bifida
- E. Fetus with congenital heart disease

7. The following one associated with an increased risk of fetal malformation.

- A. Single umbilical vein
- B. Paternal age over 50 years
- C. Maternal smoking
- D. Maternal chorionic alcoholism
- E. Pre-eclamptic toxemia (P E T)

8. After menopause

- A. There is increase vaginal acidity
- B. Gonadotrophins level falls
- C. There is increase in bone density
- D. Hormonal replacement therapy should be given routinely to all menopausal women.
- E. Any post menopausal bleeding should be investigated

9. The following are always an indication for caesarean section.
- A. Grade IV placenta preevia
 - B. Abruptio placenta
 - C. Cervical cerclage
 - D. Breech presentation
 - E. Twins pregnancy
10. Feature characteristically associated with an imperforate hymen in a girl aged 16 years includes:
- A. Absence of secondary sexual characteristic
 - B. Acute retention of urine
 - C. Closed spina bifida
 - D. Short stature
 - E. Present with secondary amenorrhoea
11. Infants of diabetic mothers are at risk of one of the following:
- A. Low haemoglobin
 - B. Hypercalcaemia
 - C. Hyperglycemia
 - D. Microsomia
 - E. Respiratory distress syndrome
12. Indication of glucose tolerance test GTT in pregnancy include all of the following EXCEPT:
- A. Previous history of gestational diabetes mellitus.
 - B. History of macrosomic baby
 - C. Glycosuria in one occasion
 - D. History of unexplained fetal death
 - E. History of babies with congenital sacral agenesis