



Qualitative Tests of Amino Acids


BCH303 [Practical]

Amino Acids:


- Amino acids role.
- There are 20 natural amino acids that are found within proteins.
→ **All of them are L- α amino acids.**



General structure of amino acids:




Classification of amino acids:




Twenty standard amino acids


Nonpolar, aliphatic R groups




Aromatic R groups




Positively charged R groups



Polar, uncharged R groups



Negatively charged R groups




Some properties of Amino Acids:

1. Amphoteric Compounds.
2. Isoelectric point (pI).
3. Optical Activity.
4. Light Absorption.


Some properties of Amino Acids:

1. Amphoteric Compounds:

- What is **Amphoteric** compounds ?
- Amphoteric properties of amino acids due to the presence of their **ionizable α -amino and α -carboxylic group** can act sometimes as acids and sometimes as bases **depending on the pH of their media** .




A. Presence of **carboxyl group** COOH that able to **donate** proton (H⁺) “acidic behavior”, and converted to COO⁻ :



High pH → Act as **acids**

B. Presence of **amino group** NH₂ that able to **accept** proton (H⁺) “basic behavior”, and converted to NH₃⁺ :




Low pH → Act as **Base**

Some properties of Amino Acids cont':

2. Isoelectric point (pI):



- It is the pH value at which the **positive charge equals** the **negative charge** (i.e. **the net charge of this molecule equals zero**) → **Zwitter ion**
- Electric field?
- Solubility?**
- Different pI.



Some properties of Amino Acids cont':

3. Optical Activity :


- Amino acids are able to rotate polarized light either to:
 - The left (**Levorotatory**) → (-) – Amino acid
 - The right (**Dextrorotatory**) → (+) – Amino acid
- **Asymmetric C atom.**
- **Glycine ?**



Some properties of Amino Acids cont':

4. Light Absorption:

- The aromatic amino acids absorb ultraviolet light at **280nm**.
- proteins ?



Practical part

Qualitative tests of amino acids

1 Solubility Test.

2 Ninhydrin test: for α -L amino acids.

3 Xanthoproteic test: for Aromatic amino acids.

4 Sakaguchi Test: for arginine.

5 Millon's test: for amino acids containing hydroxy phenyl group (Tyrosine)

6 Lead sulfite test: for of amino acids containing sulfhydryl group (- SH) (Cysteine)

Experiment 1 : Solubility Test

Objective:

- Investigate the solubility of selected amino acid in various solutions.

Principle:

- Amino acids are generally **soluble in water** and **insoluble in non-polar organic solvents** such as hydrocarbons.
- This is because the presence of **amino and carboxyl group**.


Experiment 1 : Solubility Test

Method:

1. Add 2ml of different solvents in 3 clean test tubes then place 0.5 ml of each amino acid.
2. Shake the tubes thoroughly, then leave the solution for about one minute.
3. Notice what happened to the solution .
4. Record your result .

Results:

Amino acid	Solvent	Degree of solubility
Glycine	Water	
	NaOH	
	HCl	
	Chloroform	
Arginine	Water	
	NaOH	
	HCl	
	Chloroform	
Glutamine	Water	
	NaOH	
	HCl	
	Chloroform	



soluble





insoluble

Experiment 2: Ninhydrin test

Objective:

- To detect α -L-amino acids.


Principle:



Experiment 2: Ninhydrin test

Principle cont':

- All amino acids that have a free amino group will give (purple color).
- While not free amino group-proline and hydroxy-proline (imino acids) will give a (yellow color) .



Note:

All primary amines and ammonia react similarly and produce blue/purple product but without the liberation of carbon dioxide.

Experiment 2: Ninhydrin test

Method:

- 1-Place 1 ml of each of the solutions in a test tube and add 1 ml of ninhydrin solution.
- 2- Boil the mixture over a water bath for 2 min.
- 3- Allow to cool and observe the blue-purple color formed.
- 4- Record your results.

Results:

Tube	Observation
Glycine	
Tryptophan	
Proline	

! CAUTION

Ninhydrin is a strong oxidizing agent, it should be handled with care.



Experiment 3 : Xanthoproteic test


Objective:

- To differentiate between **aromatic amino acids** which give **positive** results and other amino acids.

Principle:

- In the presence of concentrated nitric acid (HNO_3), the aromatic phenyl ring is nitrated to give yellow colored nitro-derivatives , [**nitration reaction**] → giving the solution yellow color.
- Amino acids **tyrosine** and **tryptophan** → contain activated benzene rings.
- **Phenylalanine ?**

Nitration



benzene ring is not activate

Experiment 3 : Xanthoproteic test

Method:


1. Label four tubes (1 - 4), then add 1 ml of each amino acid solutions and phenol solution to those test tubes each alone.
2. Add 1 ml of concentrated HNO_3 . then record your result
3. Now COOL THOROUGHLY under the tap and CAUTIOUSLY add 5 drops of 10M NaOH to make the solution strongly alkaline (the alkaline is added to be sure about the nitration).

! CAUTION

Concentrated HNO_3 is a toxic, it should be handled with care.

Results:

Tube	Observation	
	+ HNO_3	+NaOH
Tyrosine		
Tryptophan		
Phenylalanine		
Phenol		




Experiment 4 : Millon's test

Objective:

- This test is specific for **Tyrosine** → because it is the only amino acid containing a **phenol group**.

Principle:




Note:

All phenols (compound having benzene ring and OH attached to it) give positive results in Millon's test.

Experiment 5 : Sakaguchi Test


Objective:

- Detection of amino acid containing **guanidinium group** → test for **Arginine**.



Principle:

- In **alkaline** solution, arginine react with α -naphthol and sodium hypobromite /chlorite as an oxidize agent, to form **red** complexes as a positive result.



guanidinium group


Experiment 5 : Sakaguchi Test

Method:

1. Label 2 test tube and place in each one 2 ml of the amino acid solution .
2. Add to each tube 2ml of NaOH solution. Mix well
3. Add to each tube 5 drops of α -naphthol solution. Mix well
4. Add to each tube 5 drops of sodium hypobromite solution, and record your result .

Results:

Tube	Observation
Glycine	
Arginine	



Experiment 6 : Lead Sulfite Test

Objective:

- This test is specific for **-SH [sulfhydryl group]** containing amino acid → **Cysteine**.

Principle:

- Sulphur in **cysteine**, is converted to **sodium sulfide** by boiling with 40% NaOH.
- The Na_2S can be detected by the precipitation of **PbS** (lead sulfide) from an alkaline solution when adding lead acetate $(\text{CH}_3\text{COO})_2\text{Pb}$.

