

Cloning and its Applications

Islam M. Saadeldin

Department of Physiology, Faculty of Veterinary Medicine, Zagazig University, Zagazig 44519, Egypt

Corresponding author: Islam M. Saadeldin, Department of Physiology, Faculty of Veterinary Medicine, Zagazig University, 44519 Zagazig, Egypt, Tel: +20-11-0002-4182; Fax: +20-55-228-3683; E-mail: islamms@zu.edu.eg; islamms@snu.ac.kr

Rec date: 18 May 2015; **Acc date:** 20 May 2015; **Pub date:** 22 May 2015

Copyright: © 2015 Saadeldin MI. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Editorial

In mammalian fertilization process, female pronucleus share male pronucleus to form the zygote which undergoes mitotic division yielding totipotent cells. After morula compaction, the embryonic cells differentiate into two types of cells; the inner cell mass cells which have the pluripotent stemness criteria to form the embryo proper, and the trophectoderm which would share the placenta formation.

In vitro -without spermatozoa contribution and using somatic cells instead- the oocyte was found to have extraordinary astonishing capabilities to process the genetic materials of transferred somatic cells through different mechanisms;

1) Reprogramming the somatic cells even when it is enucleated, as in case of somatic cell nuclear transfer (cloning) which produces a copy of genetic materials of the somatic cell donor animal [1];

2) Enforcing the somatic cells nuclei to bypass an obligatory mitotic cell cycle checkpoints and undergo premature condensation [2]; and

3) Enforcing the somatic cells for entry of meiosis through the reduction division as known by semi-cloning or haploidization [3,4].

Moreover, it was found that the early meiotic ooplasm was able to induce initiation of a meiosis-like reducing division in mitotic nuclei originating from differentiated somatic cells [5]. Thus, the meiotic machinery property of the ooplasm can be helpful for generation of artificial oocytes or male germ cells through injecting male and/or female somatic cells into enucleated mature or immature oocytes [6-8].

Since the birth of Dolly the sheep, the first cloned mammal, an ever-growing number of studies worldwide have helped to substantiate the potential applications of somatic cell nuclear transfer (cloning) to overcome several problems in various biology fields such as:

Reproductive cloning: To generate copies of particular species including extinct or endangered species and for propagation of the livestock and elite animals.

Therapeutic cloning: The elite cloned embryos would be a suitable source for pluripotent embryonic stem cells that capable for differentiation into different organs for regenerative medicine purposes to avoid the transplant rejection.

Transgenesis: Production of genetically modified organisms for increasing production of specific merit (such as milk, wool and meat...

etc); model animals for human diseases (Parkinson's disease and Alzheimer diseases..etc) and generating humanized organs for xenotransplantation purposes.

However, the use of this technique is limited owing to its low efficiency and several trials showed the improvement of cloning efficiencies [9].

Therefore, the current issue of "Cloning & Transgenesis" will focus on cloning and semi-cloning regarding the trials to improve their efficiency and to reveal their potential applications in biology and medicine.

References

1. Wilmut, I, Schnieke AE, McWhir J, Kind AJ, Campbell KH (1997) Viable offspring derived from fetal and adult mammalian cells. *Nature* 385(6618): 810-813.
2. Szollosi D, Czolowska R, Soltynska MS, Tarkowski AK (1986) Ultrastructure of cell fusion and premature chromosome condensation (PCC) of thymocyte nuclei in metaphase II mouse oocytes. *Biol Cell* 56: 239-249.
3. Takeuchi T, Kaneko M, Veeck LL, Rosenwaks Z, Palermo GD (2001) Creation of viable human oocytes using diploid somatic nuclei. Are we there yet? (Abstract O-011). *Human Reproduction* 16: 5.
4. Tesarik J (2002) Reproductive semi-cloning respecting biparental embryo origin: embryos from syngamy between a gamete and a haploidized somatic cell. *Hum Reprod* 17: 1933-1937.
5. Kubelka M, Moor RM (1997) The behaviour of mitotic nuclei after transplantation to early meiotic ooplasts or mitotic cytoplasts. *Zygote* 5: 219-227.
6. Tesarik J, Nagy ZP, Sousa M, Mendoza C, Abdelmassih R (2001) Fertilizable oocytes reconstructed from patient's somatic cell nuclei and donor ooplasts. *Reprod Biomed Online* 2: 160-164.
7. Trounson A (2001) Nuclear transfer in human medicine and animal breeding. *Reprod Fertil Dev* 13: 31-39.
8. Tsai MC, Takeuchi T, Bedford JM, Reis MM, Rosenwaks Z, et al. (2000) Alternative sources of gametes: reality or science fiction? *Hum Reprod* 15: 988-998.
9. Saadeldin IM, Kim SJ, Choi YB, Lee BC (2014) Improvement of cloned embryos development by co-culturing with parthenotes: a possible role of exosomes/microvesicles for embryos paracrine communication. *Cellular reprogramming* 16: 223-234.