

King Saud University
Department Of Mathematics.
M-203 (First Mid-Term)
(Differential and Integral Calculus)

(Summer Semester 1433/1434)

Max. Marks: 25

Time: 1.30 hrs

Marking Scheme: All questions carry equal marks.

Q. No: 1 Discuss the convergence of the sequence $\left\{ \frac{1}{n} \sin \left(\frac{1}{n^2} \right) \right\}$.

Q. No: 2 Find the sum of the series $\sum_{n=1}^{\infty} \left[\frac{1}{(n+1)(n+2)} + \frac{1}{e^n} \right]$.

Q. No: 3 Find the **interval of convergence** and **radius of convergence** of the power series $\sum_{n=1}^{\infty} (-2)^n \frac{1}{\sqrt{n}} (x+3)^n$.

Q. No:4 Determine whether the series is **absolutely convergent**, **conditionally convergent**, or **divergent** $\sum_{n=2}^{\infty} \frac{(-1)^n}{n \ln(n)}$.

Q. No:5 Find the Taylor series for the function $f(x) = \cos x$ at $c = \pi$.

Q#3) Find the Interval of convergence and radius of cong. of the power series $\sum_{n=1}^{\infty} \frac{(-2)^n}{\sqrt{n}} (x+3)^n$. [Marks: 5]

Sol. $\lim_{n \rightarrow \infty} \left| \frac{(-2)^{n+1}}{\sqrt{n+1}} (x+3)^{n+1} \div \frac{(-2)^n}{\sqrt{n}} (x+3)^n \right|$
 $= 2|x+3|$

For abs. cong. $2|x+3| < 1 \Rightarrow |x+3| < \frac{1}{2}$
 $(\Rightarrow) -\frac{1}{2} < x+3 < \frac{1}{2}$
 $-\frac{7}{2} < x < -\frac{5}{2}$ (2)

At $x = -\frac{7}{2}$, we have $\sum_{n=1}^{\infty} \frac{(-2)^n}{\sqrt{n}} \left(-\frac{1}{2}\right)^n = \sum_{n=1}^{\infty} \frac{1}{\sqrt{n}}$ which is a divg. p-series.

At $x = -\frac{5}{2}$, we have $\sum_{n=1}^{\infty} \frac{(-2)^n}{\sqrt{n}} \left(\frac{1}{2}\right)^n = \sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt{n}}$ (1)
which is cong. by AST. (1)

Hence Interval of cong: $[-\frac{7}{2}, -\frac{5}{2}]$
Radius of cong: $r = \frac{1}{2}$ (1)

Q#4) Determine whether the series is absolutely cong, cond. cong, or divergent: $\sum_{n=2}^{\infty} \frac{(-1)^n}{n(\ln(n))}$ [Marks: 5]

Sol. By AST, $\sum_{n=2}^{\infty} \frac{(-1)^n}{n(\ln(n))}$ is cong. (2)

$\sum_{n=4}^{\infty} \left| \frac{(-1)^n}{n(\ln(n))} \right| = \sum_{n=2}^{\infty} \frac{1}{n(\ln(n))}$ is divg. by Integral test (to verify the hypotheses of the integral test)

$\lim_{t \rightarrow \infty} \int_2^t \frac{1}{x \ln(x)} dx = \lim_{t \rightarrow \infty} \left[\ln x \right]_2^t$ Put $\ln x = u \Rightarrow \frac{1}{x} dx = du$
 $= \lim_{t \rightarrow \infty} [\ln t - \ln 2] = \int \frac{1}{u} du = \ln(u)$
 $= \infty$; divg. (2)

Hence, $\sum_{n=2}^{\infty} \frac{(-1)^n}{n(\ln(n))}$ is cond. cong. (1)

Q #5) Find the Taylor Series for the function $f(x) = \cos x$ at $c = \pi$ [Marks: 5]

Soln. we have, the Taylor Series

$$f(x) = f(c) + f'(c)(x-c) + \frac{f''(c)}{2!}(x-c)^2 + \frac{f'''(c)}{3!}(x-c)^3 + \dots + \frac{f^{(n)}(c)}{n!}(x-c)^n + \dots$$

Here $f(x) = \cos x \Rightarrow f(\pi) = \cos \pi = -1$

$f'(x) = -\sin x \Rightarrow f'(\pi) = 0$

$f''(x) = -\cos x \Rightarrow f''(\pi) = 1$

$f'''(x) = \sin x \Rightarrow f'''(\pi) = 0$

$f^{(4)}(x) = \cos x \Rightarrow f^{(4)}(\pi) = -1$

Substituting these values in (1), we get

$$f(x) = \cos x = -1 + \frac{(x-\pi)^2}{2!} - \frac{(x-\pi)^4}{4!} + \dots + (-1)^{n+1} \frac{(x-\pi)^{2n}}{2n!}$$

$$= \sum_{n=0}^{\infty} (-1)^{n+1} \frac{(x-\pi)^{2n}}{(2n!)} \quad (3)$$