

King Saud University

College of Sciences

Department of Mathematics

151 Math Exercises

(2)

The Universal Quantifiers

By:

Malek Zein AL-Abidin

س1440

2018

The Universal Quantifier

DEFINITION 1 The *universal quantification* of $P(x)$ is the statement

“ $P(x)$ for all values of x in the domain.”

The notation $\forall x P(x)$ denotes the universal quantification of $P(x)$.

Here \forall is called the *universal quantifier*.

We read $\forall x P(x)$ as “for all $x P(x)$ ” or “for every $x P(x)$ ”

An element for which $P(x)$ is false is called a *counterexample* of $\forall x P(x)$.

Note that When all the elements in the domain can be listed —say, x_1, x_2, \dots, x_n —

it follows that the universal quantification $\forall x P(x)$ is the same as the conjunction

$$P(x_1) \wedge P(x_2) \wedge \dots \wedge P(x_n)$$

because this conjunction is true if and only if $P(x_1), P(x_2), \dots, P(x_n)$ are all true.

The Existential Quantifier

DEFINITION 2 The *existential quantification* of $P(x)$ is the proposition

“There exists an element x in the domain such that $P(x)$.”

We use the notation $\exists x P(x)$ for the existential quantification of $P(x)$.

Here \exists is called the *existential quantifier*.

Note that when all the elements in the domain can be listed —say, x_1, x_2, \dots, x_n —

it follows that the universal quantification $\exists x P(x)$ is the same as the disjunction

$$P(x_1) \vee P(x_2) \vee \dots \vee P(x_n),$$

because this disjunction is true if and only if at least one of

$$P(x_1), P(x_2), \dots, P(x_n) \text{ is true.}$$

Statement	When True?	When False?
$\forall x P(x)$	$P(x)$ is true for every x	There is an x for which $P(x)$ is false
$\exists x P(x)$	There is an x for which $P(x)$ is true	$P(x)$ is false for every x

Negating Quantified Expressions

Negation	Equivalent Statement	When Is Negation True?	When False?
$\neg \exists x P(x)$	$\forall x \neg P(x)$	For every x $P(x)$ is false	There is an x for which $P(x)$ is true
$\neg \forall x P(x)$	$\exists x \neg P(x)$	There is an x for which $P(x)$ is false	$P(x)$ is true for every x

EXAMPLE 1 Let $Q(x)$ be the statement “ $x < 2$.” What is the truth value of the quantification $\forall x Q(x)$, where the domain consists of all real numbers?

Solution: $Q(x)$ is not true for every real number x , because, for instance, $Q(3)$ “ $3 < 2$ ” is false. That is, $x = 3$ is a counterexample for the statement $\forall x Q(x)$.

Thus $\forall x Q(x)$ is false.

EXAMPLE 2 Suppose that $P(x)$ is “ $x^2 > 0$ ”. To show that the statement $\forall x P(x)$ is false where the universe of discourse consists of all integers, we give a counterexample. We see that $x = 0$ is a counterexample because $x^2 = 0$ when $x = 0$, so that x^2 is not greater than 0 when $x = 0$ ($x^2 = 0 \not> 0$)

EXAMPLE 3 What is the truth value of $\forall x P(x)$, where $P(x)$ is the statement “ $x^2 < 10$ ” and the domain consists of the positive integers not exceeding 4?

Solution: The statement $\forall x P(x)$ is the same as the conjunction $P(1) \wedge P(2) \wedge P(3) \wedge P(4)$ because the domain consists of the integers 1, 2, 3, and 4. Because $P(4)$, which is the statement “ $4^2 < 10$,” is false, ($4^2 = 16 \not< 10$) it follows that $\forall x P(x)$ is false.

EXAMPLE 4 What is the truth value of $\forall x (x^2 \geq x)$ if the domain consists of all real numbers? What is the truth value of this statement if the domain consists of all integers?

Solution: The universal quantification $\forall x (x^2 \geq x)$, where the domain consists of all real numbers, is false. For example, $(\frac{1}{2})^2 = \frac{1}{4} \not\geq \frac{1}{2}$.

Note that $x^2 \geq x \Leftrightarrow x^2 - x = x(x - 1) \geq 0 \Leftrightarrow x \leq 0$ or $x \geq 1$

It follows that $\forall x (x^2 \geq x)$ is false if the domain consists of all real numbers (because the inequality is false for all real numbers x with $0 < x < 1$). However, if the domain consists of the integers, $\forall x (x^2 \geq x)$ is true, because there are no integers x with $0 < x < 1$.

EXAMPLE 5 Let $P(x)$ denote the statement “ $x > 3$.” What is the truth value of the quantification $\exists x P(x)$, where the domain consists of all real numbers?

Solution: Because “ $x > 3$ ” is sometimes true—for instance, when $x = 4$ —the existential quantification of $P(x)$, which is $\exists x P(x)$, is true.

EXAMPLE 6 Let $Q(x)$ denote the statement “ $x = x + 1$.” What is the truth value of the quantification $\exists x Q(x)$, where the domain consists of all real numbers?

Solution: Because $Q(x)$ is false for every real number x , the existential quantification of $Q(x)$, which is $\exists x Q(x)$, is false.

EXAMPLE 7 What is the truth value of $\exists x P(x)$, where $P(x)$ is the statement “ $x^2 > 10$ ” and the universe of discourse consists of the positive integers not exceeding 4?

Solution: Because the domain is $\{1, 2, 3, 4\}$, the proposition $\exists x P(x)$ is the same as the disjunction $P(1) \vee P(2) \vee P(3) \vee P(4)$.

Because $P(4)$, which is the statement “ $4^2 > 10$,” is true, it follows that $\exists x P(x)$ is true.

EXAMPLE 8 What are the negations of the statements $\forall x (x^2 > x)$ and $\exists x (x^2 = 2)$?

Solution: (i) The negation of $\forall x (x^2 > x)$ is the statement $\neg \forall x (x^2 > x)$, which is equivalent to $\exists x \neg (x^2 > x)$. This can be rewritten as $\exists x (x^2 \leq x)$.

(ii) The negation of $\exists x (x^2 = 2)$ is the statement $\neg \exists x (x^2 = 2)$, which is equivalent to $\forall x \neg (x^2 = 2)$. This can be rewritten as $\forall x (x^2 \neq 2)$. The truth values of these statements depend on the domain.

EXAMPLE 9 Suppose that $Q(x)$ is “ $x^2 \geq 2x$ ”, where x is an integer.

(i) What is the negation of $\exists x Q(x)$?

Solution:

$$\forall x \in \mathbb{Z}, x^2 < 2x$$

(ii) What is the truth value of $\forall x Q(x)$? Justify your answer.

Solution:

False, take $x = 1$

(iii) What is the truth value of $\exists x Q(x)$? Justify your answer.

Solution:

True, take $x = 1$

EXAMPLE 10

Determine the truth value of each of the statements below given that the domain of each variable is the set of real numbers.

(i) $\exists x, (x^2 = 2)$

(ii) $\forall x, (x^2 \neq x)$.

Solution:

(i) true , because $(\sqrt{2})^2 = 2$

(ii) false , because $(1)^2 = 1$.

Exercises

Q₁. Let $P(x)$ be the statement “ $x = x^2$.” If the domain consists of the integers, what are these truth values?

a) $P(0)$

b) $P(1)$

c) $P(2)$

d) $P(-1)$

e) $\exists x P(x)$

f) $\forall x P(x)$

Q₂. Let $Q(x)$ be the statement “ $x + 1 > 2x$.” If the domain consists of all integers, what are these truth values?

a) $Q(0)$

b) $Q(-1)$

c) $Q(1)$

d) $\exists x Q(x)$

e) $\forall x Q(x)$

f) $\exists x \neg Q(x)$

g) $\forall x \neg Q(x)$

Q₃. Determine the truth value of each of these statements if the domain consists of all integers.

a) $\forall n (n + 1 > n)$

b) $\exists n (2n = 3n)$

c) $\exists n (n = -n)$

d) $\forall n (3n \leq 4n)$

Q₄. Determine the truth value of each of these statements if the domain consists of all real numbers.

a) $\exists x (x^3 = -1)$

b) $\exists x (x^4 < x^2)$

c) $\forall x ((-x)^2 = x^2)$

d) $\forall x (2x > x)$

Q₅. Determine the truth value of each of these statements if the domain consists of all integers.

a) $\forall n (n^2 \geq 0)$

b) $\exists n (n^2 = 2)$

c) $\forall n (n^2 \geq n)$

d) $\exists n (n^2 < 0)$

Q₆. Determine the truth value of each of these statements:

(1) $\forall x \in \mathbb{R}, x^2 - 4x + 4 \geq 0$

(2) $\forall x > 0, x \geq \frac{1}{x}$

(3) $\forall x \in \mathbb{Z}, ((x \geq 2) \vee (x^2 \leq 2))$

(4) $\exists x \in \{1,2,3,4\}, 2^x < x!$

$$(5) \exists x \in \mathbb{Z}^* = \mathbb{Z} - \{0\}, \frac{x-1}{x} \in \mathbb{Z}$$

$$(6) \exists x \in \mathbb{R}, x^2 = 5$$

Q₇ Write *the negation* of the below statements:

(i) Some students did not listen to the instructions.

(ii) $\exists x \in D, x^2 > 3$

(iii) If you collect enough points, you will win the game.