

# Fundamentals of Organic Chemistry

## CHEM 109

*For Students of Health Colleges*

Credit hrs.: (2+1)

*King Saud University*

College of Science, Chemistry Department

### CHAPTER 4: Alcohols, Phenols and Ethers

1

## Alcohols, Phenols and Ethers

- Alcohols, phenols and ethers may be viewed as organic *derivatives of water*.


H-O-H	R-O-H	Ar-O-H	R-O-R	R-O-Ar	Ar-O-Ar
Water	Alcohol	Phenol	Ether		

- Alcohols and phenols have a common functional group, *the hydroxyl group, -OH*.
- Alcohols are compounds whose molecules have a hydroxyl group attached to a *saturated carbon* atom.
- Phenols are compounds that have a hydroxyl group attached directly to a *benzene ring*.
- Ethers are compounds whose molecules have an oxygen atom bonded to *two carbon atoms*.

2

## Classification of Alcohols and Ethers

- Alcohols are classified as primary (1°), secondary (2°), or tertiary (3°), depending on whether one, two, or three organic groups are connected to the hydroxyl-bearing carbon atom.




- Methyl alcohol*, which is not strictly covered by this classification, is usually grouped with the primary alcohols.
- Ethers are classified as
  - Symmetrical ethers;
    - When the organic groups attached to the oxygen are identical.
  - Unsymmetrical ethers (*mixed ethers*);
    - When the organic groups attached to the oxygen are different.

3

## Nomenclature of Alcohols

## Nomenclature

- The **common names** for the simplest alcohols consist of alkyl group attached to the hydroxyl function followed by the word alcohol: Alkyl alcohol.
- In the **IUPAC system**, alcohols are named according to the following rules.
  - Select the **longest continuous carbon chain** that *contains the -OH group*.
    - Drop the *-e* ending of the parent alkane and replace it by the suffix *-ol*: Alkanol
  - When isomers are possible, the chain is numbered so as to give the functional group (-OH) the *lowest possible number*.




4

## Nomenclature of Alcohols


## Nomenclature

3. When alkyl side chains or other groups are present; they are named alphabetically and their positions are indicated by a number.

The position of the functional group (-OH) is always given the *lowest possible number* at the end of the name.



For cyclic alcohols, numbering always starts from the carbon bearing the -OH group.




5

## Nomenclature of Alcohols

## Nomenclature

4. With Unsaturated Alcohols; If a molecule contains both an -OH group and a C=C or C-C triple bond, the -OH group takes preference before the double or triple bonds in getting the lower number.

The name should include (if possible) both the hydroxyl and the unsaturated groups, *even if this does not make the longest chain the parent hydrocarbon*.




6

## Nomenclature of Alcohols

## Nomenclature

- Alcohols with More Than One Hydroxyl Group
  - Compounds with two adjacent alcohol groups are called *glycols*.  
The most important example is ethylene glycol.
  - Compounds with more than two hydroxyl groups are also known, and several, such as glycerol and sorbitol, are important commercial chemicals.




- Ethylene glycol* is used as the "permanent" antifreeze in automobile radiators and as a raw material in the manufacture of Dacron.
- Ethylene glycol* is completely miscible with water.
- Glycerol* is a syrupy, colorless, water-soluble, high-boiling liquid with a distinctly sweet taste. Its soothing qualities make it useful in shaving and toilet soaps and in cough drops and syrups.

7


## Nomenclature of Phenols

## Nomenclature

- Phenols are usually named as derivatives of the parent compounds.



- The hydroxyl group is named as a substituent when it occurs in the same molecule with carboxylic acid, aldehyde, or ketone functionalities, which have priority in naming.





8

## Nomenclature of Ethers

## Nomenclature


### Common Names

Ethers are usually named by giving the name of each alkyl or aryl group, in alphabetical order, followed by the word *ether*.

Methyl ether	$\text{CH}_3\text{—O—CH}_3$	Ethyl methyl ether	$\text{CH}_3\text{—O—CH}_2\text{CH}_3$
Ethyl ether	$\text{CH}_3\text{CH}_2\text{—O—CH}_2\text{CH}_3$	Ethyl- <i>n</i> -propyl ether	$\text{CH}_3\text{CH}_2\text{—O—CH}_2\text{CH}_2\text{CH}_3$
Vinyl ether	$\text{CH}_2=\text{CH—O—CH}=\text{CH}_2$	<i>t</i> -Butyl methyl ether	$(\text{CH}_3)_3\text{C—O—CH}_3$
Phenyl ether		Methyl phenyl ether (anisole)	

### IUPAC system

- For ethers with more complex structures, it may be necessary to name the -OR group as an alkoxy group.
- In the IUPAC system, the smaller alkoxy group is named as a substituent.



9

## Physical Properties of Alcohols and Ethers


### Physical State

- The simplest alcohol, methanol, is a liquid at room temperature. In contrast, alkanes from methane to butane are gases.
- Phenol is a colorless, crystalline, and low-melting solid and other phenols also are solids.
- Ethers are colorless compounds with characteristic, relatively pleasant odors.

### Boiling Points

- Ethers have lower boiling points (b.p.s) than alcohols with an equal number of carbon atoms.
- Ether has nearly the same b.p. as the corresponding hydrocarbon in which a -CH<sub>2</sub>- group replaces the ether's oxygen.

Because of their structures (no O-H bonds), ether molecules cannot form hydrogen bonds with one another.




Compound	Formula	bp	mol wt	Water solubility (g/100 mL, 20°C)
1-butanol	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$	118°C	74	7.9
diethyl ether	$\text{CH}_3\text{CH}_2\text{—O—CH}_2\text{CH}_3$	35°C	74	7.5
pentane	$\text{CH}_3\text{CH}_2\text{—CH}_2\text{—CH}_2\text{CH}_3$	36°C	72	0.03

10

## Physical Properties of Alcohols and Ethers

### Boiling Points

- Series of normal alcohols; The boiling points increase with increasing molecular weights.
- A comparison of boiling points among isomeric alcohols; The boiling points decrease as the number of alkyl branches from the carbinol group increases.




- Phenol and most other phenols have high boiling points.

11

## Physical Properties of Alcohols and Ethers

### Solubility

- The lower alcohols are completely miscible with water.
- As the number of carbons in the alcohol increases, the solubility in water decreases.
- Low-molecular-weight ethers, such as dimethyl ether, are quite soluble in water.  
Ether molecules can form hydrogen bonds to water.



Structure	Name	Mol wt	Bp (°C)	Solubility in H <sub>2</sub> O at 20°C
$\text{CH}_3\text{CH}_2\text{CH}_3$	propane	44	-42	insoluble
$\text{CH}_3\text{OCH}_3$	methyl ether	46	-24	soluble
$\text{CH}_3\text{CH}_2\text{OH}$	ethanol	46	78	soluble
$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_3$	n-butane	58	-0.5	insoluble
$\text{CH}_3\text{CH}_2\text{OCH}_3$	ethyl methyl ether	60	8	soluble
$\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$	1-propanol	60	97	soluble
$\text{CH}_3(\text{CH}_2)_3\text{CH}_3$	n-pentane	72	35	insoluble
$\text{CH}_3\text{CH}_2\text{OCH}_2\text{CH}_3$	ethyl ether	74	36	7.5 g/100 g
$\text{CH}_3(\text{CH}_2)_3\text{CH}_2\text{OH}$	1-butanol	74	118	7.9 g/100 g
$\text{CH}_3(\text{CH}_2)_5\text{CH}_3$	n-heptane	100	98	insoluble
$\text{CH}_3(\text{CH}_2)_2\text{O}(\text{CH}_2)_2\text{CH}_3$	n-propyl ether	102	91	0.2 g/100 g
$\text{CH}_3(\text{CH}_2)_4\text{CH}_2\text{OH}$	1-hexanol	102	157	0.6 g/100 g

- Phenol and most other phenols are slightly soluble in water.

12


## Hydrogen Bonding in Alcohols and Ethers

- The **boiling points** (bp's) of alcohols are much higher than those of ethers or hydrocarbons with similar molecular weights.

	CH <sub>3</sub> CH <sub>2</sub> OH	CH <sub>3</sub> OCH <sub>3</sub>	CH <sub>3</sub> CH <sub>2</sub> CH <sub>3</sub>
mol wt	46	46	44
bp	+78.5°C	-24°C	-42°C

Why? Because alcohols form hydrogen bonds with one another.

The O-H bond is polarized by the high electronegativity of the oxygen atom and places a partial positive charge on the hydrogen atom and a partial negative charge on the oxygen atom.



Two or more alcohol molecules thus become loosely bonded to one another through hydrogen bonds.

- Consequently, **alcohols** have relatively high boiling points because they must supply enough heat to break the hydrogen bonds before each molecule.
- Hydrogen bonds are weaker than ordinary covalent bonds.

13

## Hydrogen Bonding in Alcohols and Ethers

- The lower molecular-weight alcohols and ethers can form H-bond with water molecules.
- This accounts for the complete miscibility of the lower alcohols and ethers with water.
- However, as the organic chain lengthens and the alcohol becomes relatively more hydrocarbon like, its water solubility decreases.




Table 7.1 Boiling Point and Water Solubility of Some Alcohols



Name	Formula	bp, °C	Solubility in H <sub>2</sub> O g/100 g at 20°C
methanol	CH <sub>3</sub> OH	65	completely miscible
ethanol	CH <sub>3</sub> CH <sub>2</sub> OH	78.5	completely miscible
1-propanol	CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> OH	97	completely miscible
1-butanol	CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> OH	117.7	7.9
1-pentanol	CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> OH	137.9	2.7
1-hexanol	CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> OH	155.8	0.59

14

## The Acidity of Alcohols and Phenols

- Like water, alcohols and phenols are weak acids.


The hydroxyl group can act as a proton donor, and dissociation occurs in a manner similar to that for water



15

## The Acidity of Alcohols and Phenols

- Phenols are stronger acids than alcohols mainly because the corresponding phenoxide ions are stabilized by resonance.



The negative charge of an *alkoxide ion* is concentrated on the oxygen atom, but the negative charge on a *phenoxide ion* can be delocalized to the ortho and para ring positions through resonance.

Because *phenoxide ions* are stabilized in this way, the equilibrium for their formation is more favorable than that for *alkoxide ions*

16







## Preparation of Alcohols

### 1. Hydration of Alkenes

- a. Addition of water to a double bond in the presence of an *acid catalyst, H<sup>+</sup>*.




- b. The addition follows *Markovnikov's rule*.



- c. It is *not possible to prepare primary alcohols* except Ethanol.

### 2. Oxidation of Cycloalkenes


Alkenes react with alkaline potassium permanganate to form glycols.



19



## Preparation of Alcohols

### 3. Nucleophilic Substitution of Alkyl Halide



### 4. Reduction of Ketones, and Aldehydes

*Aldehydes and ketones are easily reduced to primary and secondary alcohols, respectively.*




20



## Preparation of Ethers


### 1) Dehydration of Alcohols

- It takes place in the presence of acid catalysts ( $\text{H}_2\text{SO}_4$ ,  $\text{H}_3\text{PO}_4$ ) (intermolecular reaction)




- Example;

The most important commercial ether is diethyl ether. It is prepared from ethanol and sulfuric acid.



- When ethyl alcohol is dehydrated by sulfuric acid at  $180^\circ\text{C}$ , the dominant product is ethylene.



23

## Preparation of Ethers

### 1) Dehydration of Alcohols

- To prepare ethyl ether

- Dissolve ethyl alcohol in sulfuric acid at ambient temperature.
- Heat the solution to  $140^\circ\text{C}$  while adding more alcohol.



24

## Preparation of Ethers

### 2) Williamson Synthesis

This method has two steps;

- 1) An alcohol is converted to its alkoxide by treatment with a reactive metal (sodium or potassium).



- 2) Displacement is carried out between the alkoxide and an alkyl halide.



- To obtain the best yields of mixed dialkyl ethers, we select a 1° rather than a 2° or 3° alkyl halide and react it with a sodium alkoxide
- To prepare an alkyl aryl ether, we must be careful not to pick a combination in which one of the reagents has a halogen directly attached to an aromatic ring.

25

## Preparation of Ethers

- Example 1; Preparation of *t*-butyl methyl ether,  $(\text{CH}_3)_3\text{C}-\text{O}-\text{CH}_3$ .


- In theory, this could be done by either of two reactions.

1. You could react sodium methoxide,  $\text{CH}_3\text{O}^- \text{Na}^+$ , with *t*-butyl chloride,  $(\text{CH}_3)_3\text{C}-\text{Cl}$ .

*This combination leads to dehydrohalogenation to an alkene, an elimination reaction.*

2. You could react sodium *t*-butoxide,  $(\text{CH}_3)_3\text{C}-\text{O}^- \text{Na}^+$ , with methyl chloride,  $\text{CH}_3\text{Cl}$ .

*This route gives the desired ether by substitution.*




26

## Preparation of Ethers

**Example 2;** Assume you need to synthesize methyl phenyl ether (anisole),  $\text{CH}_3\text{-O-C}_6\text{H}_5$ , by the Williamson method.

- In theory, you could obtain anisole in either of two ways.



27

## Reactions of Alcohols, Phenols and Ethers

- Alcohols undergo two kinds of reactions:
  - Those that involve the breaking of the oxygen-hydrogen bond (CO-H).
  - Those that involve the rupture of the carbon-oxygen bond (C-OH).
- Phenols do not participate in reactions where the C-OH bond is broken.
- Ethers are quite stable compounds.
  - The ether linkage does not react with bases, reducing agents, oxidizing agents, or active metals.
  - Ethers react only under strongly acidic conditions.

28

## Reactions of Alcohols

## Reactions of Alcohols, Phenols and Ethers

A) Those that involve the rupture of the carbon-oxygen bond (C-OH).


1) **Nucleophilic Substitution Reaction**; The Reaction of Alcohols with Hydrogen Halides: Alkyl Halides

*Alcohols react with hydrogen halides (HCl, HBr and HI) to give alkyl halides.*



2) **Dehydration of Alcohols: Formation of Alkenes**

*Alcohols can be dehydrated by heating them with strong acid.*



29


## Reactions of Alcohols

## Reactions of Alcohols, Phenols and Ethers



B) **Oxidation Reactions**

Alcohols with at least one hydrogen attached to the hydroxyl-bearing carbon can be oxidized to carbonyl compounds.

o **Primary alcohols** give **aldehydes**, which may be further oxidized to **carboxylic acids**.



▪ **Primary alcohols**, oxidation can be stopped at aldehyde stage by special reagents, such as "pyridinium chlorochromate (PCC)".




30

## Reactions of Alcohols


## Reactions of Alcohols, Phenols and Ethers

### B) Oxidation Reactions

- Primary alcohols yield aldehydes when treated with mild oxidizing agents such as hot metallic copper or  $\text{CrO}_3$  in pyridine.



- Primary alcohols; when treated with stronger oxidizing agents, such as chromic acid,  $\text{H}_2\text{Cr}_2\text{O}_7$ , or neutral potassium permanganate,  $\text{KMnO}_4$ , the intermediate aldehydes formed initially are oxidized further to carboxylic acids.




31

## Reactions of Alcohols

## Reactions of Alcohols, Phenols and Ethers

### B) Oxidation Reactions

- Secondary alcohols, when treated with any of the oxidizing agents mentioned previously, yield ketones.



- Tertiary alcohols, having no hydrogen atom on hydroxyl-bearing carbon, do not undergo oxidation.


32



## Reactions of Phenols

## Reactions of Alcohols, Phenols and Ethers

- Halogenation takes place *without catalyst*.




- The products depend on the solvent used.
  - In *aprotic solvents* (solvents that do not release protons) ( $\text{CCl}_4$ ,  $\text{CS}_2$ ) - bromination gives a mixture of *o*- and *p*-bromophenol.
  - In *protic solvents* (solvents that can release protons) ( $\text{H}_2\text{O}$ ) - halogenation gives a trisubstituted phenol is produced.

33


## Reactions of Ethers

## Reactions of Alcohols, Phenols and Ethers


- When **ethers** are heated in concentrated acid solutions, the ether linkage is broken.



- The acids most often used in this reaction are HI, HBr, and HCl.
- If an *excess of acid* is present, the alcohol initially produced is converted into an alkyl halide by the reaction.



For example,



34