



Fundamentals of Organic Chemistry

CHEM 109

For Students of Health Colleges

Credit hrs.: (2+1)

King Saud University

College of Science, Chemistry Department

CHEM 109

CHAPTER 4. ALCOHOLS, PHENOLS AND ETHERS

Learning Objectives



At the end of this chapter, students will able to:

- know the difference in structure between alcohols, phenols and ethers.
- Know the different classes of alcohols.
- Know how to name alcohols, phenols and ethers using IUPAC method.
- Recognize the basic properties (structure, physical and chemical properties) of alcohols, phenols and ethers .
- Recognize the effect of hydrogen bonds on their physical properties.
- Recognize the acidic properties of alcohols and phenols.
- know the different methods for the preparation of alcohols, phenols and ethers .
- Know the chemical reactions of these compounds.

Alcohols, Phenols and Ethers

3

- Alcohols, phenols and ethers may be viewed as organic derivatives of water.
- Alcohols and phenols have a common functional group, the hydroxyl group, -OH.




- Alcohols are compounds whose molecules have a hydroxyl group attached to a saturated carbon atom.
- Phenols are compounds that have a hydroxyl group attached directly to a benzene ring.
- Ethers are compounds whose molecules have an oxygen atom bonded to two carbon atom.

Classification of Alcohols and Ethers

4

- Alcohols are classified as primary (1°), secondary (2°), or tertiary (3°), depending on whether one, two, or three organic groups are connected to the hydroxyl-bearing carbon atom.




- Methyl alcohol, which is not strictly covered by this classification, is usually grouped with the primary alcohols.
- Ethers are classified as
 - Symmetrical ethers;
 - When the organic groups attached to the oxygen are identical.
 - Unsymmetrical ethers (mixed ethers);
 - When the organic groups attached to the oxygen are different.

Nomenclature of Alcohols

5

- The **common names** for the simplest alcohols consist of alkyl group attached to the hydroxyl function followed by the word alcohol: *Alkyl alcohol*.
- In the **IUPAC system**, alcohols are named according to the following rules.
 - Select the **longest continuous carbon chain that contains the -OH group**. Drop the **-e** ending of the parent alkane and replace it by the suffix **-ol**: *Alkanol*
 - When **isomers are possible**, the chain is numbered so as to give the functional group (-OH) the **lowest possible number**.




Nomenclature of Alcohols


6

- When **alkyl side chains or other groups are present**; they are named alphabetically and their positions are indicated by a number.

The position of the functional group (-OH) is always given the **lowest possible number** at the end of the name.



For cyclic alcohols, numbering always starts from the carbon bearing the -OH group.




Nomenclature of Alcohols

7

4. **With Unsaturated Alcohols;** If a molecule contains both an -OH group and a C=C or C-C triple bond, the -OH group takes preference before the double or triple bonds in getting the lower number.

The name should include (if possible) both the hydroxyl and the unsaturated groups, **even if this does not make the longest chain the parent hydrocarbon.**




Nomenclature of Alcohols

8

Alcohols with More Than One Hydroxyl Group

- Compounds with two adjacent alcohol groups are called **glycols**.
The most important example is ethylene glycol.
- Compounds with more than two hydroxyl groups are also known, and several, such as glycerol and sorbitol, are important commercial chemicals.




- Ethylene glycol is used as the “permanent” antifreeze in automobile radiators and as a raw material in the manufacture of Dacron.
- Ethylene glycol is completely miscible with water.
- Glycerol is a syrupy, colorless, water-soluble, high-boiling liquid with a distinctly sweet taste. Its soothing qualities make it useful in shaving and toilet soaps and in cough drops and syrups.


Nomenclature of Phenols

9

- Phenols are usually named as derivatives of the parent compounds.



- The hydroxyl group is named as a substituent when it occurs in the same molecule with carboxylic acid, aldehyde, or ketone functionalities, which have **priority in naming**.




Nomenclature of Ethers

10


Common Names

Ethers are usually named by giving the name of each alkyl or aryl group, in alphabetical order, followed by the word *ether*.



IUPAC system

- For ethers with more complex structures, it may be necessary to **name the -OR group** as an **alkoxy** group.
- In the IUPAC system, the smaller alkoxy group is named as a substituent.



Physical Properties of Alcohols and Ethers

11


Physical State

- The simplest **alcohol**, methanol, is a liquid at room temperature. In contrast, alkanes from methane to butane are gases.
- Phenol** is a colorless, crystalline, and low-melting solid and other phenols also are solids, .
- Ethers** are colorless compounds with characteristic, relatively pleasant odors.

Boiling Points

- Ethers** have lower boiling points (bps) than **alcohols** with an equal number of carbon atoms.
- Ether** has nearly the same b.p. as the corresponding hydrocarbon in which a -CH₂- group replaces the ether's oxygen.

Because of their structures (no O-H bonds), ether molecules cannot form hydrogen bonds with one another.




Compound	Formula	bp	mol wt	Water solubility (g/100 mL, 20°C)
1-butanol	CH ₃ CH ₂ CH ₂ CH ₂ OH	118°C	74	7.9
diethyl ether	CH ₃ CH ₂ -O-CH ₂ CH ₃	35°C	74	7.5
pentane	CH ₃ CH ₂ -CH ₂ -CH ₂ CH ₃	36°C	72	0.03

Physical Properties of Alcohols and Ethers

12

Boiling Points

- Series of normal alcohols;** The boiling points increase with increasing molecular weights.
- A comparison of boiling points among **isomeric alcohols;** The boiling points decrease as the number of alkyl branches from the carbinol group increases.



- Phenol** and most other phenols have high boiling points.


Physical Properties of Alcohols and Ethers

13

Solubility

- The lower **alcohols** are completely miscible with water.
- As the number of carbons in the alcohol increases, the solubility in water decreases.
- Low-molecular-weight ethers, such as dimethyl ether, are quite soluble in water.

Ether molecules can form hydrogen bonds to water.



Structure	Name	Mol.wt.	Bp (°C)	Solubility in H ₂ O At 20 °C
CH ₃ CH ₂ CH ₃	propane	44	-42	insoluble
CH ₃ OCH ₃	methyl ether	46	-24	soluble
CH ₃ CH ₂ OH	ethanol	46	78	soluble
CH ₃ CH ₂ CH ₂ CH ₃	n-butane	58	-0.5	insoluble
CH ₃ CH ₂ OCH ₂ CH ₃	ethyl methyl ether	60	8	soluble
CH ₃ CH ₂ CH ₂ OH	1-propanol	60	97	soluble
CH ₃ (CH ₂) ₃ CH ₃	n-pentane	72	35	insoluble
CH ₃ CH ₂ OCH ₂ CH ₃	ethyl ether	74	36	7.5 g/100 g
CH ₃ (CH ₂) ₂ CH ₂ OH	1-butanol	74	118	7.9 g/100 g
CH ₃ (CH ₂) ₅ CH ₃	n-heptane	100	98	insoluble
CH ₃ (CH ₂) ₂ O(CH ₂) ₂ CH ₃	n-propyl ether	102	91	0.2 g/100 g
CH ₃ (CH ₂) ₄ CH ₂ OH	1-hexanol	102	157	0.6 g/100 g

- Phenol** and most other phenols are slightly soluble in water .

Hydrogen Bonding in Alcohols and Ethers


14

- The **boiling points** (bp's) of alcohols are much higher than those of ethers or hydrocarbons with similar molecular weights.

	CH ₃ CH ₂ OH	CH ₃ OCH ₃	CH ₃ CH ₂ CH ₃
mol wt	46	46	44
bp	+78.5°C	-24°C	-42°C

Why? Because alcohols form **hydrogen bonds** with one another.

The O-H bond is polarized by the high electronegativity of the oxygen atom and places a partial positive charge on the hydrogen atom and a partial negative charge on the oxygen atom.




- Consequently, **alcohols** have relatively high boiling points because they must supply enough heat to break the hydrogen bonds before each molecule.
- Hydrogen bonds are weaker than ordinary covalent bonds.

Hydrogen Bonding in Alcohols and Ethers

15

- The lower molecular-weight alcohols and ethers can form H-bond with water molecules.
- This accounts for the complete miscibility of the lower alcohols and ethers with water.



- However, as the organic chain lengthens and the alcohol becomes relatively more hydrocarbon like, its water solubility decreases.

Table 7.1 Boiling Point and Water Solubility of Some Alcohols



Name	Formula	bp, °C	Solubility in H ₂ O g/100 g at 20°C
methanol	CH ₃ OH	65	completely miscible
ethanol	CH ₃ CH ₂ OH	78.5	completely miscible
1-propanol	CH ₃ CH ₂ CH ₂ OH	97	completely miscible
1-butanol	CH ₃ CH ₂ CH ₂ CH ₂ OH	117.7	7.9
1-pentanol	CH ₃ CH ₂ CH ₂ CH ₂ CH ₂ OH	137.9	2.7
1-hexanol	CH ₃ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ OH	155.8	0.59

The Acidity of Alcohols and Phenols

16

- Like water, alcohols and phenols are weak acids.


The hydroxyl group can act as a proton donor, and dissociation occurs in a manner similar to that for water



The Acidity of Alcohols and Phenols

17

- **Phenols are stronger acids than alcohols** mainly because the corresponding phenoxide ions are stabilized by resonance.




The negative charge of an **alkoxide ion** is concentrated on the oxygen atom, but the negative charge on a phenoxide ion can be delocalized to the ortho and para ring positions through resonance.

Because **phenoxide ions** are stabilized in this way, the equilibrium for their formation is more favorable than that for alkoxide ions

The Acidity of Alcohols and Phenols

18

- All **electron-withdrawing groups increase acidity** by stabilizing the conjugate base. **Electron-donating groups decrease acidity** because they destabilize the conjugate base.



Preparation of Alcohols


21

○ Nucleophilic Substitution of Alkyl Halide



○ Reduction of Ketones, and Aldehydes

Aldehydes and ketones are easily reduced to primary and secondary alcohols, respectively.




Preparation of Alcohols

22

○ Addition of Grignard's Reagent to Aldehydes and Ketones

General equations




Preparation of Phenols

23

○ The Alkali Fusion of Sulfonates

The alkali fusion of sulfonates involves the following steps;

1. **Sulfonation** of an aromatic ring.
2. **Melting (fusion)** of the aromatic sulfonic acid with sodium hydroxide to give a **phenoxide salt**.
3. **Acidification** of the phenoxide with HCl to produce the **phenol**.



Uses of Alcohols and Phenols


- Drinks – The “alcohol” in alcoholic drinks is simply ethanol.
- As a fuel – Ethanol burns to give carbon dioxide and water and can be used as a fuel in its own right, or in mixtures with petrol (gasoline).
- As a solvent – Ethanol is widely used as a solvent. It is relatively safe, and can be used to dissolve many organic compounds which are insoluble in water. It is used, for example, in many perfumes and cosmetics.
- Isopropanol; rubbing alcohol, rapid evaporation, Antiseptic, more toxic than ethanol, but induces vomiting and used for manufacture of acetone.
- Phenol is used for resins, and pharmaceuticals.

Preparation of Ethers

25


1) Dehydration of Alcohols

- It takes place in the presence of acid catalysts (H_2SO_4 , H_3PO_4) (intermolecular reaction)




- Example;**

The most important commercial ether is diethyl ether. It is prepared from ethanol and sulfuric acid.



- When ethyl alcohol is dehydrated by **sulfuric acid at 180°C** , the dominant product is **ethylene**.



Preparation of Ethers

26

2) Williamson Synthesis

- This method has two steps;
 - An alcohol is converted to its alkoxide by treatment with a reactive metal (sodium or potassium).



- Displacement is carried out between the alkoxide and an alkyl halide.



- To obtain the best yields of mixed dialkyl ethers, we select a 1° rather than a 2° or 3° alkyl halide and react it with a sodium alkoxide
- To prepare an alkyl aryl ether, we must be careful not to pick a combination in which one of the reagents has a halogen directly attached to an aromatic ring.

Preparation of Ethers

27

○ Example 1; Preparation of *t*-butyl methyl ether, $(\text{CH}_3)_3\text{C-O-CH}_3$.


➤ In theory, this could be done by either of two reactions.

1. You could react sodium methoxide, $\text{CH}_3\text{O}^-\text{Na}^+$, with *t*-butyl chloride, $(\text{CH}_3)_3\text{C-Cl}$.

This combination leads to dehydrohalogenation to an alkene, an elimination reaction.

2. You could react sodium *t*-butoxide, $(\text{CH}_3)_3\text{C-O}^-\text{Na}^+$, with methyl chloride, CH_3Cl .

This route gives the desired ether by substitution.




Preparation of Ethers

28

Example 2; Assume you need to synthesize methyl phenyl ether (anisole), $\text{CH}_3\text{-O-C}_6\text{H}_5$, by the Williamson method.

➤ In theory, you could obtain anisole in either of two ways.



Reactions of Alcohols

31

A) Those that involve the rupture of the carbon-oxygen bond (C-OH).

1) Nucleophilic Substitution Reaction; The Reaction of Alcohols with Hydrogen Halides:


Alkyl Halides

Alcohols react with hydrogen halides (HCl, HBr and HI) to give alkyl halides.



2) Dehydration of Alcohols: Formation of Alkenes

Alcohols can be dehydrated by heating them with strong acid.




Reactions of Alcohols

32



B) Oxidation Reactions

Alcohols with at least one hydrogen attached to the hydroxyl-bearing carbon can be oxidized to carbonyl compounds.

- **Primary alcohols** give **aldehydes**, which may be further oxidized to **carboxylic acids**.



- **Primary alcohols**, oxidation can be stopped at aldehyde stage by special reagents, such as "pyridinium chlorochromate (PCC)".




Reactions of Alcohols


33

B) Oxidation Reactions

- Primary alcohols yield aldehydes when treated with *mild oxidizing agents* such as hot metallic copper or CrO_3 in pyridine.



- Primary alcohols; when treated with *stronger oxidizing agents*, such as chromic acid, $\text{H}_2\text{Cr}_2\text{O}_7$, or neutral potassium permanganate, KMnO_4 , the intermediate aldehydes formed initially are oxidized further to carboxylic acids.




Reactions of Alcohols

34

B) Oxidation Reactions

- Secondary alcohols, when treated with any of the oxidizing agents mentioned previously, yield ketones.




- Tertiary alcohols, having no hydrogen atom on hydroxyl-bearing carbon, do not undergo oxidation.

Reactions of Phenols

35

- Halogenation takes place *without catalyst*.




- The products depend on the solvent used.
 - In *aprotic solvents* (solvents that do not release protons) (CCl_4 , CS_2)-bromination gives a mixture of *o*- and *p*-bromophenol.
 - In *protic solvents* (solvents that can release protons) (H_2O)-halogenation gives a trisubstituted phenol is produced.

Reactions of Ethers

36

- Cleavage of Ethers by Hot Concentrated Acids


- When **ethers** are heated in concentrated acid solutions, the ether linkage is broken.



- The acids most often used in this reaction are HI, HBr, and HCl.
- If an excess of acid is present, the alcohol initially produced is converted into an alkyl halide by the reaction.



For example,



Uses of Ethers



- Ether is used as a mild anesthetic and as a solvent in industries
- It is used as an antiseptic to prevent infection when an injection is administered.
- Dimethyl ether is used as refrigerant and as solvent at low temperature.
- Diethyl ether is a common ingredient as an aesthesia in surgery.
- Diethyl ether is common solvent for oils, gums, resins etc.
- We use phenyl ether as a heat transfer medium because of its high boiling point.