

College of Science.
Department of Statistics & Operations
Research

كلية العلوم
قسم الإحصاء وبحوث العمليات

Final Exam
Academic Year 1442-1443 Hijri- First Semester

| Exam Information معلومات الامتحان | | | |
|-----------------------------------|---|------------|-------------------|
| Course name | Modeling and Simulation النمذجة والمحاكاة | | اسم المقرر |
| Course Code | OPER 441 441 بحث | | رمز المقرر |
| Exam Date | 2021-12-23 | 1443-05-19 | تاريخ الامتحان |
| Exam Time | 08: 00 AM | | وقت الامتحان |
| Exam Duration | 3 hours | ثلاث ساعات | مدة الامتحان |
| Classroom No. | | | رقم قاعة الاختبار |
| Instructor Name | | | اسم استاذ المقرر |

| Student Information معلومات الطالب | | |
|------------------------------------|------------------------|----------------|
| Student's Name | هيفاء محمد حمد ال جياد | اسم الطالب |
| ID number | 438201897 | الرقم الجامعي |
| Section No. | | رقم الشعبة |
| Serial Number | | الرقم التسلسلي |
| General Instructions: | | تعليمات عامة: |

- Your Exam consists of PAGES (except this paper) عدد صفحات الامتحان صفحة. (باستثناء هذه الورقة)
- Keep your mobile and smart watch out of the classroom. يجب ابقاء الهواتف والساعات الذكية خارج قاعة الامتحان.
-

هذا الجزء خاص بأستاذ المادة

This section is ONLY for instructor

| # | Course Learning Outcomes (CLOs) | Related Question (s) | Points | Final Score |
|---|--|----------------------|--------|-------------|
| 1 | To know the basics of pseudo random generation and apply different methods of random generation techniques | | | |
| 2 | Chose and fit theoretical distribution on collected data | | | |
| 3 | Generate random variates from different probability functions and directly from collected data | | | |
| 4 | build simple simulation models of real-life problems | | | |
| 5 | Define and compute performance measures from simulation models | | | |
| 6 | Recognize and analyze simple models and its main elements for simulation | | | |
| 7 | Understanding how to use computer software (ECXEL) for simulation models | | | |
| 8 | use appropriate statistical techniques to analyze and evaluate outputs of simulation models | | | |

| | Q. #1 | Q. #2 | Q. #3 | Q. #4 | Q. #5 | Q. #6 | Total |
|-------|-------|-------|-------|-------|-------|-------|-------|
| | 23 | 20 | 13 | 15 | 24 | 20 | 115 |
| Score | 10 | 12 | 8 | 15 | 20 | 17 | 82 |

10
23

Question # 1:

Consider a car insurance company collected the following report about their clients:

| Age | Male | | | | Female | | | |
|--------------|--------------------|----------------------------|------------------------|------------------------|--------------------|----------------------------|------------------------|------------------------|
| | Total # of clients | Total #accidents per group | Min Claim (per client) | Max Claim (per client) | Total # of clients | Total #accidents per group | Min Claim (per client) | Max Claim (per client) |
| 18 - 28 | 113 | 284 | 350 | 9000 | 312 | 435 | 550 | 7500 |
| 28 - 38 | 258 | 314 | 950 | 13000 | 271 | 361 | 1050 | 11300 |
| 38 - 48 | 546 | 395 | 1300 | 6500 | 687 | 302 | 1450 | 9300 |
| 48 - 58 | 420 | 150 | 2000 | 11000 | 348 | 129 | 2800 | 11800 |
| 58 - 68 | 302 | 145 | 4000 | 20000 | 354 | 159 | 3700 | 32000 |
| 68 - 90 | 93 | 23 | 15000 | 80000 | 12 | 43 | 21000 | 65000 |
| Total | 1732 | 1311 | | | 1984 | 1429 | | |

Answer the following questions:

- How to simulate the gender type of the client from the table? Write the algorithm.

$$F_x(x) = \sum_{i=1}^n I_{[0, x]}(F(x_i))$$
 Sum of uniform dist.

$$f_x(x) = \frac{1}{b-a+1}$$

$$F_x(x) = F_{Male} + F_{Female}$$
 use convolution with parameter and Inverse transform
 $a = \text{min claim}$
 $b = \text{max claim}$

- Give a simulated gender type of the 4 clients using the following uniform values:

| | | | | |
|-----------------|--------|--------|--------|--------|
| $u \sim U[0,1]$ | 0.1466 | 0.7061 | 0.8585 | 0.4944 |
| | | | | |

- Given that you want to simulate the age of a male client between 28 and 38, what you will do assuming uniform distribution?

$$f(x) = \frac{1}{b-a}$$

$$F(x) = \frac{x-a}{b-a} = u \Rightarrow F^{-1}(u) = u(b-a) + a$$

$$F^{-1}(u) = u(b-a+1) + a$$

$$= u(38-28+1) + 28$$

- How to simulate the exact age of the client from the table assuming uniform distribution? Write the algorithm.

$$F^{-1}(u) = u(b-a) + a$$
 age

cont.

5. Give a simulated exact age (round to 1 decimal digit ##.#) for 4 clients using the following uniform values assuming uniform distribution in each category:

| | | | | |
|-----------------|--------|--------|--------|--------|
| $u \sim U[0,1]$ | 0.5915 | 0.1998 | 0.9780 | 0.6281 |
| | 23.9 | | | |
| | | | | |

6. How to generate the amount of the claim for each client from the table assuming integer uniform? Write the general algorithm.

$$F^{-1}(u) = u(b-a) + a$$

age

7. Give a simulated exact amount of the claim for 4 clients using the following uniform values assuming discrete uniform distribution in each category:

| | | | | |
|-----------------|---------|--------|--------|--------|
| $u \sim U[0,1]$ | 0.3887 | 0.0739 | 0.9482 | 0.7715 |
| | 3712.64 | | | |
| | | | | |

discrete dist.

$$F^{-1}(u) = u(b-a+1) + a$$

claim

?

$$\frac{1}{5} - \frac{x}{20} = u \rightarrow \frac{1}{5} - u = \frac{x}{20}$$

$$x^2 = 4 - 20u \rightarrow x = \sqrt{4 - 20u}$$

Question # 2:

Consider an investment with a monthly return on investment. The monthly percentage of return on investment is a random variable $X\%$ given by the following probability function:

$$f(x) = \begin{cases} \frac{|x|}{10}, & -2 \leq x \leq 4 \\ 0, & \text{otherwise.} \end{cases} \quad \text{where} \quad f(x) = \begin{cases} \frac{-x}{10}; & -2 \leq x \leq 0 \\ \frac{x}{10}; & 0 \leq x \leq 4 \end{cases}$$

Answer the following:

1) Write the inverse algorithm for generating the monthly percentage of return on investment.

$$F(x) = \int_{-2}^x \frac{-x}{10} dx = \left[-\frac{x^2}{20} \right]_{-2}^x = -\frac{x^2}{20} + \frac{(-2)^2}{20} = \frac{1}{5} - \frac{x^2}{20} \quad \text{for } -2 \leq x \leq 0$$

$$F(x) = \int_0^x \frac{x}{10} dx = \left[\frac{x^2}{20} \right]_0^x = \frac{x^2}{20} \quad \text{for } 0 \leq x \leq 4$$

$$F(x) = \begin{cases} \frac{1}{5} - \frac{x^2}{20} & -2 \leq x \leq 0 \\ \frac{x^2}{20} & 0 \leq x \leq 4 \end{cases}$$

$$F^{-1}(u) = \begin{cases} \sqrt{4 - 20u} & \text{for } 0 \leq u \leq \frac{1}{5} \\ \sqrt{20u} & \text{for } \frac{1}{5} < u \leq 1 \end{cases}$$

2) Using simulation and the following $U[0,1]$ numbers, evaluate the results of the investment for 6 months with initial budget of 100,000 SR.

| Mon. | u | X(n) | AT(n) |
|------|-------|---------|---------|
| 1 | 0.032 | 1.833 | 1.833 |
| 2 | 0.138 | 1.11355 | 2.94655 |
| 3 | 0.623 | 3.5248 | 6.47 |
| 4 | 0.691 | 3.717 | ? |
| 5 | 0.006 | 0.346 | |
| 6 | | | |

→ stop here

3) From simulation, compute the average and standard deviation of the monthly percentage of return on investment.

$$\text{Average} = \int x f(x)$$

$$\text{Average} = 2$$

4) From simulation compute the probability that the company will have profit more than 10000 SR per month.

$$P(x > 10,000) =$$

5) From simulation output, what is the probability of losing.

$$X < 0$$

Patient Arrive ~ Poisson (8)
 health Cond. (serious, stable; Critical)

8/3

Question #3:

Patients arrive to a hospital's emergency room according to a Poisson process with rate 8 patients per hour. Patients come in three different health conditions. The patients are categorized according to their condition as critical, serious, or stable. In the past year, statistics show that:

- i. 10% of the emergency room patients were critical and take random amount of treatment of Erlang with parameters $\alpha = 2$ and $\lambda = 0.5$ per hour
- ii. 30% of the emergency room patients were serious; and take random amount of treatment of Exponential with average time of 2 hours.
- iii. the rest of the emergency room patients were stable; and take random amount of treatment of integer uniform between 15 min and 30 min.

0/60

After treatment at the ER, statistics show that: 40% of the critical patients died; 10% of the serious patients died; and 1% of the stable patients died.

Answer the following with **True or False**:

$$\begin{array}{l|l} T(n) & PAT(n) \\ \hline T(n) & T(n) = PAT(n) \\ T(n+1) & T(n) + T(n+1) = PAT(n+1) \end{array}$$

| | | |
|---|-----|--|
| F | 1. | To simulate the ER system, we have to simulate 3 independent random processes only: Patients Arrival Time PAT(n), Patient Treatment Time PTT(n) and Patient Exit Conditions PEC(n). + Patient Come Condition |
| T | 2. | To simulate the patient's arrival to ER system, we need to use the inverse transform for exponential distribution with mean value = 1/8. $E(X) = \frac{1}{\lambda} = 8 \rightarrow \lambda = 1/8$ |
| T | 3. | The time between arrivals T(n) is simulated by the function: $-8 \ln(1-u)$ where $u \sim U[0,1]$ |
| F | 4. | The arrival time of the patient (n) = PAT(n) is equal to PAT(n-1) - T(n). Where T(n) time between arrivals from simulation. $\hookrightarrow = PAT(n-1) + T(n)$ |
| T | 5. | The condition of the patient CP(n) can be generated as a random process with discrete uniform with values 1, 2, 3. |
| T | 6. | To simulate the condition of the patient CP(n), get $u_1 \sim U[0,1]$: if $0 \leq u_1 \leq 0.1$ then CP(n)=1 (Critical). If $0.1 < u \leq 0.3$ then CP(n)=2 (serious). If $0.3 < u \leq 1$ then CP(n)=3 (stable) |
| T | 7. | Patient Treatment Time PTT(n) is a random variable depends on the condition of the patient CP(n). |
| F | 8. | If the condition of the patient CP(n)=2 (serious) then PTT(n)~DU (15, 30 min); using discrete inverse method: take $u_2 \sim U[0,1]$ and PTT(n) = 15 + int[(1+30 - 15) u ₂] Stable |
| F | 9. | If the condition of the patient CP(n) = 1 (critical) then we generate the Patient Treatment Time PTT(n) ~ Er($\alpha = 2$ and $\lambda = 0.5$ per hour) by using inverse transform method directly. \hookrightarrow convolution because erlang is sum of exponential. |
| T | 10. | It is always possible to use acceptance/rejection method to generate the Patient Treatment Time PTT(n) ~ Er($\alpha = 2$ and $\lambda = 0.5$ per hour). |
| T | 11. | Patient Exit Conditions PEC(n) is also a random variable depends on the condition of the patient CP(n). |
| T | 12. | If condition of the patient CP(n) = 2 (serious), then Patient Exit Conditions PEC(n) ~ Bernoulli (p = 0.1) with $w \sim U[0,1]$ If $w < 0.1$ then PEC(n) = Lived, Else PEC(n) = Died |
| F | 13. | If condition of the patient CP(n) = 3 (stable), then Patient Exit Conditions PEC(n) ~ Bernoulli (p = 0.01) with $w \sim U[0,1]$ If $w < 0.99$ then PEC(n) = Lived, Else PEC(n) = Died |
| T | 14. | The patient Departure time DT(n) is equal to patient arrival time PAT(n) + Patient Treatment Time PTT(n). |

15
15

Question #4:

An insurance policy pays for the insured 1000 SR per day spent admitted in a hospital for up to three days. If the insured spend more than three days, the insurance company pays 500 SR per day for each extra day of hospitalization thereafter. The number of days that an insured customer spends in the hospital, X , is a discrete random variable with probability function:

$$P[X = k] = \begin{cases} \frac{6-k}{15}, & k = 1, 2, 3, 4, 5 \\ 0, & \text{otherwise.} \end{cases}$$

Answer the following:

- 1) Write the inverse algorithm for generating number of days for the insured clients
- 2) Write the simulation algorithm and use the following $U[0,1]$ numbers, give the results for the 9 insured clients.

| u | Days | Cost (n) | u | Days | Cost (n) | u | Days | Cost (n) |
|-------|------|----------|-------|------|----------|-------|------|----------|
| 0.032 | 1 | 1000 | 0.691 | 3 | 3000 | 0.823 | 4 | 3500 |
| 0.138 | 1 | 1000 | 0.006 | 1 | 1000 | 0.776 | 3 | 3000 |
| 0.623 | 3 | 3000 | 0.413 | 2 | 2000 | 0.306 | 1 | 1000 |

- 3) Compute the average and standard deviation of amount that the insurance company will pay per claim.
- 4) Compute the average and standard deviation of number of days that a client spends in the hospital.
- 5) From simulation compute the probability that the company will pay more than 2000 SR.

(1) $F(x) = \frac{1}{b-a+1} = u$

4) $E(x) = \sum k P(k) = \frac{7}{3}$
 $Var(x) = E[x^2] - (E(x))^2 = \frac{14}{9}$
 Standard deviation = $\sqrt{Var(x)} = 1.247$

| k | 1 | 2 | 3 | 4 | 5 |
|---------------|--------|--------|---------|---------|--------|
| $P(x)$ | $5/15$ | $4/15$ | $3/15$ | $2/15$ | $1/15$ |
| $P(x \leq x)$ | $5/15$ | $9/15$ | $12/15$ | $14/15$ | 1 |
| | 0.33 | 0.6 | 0.8 | 0.93 | |

5) For the company to pay more than 2000 $X \geq 3$

$P(X \geq 3) = P(X=3) + P(X=4) + P(X=5)$

$$F^{-1}(x) = \begin{cases} 1 & 0 < u < \frac{5}{15} \\ 2 & \frac{5}{15} < u < \frac{9}{15} \\ 3 & \frac{9}{15} < u < \frac{12}{15} \\ 4 & \frac{12}{15} < u < \frac{14}{15} \\ 5 & \frac{14}{15} < u < 1 \end{cases}$$

(3) Average = $\frac{\sum \text{Cost}}{\text{total time}} = \frac{18500}{9} = 2055.56$

$Var(x) = \frac{\sum (X_i - \bar{X})^2}{n} = \frac{1}{9} [4(1000 - 2055.56)^2 + 3(3000 - 2055.56)^2 + (2000 - 2055.56)^2 + (3500 - 2055.56)^2]$
 $= 1024691.358$

Standard deviation = $\sqrt{Var(x)}$ 6 out of 8
 $= 1012.27$

20
24

Question #5: Consider the following are functions in EXCEL.

| | | | | | |
|--------------|--------------------|--------------|---------------------|--------------|---------------------|
| A | RANDBETWEEN(a,b) | E | GAMMA.DIST(a,b,c,d) | J | RAND() |
| B | NORM.DIST(a,b,c,d) | F | DATA TABLE | J | BINOM.INV(a,b,c) |
| C | GAMMA.INV(a,b,c,d) | G | BINOM.DIST(a,b,c) | K | CONFIDENCE.T(a,b,c) |
| D | VLOOKUP(a,b,c) | H | NORM.INV(a,b,c) | | |

Read each statement and assign the letter of EXCEL function from the table to the statement.

| The Statement | Excel Function |
|---|----------------|
| 1. Used to compute the PDF function of the Normal distribution with $\mu=a$ and $\sigma=b$ at the value a | B ✓ |
| 2. Used to generates random values from relative frequency table. | I ✗ |
| 3. Used generate integer random values with equal distributions between (a) and (b) including the (a) and (b) | A |
| 4. Used to generate random values from Exponential distribution | C |
| 5. Used to compute the upper limit of the confidence interval from sample | K |
| 6. Used generate integer random values with equal distributions between (a) and (b) without the (a) and (b) | A |
| 7. Used to generate random values from Erlang distribution | C |
| 8. Used to compute the PDF function of the Gamma distribution at the value a | E |
| 9. Used to compute the CDF function of the Normal distribution with $\mu=a$ and $\sigma=b$ at the value a | D ✗ |
| 10. Used to Generates random values from Gamma distribution | C |
| 11. Used to generate random values from of integer values (a) to (b) with different probabilities | A/I ✗ |
| 12. Used to compute the CDF function of the Gamma distribution at the value a | E |
| 13. Used to generate continuous random values between a and b with uniform distribution | I |
| 14. Used to compute the PMF function of the Binomial distribution with value a | G |
| 15. Used to run the simulation model for many times and record the measures from every run | D ✗ |
| 16. Used to compute the CDF function of the Binomial distribution with at the value a | G |
| 17. Used to compute the CDF function of the Exponential distribution at the value a | E |
| 18. Used to generate random values from standard deviation | H |
| 19. Used to compute the CDF function of the Erlang distribution at the value a | E |
| 20. Used to Generates random values from normal distribution with mean b and standard deviation c | H |
| 21. Used to generate continuous random values between 0 and 1 with uniform distribution | I |
| 22. Used to compute $Z_{\alpha/2}$ | F ✗ |
| 23. Used to compute the lower limit of the confidence interval from sample | K |
| 24. Used to generate random values from binomial distribution | J |
| 25. Used to compute the half width of the confidence interval from sample | K |

Time between arrivals \sim Erlang ($r=2, \lambda=0.5$)
 repair time \sim Weibull ($\alpha=2, \beta=3$)

Question #6:

Customer arrive to a car repair station for maintenance according to a random process with time between arrivals (in hours) follows Erlang distribution with parameters $r=2$ and $\lambda = 0.5$. The arriving cars come to a single repairman to perform the maintenance. It is assumed that the repair time (in hours) follows a random time that follows a Weibull distribution with parameters $\alpha = 2$ and $\beta=3$. The cars are repaired according to the order of their arrival. The repair station has a space that can hold any number of cars. The following data was collected.

17
20

الوقت
المغيب

1. Define all the variables and random process and the logic of the system $TBC(n), AT(n)$
2. Draw the flow chart for the system for simulation of 10 cars. $ST(n), WT(n), IT(n)$
3. Complete the results of simulation in the table below? Use uniform values as needed starting from u_1 $ET(n)$

| car# | time btwn cars | Arrival Time | Service time | start time | Wait? (0/1) | Waiting Time | Exit time | Idle Time |
|-----------|----------------|--------------|--------------|------------|-------------|--------------|-----------|-----------|
| u_1 | 0.164 | 0.092 | 0.588 | 0.260 | 0.608 | 0.094 | 0.722 | 0.260 |
| u_2 | 0.938 | 0.412 | 0.879 | 0.138 | 0.018 | 0.371 | 0.247 | 0.138 |
| 1 | 5.919 | 5.919 | 1.92 | 5.919 | 0 | 0 | 7.839 | 1.92 |
| u_1 | 0.129 | 0.874 | 0.841 | 0.308 | 0.165 | 0.136 | 0.368 | 0.308 |
| u_2 | 0.355 | 0.992 | 0.690 | 0.040 | 0.661 | 0.904 | 0.308 | 0.040 |
| 2 | 1.15 | 7.069 | 2.45 | 7.839 | 1 | 0.77 | 10.289 | 2.45 |
| u_1 | 0.130 | 0.371 | 0.220 | 0.090 | 0.927 | 0.680 | 0.032 | 0.090 |
| u_2 | 0.012 | 0.563 | 0.668 | 0.262 | 0.416 | 0.607 | 0.737 | 0.262 |
| 3 | 0.3026 | 7.371 | 1.25 | 10.289 | 1 | 2.918 | 11.539 | 1.25 |
| u_1 | 0.999 | 0.176 | 0.429 | 0.131 | 0.830 | 0.371 | 0.362 | 0.131 |
| u_2 | 0.020 | 0.071 | 0.079 | 0.221 | 0.891 | 0.190 | 0.067 | 0.221 |
| 4 | 13.86 | 21.23 | 1.64 | 21.23 | 0 | 0 | 22.87 | 1.64 |
| u_1 | 0.840 | 0.021 | 0.492 | 0.843 | 0.428 | 0.062 | 0.112 | 0.843 |
| u_2 | 0.128 | 0.137 | 0.340 | 0.038 | 0.446 | 0.848 | 0.944 | 0.038 |
| 5 | 3.939 | 25.17 | 1.75 | 25.17 | 0 | 0 | 26.92 | 1.75 |
| u_1 | 0.584 | 0.211 | 0.855 | 0.344 | 0.844 | 0.296 | 0.311 | 0.344 |
| u_2 | 0.577 | 0.647 | 0.659 | 0.850 | 0.977 | 0.558 | 0.646 | 0.850 |
| 6 | 3.474 | 28.64 | 2.49 | 28.64 | 0 | 0 | 31.13 | 2.49 |
| u_1 | 0.684 | 0.815 | 0.407 | 0.053 | 0.072 | 0.969 | 0.442 | 0.053 |
| u_2 | 0.305 | 0.682 | 0.212 | 0.045 | 0.783 | 0.014 | 0.076 | 0.045 |
| 7 | 3.03 | 31.67 | 1.61 | 31.67 | 0 | 0 | 33.28 | 1.61 |
| u_1 | 0.318 | 0.117 | 0.843 | 0.614 | 0.217 | 0.009 | 0.532 | 0.614 |
| u_2 | 0.716 | 0.408 | 0.439 | 0.874 | 0.429 | 0.332 | 0.287 | 0.874 |
| 8 | 3.28 | 34.95 | 2.45 | 34.95 | 0 | 0 | 37.4 | 2.45 |
| u_1 | 0.108 | 0.034 | 0.028 | 0.328 | 0.325 | 0.243 | 0.825 | 0.328 |
| u_2 | 0.532 | 0.833 | 0.628 | 0.280 | 0.144 | 0.157 | 0.131 | 0.280 |
| 9 | 1.74 | 36.69 | 0.61 | 37.4 | 1 | 0.71 | 38.01 | 0.61 |
| u_1 | 0.918 | 0.115 | 0.798 | 0.260 | 0.960 | 0.750 | 0.268 | 0.260 |
| u_2 | 0.835 | 0.991 | 0.569 | 0.823 | 0.056 | 0.603 | 0.848 | 0.823 |
| 10 | 8.605 | 45.29 | 2.338 | 45.29 | 0 | 0 | 47.62 | 2.33 |

ok

4. From the simulation run, what is the percentage of customers who wait in line? =
5. What is the probability that the repairman is IDLE during the simulation Time?
6. What is the average waiting time in line? $\Rightarrow \frac{\text{waiting time}}{\text{Total time}} =$

* Customer Wait

* Total Costumer

$= \frac{3}{16}$

8 out of 8

$= \frac{0.77 + 2.918 + 0.71}{45.29}$