

King Saud University
Faculty of Sciences
Department of Mathematics

Second Examination

Math 106

Semester I

1437-1438

Time: 1H30

Exercise 1 : (3+3+3)

a) Find the following limit

$$\lim_{x \rightarrow 0^+} (\tan x)^{\tan x}.$$

b) Evaluate the following integral $\int \sinh^{-1} x dx$.

c) Evaluate the following integral $\int \tan^5 x \sec x dx$.

Exercise 2 : (2+3+3)

a) Find $\int \frac{1}{\sqrt{-x^2 + 6x - 5}} dx$.

b) Evaluate $\int \frac{\sqrt{9 - x^2}}{x^2} dx$.

c) Compute the integral $\int \frac{2x^2 - x + 2}{x(x^2 + 2)} dx$.

Exercise 3 : (3+2+3)

a) Find $\int \frac{dx}{1 - \sin x}$.

b) Does the integral $\int_{-\infty}^0 x e^x dx$ converges? If it converges find its value.

c) Sketch the region bounded by the curves $y = x^2 - 8$, $y = -x^2$ and find its area.

106Midterm2 Solutions (Sem1-37/38)

Question1 a) $\ln y = \tan x \ln \tan x$ (0,5) + 2 + (0,5)

$$\lim_{x \rightarrow 0^+} \ln y = \lim_{x \rightarrow 0^+} \frac{\ln \tan x}{\frac{1}{\tan x}} = \lim_{x \rightarrow 0^+} \frac{\sec^2 x \tan^2 x}{\tan x (-\sec^2 x)} = - \lim_{x \rightarrow 0^+} \tan x = 0$$

$$\text{So } \lim_{x \rightarrow 0^+} \tan x^{\tan x} = e^0 = 1$$

Note this is the same as $\lim_{u \rightarrow 0^+} u^u$

$$\text{b) } \int \sinh^{-1} x dx = x \sinh^{-1} x - \int \frac{x dx}{\sqrt{1+x^2}} \quad (1,5) + (1,5)$$

$$= x \sinh^{-1} x - \sqrt{1+x^2} + C$$

$$\text{c) } \int \tan^5 x \sec x = \int \tan^4 x \sec x dx = \int (\sec^2 x - 1)^2 \sec x \tan x dx$$

$$= \frac{(\sec x)^5}{5} - \frac{2}{3} \sec^3 x + \sec x + C \quad (1) + (1) + (1)$$

Question2

$$\text{a) } \int \frac{dx}{\sqrt{-x^2+6x-5}} = \int \frac{dx}{\sqrt{4-(x-3)^2}} = \sin^{-1} \frac{(x-3)}{2} + C \quad (1) + (1)$$

$$\text{b) } \int \frac{\sqrt{9-x^2}}{x^2} dx = \int \frac{9 \cos^2 \theta d\theta}{9 \sin^2 \theta} = \int (\csc^2 \theta - 1) d\theta \quad x = 3 \sin \theta$$

$$= -\cot \theta - \theta + C$$

$$= -\frac{\sqrt{9-x^2}}{x} - \sin^{-1} \left(\frac{x}{3} \right) + C \quad (1) + (1) + (1)$$

$$c) \frac{2x^2 - x + 2}{x(x^2 + 2)} = \frac{1}{x} + \frac{x-1}{x^2+2} \quad (1, 5) + (1, 5)$$

$$\int \frac{(2x^2 - x + 2)dx}{x(x^2 + 2)} = \ln |x| + \frac{1}{2} \ln(x^2 + 2) - \frac{1}{\sqrt{2}} \tan^{-1} \left(\frac{x}{\sqrt{2}} \right) + C$$

Question 3

$$a) \int \frac{dx}{1 - \sin x} = 2 \int \frac{du}{u^2 - 2u + 1} \quad u = \tan \left(\frac{x}{2} \right), dx = \frac{2du}{1+u^2} \quad (1, 5)$$

$$= 2 \int \frac{du}{(u-1)^2} = \frac{-2}{u-1} + C = \frac{-2}{\tan \left(\frac{x}{2} \right) - 1} + C \quad (1, 5)$$

$$b) \int_c^0 x e^x dx = [x e^x - e^x]_c^0 = -1 - c e^c + e^c \quad (1)$$

$$\lim_{c \rightarrow -\infty} (-1 - c e^c + e^c) = -1,$$

$$\text{so } \int_{-\infty}^0 x e^x dx \text{ cv and } \int_{-\infty}^0 x e^x dx = -1 \quad (1)$$

$$c) \text{ Intersection points: } x^2 - 8 = -x^2 \text{ so } x = \pm 2 \quad (0, 5)$$

$$A = 2 \int_0^2 [-x^2 - (x^2 - 8)] dx = 2 \left[-\frac{2}{3} x^3 + 8x \right]_0^2 = \frac{64}{3} \quad (1, 5)$$

