

Question 1 :

1. $AB + AC - D = 0 \iff A(B + C) = D \Rightarrow |A| |B + C| = |D|.$

$$B + C = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 1 \\ 0 & 0 & 3 \end{pmatrix}, \text{ then } |B + C| = -3 \text{ and } |A| = -2.$$

2. $RS + R - 2I = 0 \iff R(S + I) = 2I.$ Then

$$R^{-1} = \frac{1}{2}(S + I) = \begin{pmatrix} 1 & 0 & 1 \\ 1 & 2 & 2 \\ 0 & 1 & 3 \end{pmatrix}.$$

3. $a - b - 2c - 3d = 0 \iff a = b + 2c + 3d.$ The matrices in W are in the form

$$\begin{pmatrix} b + 2c + 3d & b \\ c & d \end{pmatrix} = b \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix} + c \begin{pmatrix} 2 & 0 \\ 1 & 0 \end{pmatrix} + d \begin{pmatrix} 3 & 0 \\ 0 & 1 \end{pmatrix}.$$

Then $\left\{ \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 2 & 0 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 3 & 0 \\ 0 & 1 \end{pmatrix} \right\}$ is a basis of the vector subspace W .

Question 2 :

The extended matrix of the system is: $\left[\begin{array}{ccc|c} 1 & m & 2 & 3 \\ 4 & 6 + m & -m & 13 - m \\ 1 & 2(m - 1) & m + 4 & m + 2 \end{array} \right].$

The matrix $\left[\begin{array}{ccc|c} 1 & m & 2 & 3 \\ 0 & m - 2 & m + 2 & m - 1 \\ 0 & 0 & 2(m - 1) & m - 1 \end{array} \right]$ is row equivalent to the extended matrix of the system.

- a) If $m \neq 1$ and $m \neq 2$ the system has a unique solution.
- b) If $m = 1$ the system has infinite solutions.
- c) If $m = 2$ the system has no solution.

Question 3 :

1. If $u_1 - 2u_2 + 3u_6 = 5u_3 + 7u_4 - 6u_5$, then $u_1 - 2u_2 - 5u_3 - 7u_4 + 6u_5 + 3u_6 = 0$. This is a linear combination of the vectors $u_1, u_2, u_3, u_4, u_5, u_6$ which are linearly independent. This is impossible. Then $u_1 - 2u_2 + 3u_6 \neq 5u_3 + 7u_4 - 6u_5$.

2. (a) The standard matrix of T is $\begin{pmatrix} 1 & -2 & 1 & 3 \\ 2 & -3 & 0 & 2 \\ -1 & 0 & 3 & 5 \end{pmatrix}.$

(b) The reduced echelon form of the matrix of T is $\begin{pmatrix} 1 & 0 & -3 & -5 \\ 0 & 1 & -2 & -4 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$ Then

$\{(3, 2, 1, 0), (5, 4, 0, 1)\}$ is a basis for kernel T .

(c) Using the reduced echelon form of the matrix of T we deduce that $\{(1, 2, -1), (2, 3, 0)\}$ is a basis for Image T .

Question 4 :

1. ${}_B P_C = \begin{pmatrix} 1 & 0 & 0 \\ -1 & 1 & 0 \\ 0 & -1 & 1 \end{pmatrix}$. $[v]_B = {}_B P_C [v]_C = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$

2. $[T(w)]_B = [T]_B [w]_B = \begin{pmatrix} 1 \\ -3 \\ -2 \end{pmatrix}$.

3. $T(u_1) = u_1 + u_2 + 2u_3 = au_1 - \frac{b}{5}u_2 + cu_3$. Then $a = 1, b = -5, c = 2$.

Question 5 :

1. $u_1 = \frac{1}{\sqrt{2}}(1, 1, 0)$, $\langle v_2, u_1 \rangle = \sqrt{2}$. Then $u_2 = (0, 0, 1)$.

2. (a) $\|u + v\|^2 = 11$.

(b) $\langle u, v \rangle = 0$, then $\cos \theta = 0$ and $\theta = \frac{\pi}{2}$.

Question 6 :

1. The eigenvalues of B are 1, 2. B is diagonalizable. $D = \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}$,

$P = \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}$ and $B = PDP^{-1}$. Then $B^{10} = PD^{10}P^{-1} = \begin{pmatrix} 1 & 2^{11} - 2 \\ 0 & 2^{10} \end{pmatrix}$.

2. Let $A = \begin{pmatrix} 1 & -1 & 3 \\ 0 & 1 & 0 \\ 0 & 3 & -2 \end{pmatrix}$.

(a) The characteristic polynomial of A is $q_A(\lambda) = (1 - \lambda)^2(2 + \lambda)$.

(b) The eigenvalues of A are 1 and -2 .

The eigenspace E_1 is generated by the vector $(1, 0, 0)$

and the eigenspace E_{-2} is generated by the vector $(1, 0, -1)$.

(c) A is not diagonalizable since $\dim(E_1) = 1$ and the algebraic multiplicity of 1 is 2.