

Reprinted from

PHARMACEUTICA ACTA HELVETIAE

Pharmaceutica Acta Helvetiae 73 (1999) 247–250

Effect of valproic acid on the pharmacokinetic profile of oxcarbazepine in the rat

K.M. Matar ^{a,*}, P.J. Nicholls ^b, S.A. Bawazir ^a, M.I. Al-Hassan ^a, A. Tekle ^a

^a Department of Clinical Pharmacy, College of Pharmacy, King Saud University, P.O. Box 2457, Riyadh 11451, Saudi Arabia

^b Division of Pharmacology, Welsh School of Pharmacy, University of Wales, Cardiff, Wales, UK



ELSEVIER

PHARMACEUTICA ACTA HELVETIAE

OFFICIAL JOURNAL OF THE SWISS PHARMACEUTICAL SOCIETY

Editor-in-Chief

Gerd Folkers, *Pharmaceutica Acta Helvetiae*, Departement Pharmazie, ETH Zürich-Irchel, Winterthurerstr. 190, CH-8057 Zürich, Switzerland. Telephone: (+41-1) 6356060, Fax: (+41-1) 6356884, E-mail: folkers@pharma.ethz.ch.

Associate Editors

H.P. Merkle Zürich, Switzerland	O. Sticher Zürich, Switzerland	P. Speiser Zürich, Switzerland	Y. Arata Tokyo, Japan	H. Schreier Novato, CA, USA
------------------------------------	-----------------------------------	-----------------------------------	--------------------------	--------------------------------

Editorial Advisory Board

D. Abraham (Richmond)	T. Kissel (Marburg/Lahn)	U. Rüegg (Lausanne)
T.L. Clerc (Bern)	H. Kubinyi (Ludwigshafen)	N. Tüller (Bern)
Th. Dingermann (Frankfurt/Main)	M. Mesnil (Bern-Liebefeld)	S. van Boeckel (Oss)
H. Fenner (Gelterkinden)	E. Meyer (College Station)	H.P. Weber (Dornach)
R. Gurny (Genève)	E. Mutschler (Frankfurt/Main)	H. Wunderli (Zürich)
O. Gutbrod (Leverkusen)	V. Pliska (Zürich)	

Aims and Scope: *Pharmaceutica Acta Helvetiae* covers all disciplines important for and related to pharmaceutical research, from discovery of a drug up to clinical pharmacy practice. Topics may be: analytics, biochemistry, biopharmaceutics, biotechnology, cell biology, cell cultures, clinical pharmacy, crystallography, drug delivery, drug disposition, gene technology, medicinal chemistry, metabolism, molecular modelling, molecular pharmacology, peptide and protein chemistry, pharmaceutical technology, pharmacognosy, pharmacology, pharmacokinetics, physical pharmacy, protein design, synthesis, theoretical chemistry and toxicology.

Manuscripts in the following categories will be considered for publication:

Reviews: Reviews are invited papers dealing with recent findings, new methodologies and alternative views of great importance for pharmaceutical sciences.

Articles: Articles are original papers of research in the fields mentioned above. They are expected to present new findings of considerable importance and preference is given to multidisciplinary approaches.

Communications: Communications are short papers of outstanding importance for the aforementioned research fields. Preparation on diskette is required. Instructions are available from the Publisher.

Publication information: *Pharmaceutica Acta Helvetiae* (ISSN 0031-6865). For 1999 volume 74 is scheduled for publication. Subscription prices are available upon request from the Publisher. Subscriptions are accepted on a prepaid basis only and are entered on a calendar year basis. Issues are sent by surface mail except to the following countries where Air delivery via SAL mail is ensured: Argentina, Australia, Brazil, Canada, Hong Kong, India, Israel, Japan, Malaysia, Mexico, New Zealand, Pakistan, PR China, Singapore, South Africa, South Korea, Taiwan, Thailand, USA. For all other countries airmail rates are available upon request. Claims for missing issues should be made within six months of our publication (mailing) date. Special price for members of the Swiss Pharmaceutical Society: Dfl. 125.00 including postage and handling. Information available from the Society. Contact person: Dr. M. Mesnil, Société Suisse de Pharmacie, Stationsstrasse 12, CH-3097 Bern-Liebefeld, Switzerland.

Back volumes: Please contact the Society.

© The paper used in this publication meets the requirements of ANSI/NISO Z39.48-1992 (Permanence of Paper).

Printed in The Netherlands.

Effect of valproic acid on the pharmacokinetic profile of oxcarbazepine in the rat

K.M. Matar ^{a,*}, P.J. Nicholls ^b, S.A. Bawazir ^a, M.I. Al-Hassan ^a, A. Tekle ^a

^a Department of Clinical Pharmacy, College of Pharmacy, King Saud University, P.O. Box 2457, Riyadh 11451, Saudi Arabia

^b Division of Pharmacology, Welsh School of Pharmacy, University of Wales, Cardiff, Wales, UK

Received 30 July 1998; revised 23 October 1998; accepted 28 October 1998

Abstract

The pharmacokinetics of oxcarbazepine (30 mg kg⁻¹, po), administered for 1 week, was studied in rats pre-treated for 2 weeks with valproic acid (100 mg kg⁻¹, po). Oxcarbazepine (OXC) plasma levels were measured over a period of 24 h from dosing, using a sensitive HPLC method. No significant changes were observed in the mean values of OXC pharmacokinetic parameters (C_{max} , T_{max} , $t_{1/2}$ and $AUC_{0-\infty}$) between the control and the pre-treated groups. The findings of this study suggest that OXC metabolism in the rat is apparently not affected by valproic acid, and the lack of effect may be attributed to the different pathways of biotransformation of the two drugs. © 1999 Published by Elsevier Science B.V. All rights reserved.

Keywords: Oxcarbazepine; Valproic acid; Pharmacokinetic interaction; Rat

1. Introduction

Oxcarbazepine (OXC), the 10,11-dihydro-10-oxo-carbamazepine, is a new antiepileptic drug having the same anticonvulsant activity as carbamazepine (CBZ) but a completely different metabolic profile. In humans, OXC is rapidly reduced by a cytosolic aldo-keto reductase enzyme to the active metabolite, 10-hydroxy-carbamazepine (MHD), Fig. 1 (Faigle and Menge, 1990). The bulk of this metabolite undergoes conjugation reaction with a glucuronyl moiety through the actions of microsomal glucuronyl-transferase prior to its excretion from the body, while a small fraction is further oxidized to an inactive 10,11-*trans*-dihydroxymetabolite (DHD) common to both OXC and CBZ (Lloyd et al., 1994). In rats, the reduction of OXC to MHD plays a minor role and oxidative reactions predominate (Wagner and Schmid, 1987). CBZ, on

the other hand, is mainly oxidized in the liver to its active metabolite, carbamazepine-10-11-epoxide (CBZ-E) by microsomal mono-oxygenase oxidative enzymes of the cytochrome *P*-450 system (Eadie, 1991). The OXC metabolic profile suggests that it may have fewer drug interactions in man compared with CBZ.

During concomitant administration of valproic acid (VPA) with other antiepileptic drugs (AEDs), VPA has been observed to influence pharmacokinetics of phenobarbitone, phenytoin, ethosuximide and CBZ (Riva et al., 1996). Unlike the other AEDs, VPA is an inhibitor of hepatic oxidative metabolism. A cytochrome *P*-450 microsomal enzyme system appears to rapidly metabolize VPA in the liver. The drug undergoes oxidation reactions to metabolites which are finally conjugated for clearance from the body (Eadie, 1991). The coadministration of VPA with CBZ is very effective in the management of epilepsy; but it has been reported to precipitate CBZ toxicity (Ketter et al., 1991). In this regard VPA was shown in some studies not to have significant influence on plasma CBZ concentrations but it did increase the levels of the main

* Corresponding author. Tel.: +966-1467-7448; fax: +966-1476-6383




Fig. 1. Metabolic pathways of oxcarbazepine in man (A) and rat (B).

metabolite, CBZ-E (Robbins et al., 1990). The purpose of this study was to investigate the possible influence of repeated doses of VPA on the pharmacokinetics of OXC, a drug closely related to CBZ, in rats.

2. Materials and methods

2.1. Materials

OXC, 10-OH-CBZ and 10-11-epoxycarbamazepine were kindly supplied by Ciba-Geigy (Basle, Switzerland). Sodium valproate, 40 mg ml⁻¹ syrup (Epilim®) was from Sanofi Pharma (Manchester, UK). Carboxymethylcellulose (CMC) was purchased from BDH Chemicals (Poole, UK). Acetonitrile (HPLC grade) was purchased from E. Merck (Darmstadt, Germany). HPLC water was prepared using a Milli-Q Water System (Millipore, Bedford, MA, USA). All other reagents were of analytical grade.

2.2. Subjects and experimental procedure

Adult male Sprague-Dawley rats weighing 250–300 g were used in this study. Animals were randomly divided into two (treatment and control) groups. OXC solutions were freshly prepared in 1% carboxymethylcellulose (CMC), and sodium valproate was used as the commercial preparation, 40 mg ml⁻¹ syrup (Epilim®), and were administered orally with a feeding needle.

The treatment group ($n = 12$), received VPA (100 mg kg⁻¹, po) daily for 7 days and then in combination with

OXC (30 mg kg⁻¹, po) for another week. On the other hand, the control group ($n = 12$) received 1% CMC for 1 week and then OXC (30 mg kg⁻¹, po) was given daily for another week. At the end of the 14-day treatment period, the animals were fasted overnight (water given ad libitum), and cannulation of the left femoral artery with polythene tubing (0.50 mm I.D., 1.0 mm O.D., Portex, Hythe, Kent, UK), was carried out under light ether anaesthesia. The other end of the tubing was passed under the emerging at the back of the neck. After flushing it with heparinized saline, the open end of the cannula was closed with a pin plug. Following the surgical procedure, the animals were kept in individual cages and allowed to recover from anaesthesia.

After recovery, the animals were given oral doses of OXC (30 mg kg⁻¹, po) alone or in combination with VPA, (100 mg kg⁻¹, po) and returned to their restraining cages. During the subsequent serial sampling of blood, a few drops of blood were routinely discarded to avoid potential contamination and/or dilution with the heparinized saline filling the length of the cannula. In this manner, blood samples (ca. 0.3 ml) for the measurement of OXC were drawn through the indwelling cannula into small plastic centrifuge tubes (LIP, Yorkshire, UK) just before and 0.5, 1, 2, 4, 6, 9, 12, 24 and 36 h after OXC dosing. The cannula was flushed with an equal volume of normal saline to maintain the circulatory volume in the animal. The blood samples were then immediately centrifuged at 3000 rpm for 5 min, and aliquotes 100 µl of plasma samples were stored at -20°C until assay.

2.3. Drug analysis

OXC was determined using a previously described HPLC method (Matar et al., 1995). Briefly, to 0.1 ml of rat plasma, 20 μ l of 10,11-epoxycarbamazepine as internal standard (2 μ g ml⁻¹) and 0.5 ml of dichloromethane were added.

Chromatography was performed on a reverse phase Novapak C₁₈ column (4 μ m, 150 mm \times 3.9 mm I.D.). The mobile phase consisted of a mixture of 20% v/v acetonitrile in water and the pump (M 501) flow rate was 1.5 ml/min. The effluent was monitored at 215 nm with a variable wavelength UV detector (M 481).

The lower limit of detection of OXC was 50 ng/ml. The interday coefficient of variation ranged between 6.3 to 8.3%.

2.4. Pharmacokinetic and statistical analysis

Pharmacokinetic parameters for OXC were calculated. These included maximum plasma concentration (C_{max}) and time to attain the maximum concentration (T_{max}) which were directly determined from the resulting concentration–time profiles. The elimination rate constant (K_{el}) was calculated from log-linear least-squares regression of the terminal phase data points.

The elimination half-life ($t_{1/2}$) was calculated as 0.693 divided by the elimination rate constant. Non-compartmental model was applied to calculate the area under the plasma concentration–time curve (AUC), which was determined by the linear trapezoidal rule. The area from the last point to infinity was determined by dividing the last plasma concentration by the elimination rate constant. The




Fig. 2. Mean (\pm S.E.M.) plasma concentration–time profile of oxcarbazepine (o) when administered alone (30 mg kg⁻¹, po) and after pre-treatment with (100 mg kg⁻¹) valproic acid (•) in rats ($n = 12$).

Table 1

Pharmacokinetic parameters (mean \pm S.E.M) for OXC (30 mg kg⁻¹, po) administered alone and in combination with VPA (100 mg kg⁻¹, po) in rats ($n = 12$)

Pharmacokinetic parameter	OXC alone	OXC with VPA
C_{max} (μ g ml ⁻¹)	4.36 \pm 0.43	3.21 \pm 0.52 *
T_{max} (h)	1.94 \pm 0.37	2.53 \pm 0.34 *
$t_{1/2}$ (h)	11.74 \pm 2.00	9.75 \pm 1.76 *
AUC _{0-∞} (μ g h ml ⁻¹)	46.99 \pm 6.65	36.46 \pm 5.70 *

* Student's t -test, $P > 0.05$.

programme used for the determination of the OXC pharmacokinetic parameters was Kinetica software 2.0.1, 1996–2006 (MicroPharm International, USA). Pharmacokinetic parameters are presented as (mean \pm S.E.M.).

The significance level chosen was $P < 0.05$ using a two-tailed unpaired Student's t -test for comparing two means. Evaluation was performed using STAT100 software, version 1.24, 1995–1996 (Biosoft, Cambridge, UK).

3. Results

The mean plasma concentration–time profiles of OXC after oral administration of OXC (30 mg kg⁻¹) both alone or in combinations with VPA (100 mg kg⁻¹, po) are shown in Fig. 2. The computed pharmacokinetic parameters are presented in Table 1. VPA did not produce any significant effect on the mean value of OXC pharmacokinetic parameters. Slight but not statistically significant decrease was observed in the maximum plasma concentration during pre-treatment with VPA.

4. Discussion

OXC is chemically related to CBZ but it has completely different metabolic profile. In humans, OXC is rapidly and almost completely reduced to its active metabolite (MHD) with only a minimal amount of the parent drug being found in the blood. This metabolite may partially undergo a further reduction to the DHD form prior to its systemic clearance as a glucuronide conjugate (Lloyd et al., 1994). Therefore, the pharmacokinetics of OXC and MHD are completely controlled by two non-oxidative enzymatic processes (reduction and conjugation). In rats, this reductive biotransformation plays a minor role and levels of the parent drug greatly exceed MHD concentration in plasma (Wagner and Schmid, 1987). The dependent conversion of MHD to DHD plays a minor role in humans, but it

constitutes a major metabolic pathway in rats. By contrast, the principal metabolic route for the structurally related drug, CBZ, involves a microsomal oxidation to its active metabolite (CBZ-E), and subsequent hydrolysis of the latter to an inactive metabolite, *trans*-10,11-dihydroxy-10-11-dihydrocarbamazepine (CBZ-diol).

The metabolic route for VPA involves initial oxidation by the microsomal cytochrome *P*-450 system followed by conjugation to glucuronic acid derivative. Rembeck et al. (1987) reported a decrease in CBZ levels when VPA was concomitantly used with CBZ. Another study reported a significant increase in CBZ-E levels due to inhibition of epoxide hydrolase by VPA (Robbins et al., 1990).

The objective of this study was to find the potential interaction of VPA with OXC, a drug closely related to CBZ. Despite its close structural similarity to CBZ, OXC exhibits a completely different metabolic profile to that of CBZ in both human and laboratory animals. The results of the present study indicate that pre-treatment with VPA did not significantly alter the pharmacokinetics of OXC in the rat. In addition to that, the plasma levels of MHD had not been followed because most of its levels were below the detection limit (20 ng ml^{-1}). This finding is in a good agreement with an earlier report on the interaction of OXC with VPA in healthy subjects and epileptic patients (Tartara et al., 1993). This lack of change in plasma OXC levels in rats pretreated with VPA could be explained on the different metabolic pathways of OXC, which is mainly reduction while VPA undergoes oxidation with a microsomal *P*-450 system.

References

- Eadie, M., 1991. Formation of active metabolites of anticonvulsant drugs; a review of their pharmacokinetic and therapeutic significance. *Clin. Pharmacokinet.* 21, 27–41.
- Faigle, J., Menge, G., 1990. Metabolic characteristics of oxcarbazepine (Trileptal®) and their beneficial implications for enzyme induction and drug interactions. *Behav. Neurol.* 3, 21–30, Suppl. 1.
- Ketter, T., Post, R., Worthington, K., 1991. Principles of clinically important drug interactions with carbamazepine: Part 1. *J. Clin. Psychopharmacol.* 11, 198–203.
- Lloyd, P., Flesch, G., Dieterle, W., 1994. Clinical pharmacology and pharmacokinetics of oxcarbazepine. *Epilepsia* 35, S10–S13, Suppl. 3.
- Matar, K.M., Nicholls, P.J., Al-Hassan, M.I., Tekle, A., 1995. Rapid micromethod for simultaneous measurement of oxcarbazepine and its active metabolite in plasma by high performance liquid chromatography. *J. Clin. Pharm. Ther.* 20, 229–234.
- Rembeck, B., May, T., Juergens, U., 1987. Serum concentrations of carbamazepine and its epoxide and diol metabolites in epileptic patients: the influence of dose and comedication. *Ther. Drug. Monit.* 9, 298–303.
- Riva, R., Albani, F., Contin, M., Baruzzi, A., 1996. Pharmacokinetic interactions between antiepileptic drugs; clinical considerations. *Clin. Pharmacokinet.* 31, 470–493.
- Robbins, D., Wedlund, P., Kuhri, R., Baumann, R., Levy, R., Chang, S., 1990. Inhibition of epoxide hydrolase by valproic acid in epileptic patients receiving carbamazepine. *Br. J. Clin. Pharmacol.* 29, 759–762.
- Tartara, A., Galimberti, C.A., Manni, R., Morini, R., Limido, G., Gatti, G., Bartoli, A., 1993. The pharmacokinetics of oxcarbazepine and its active metabolite 10-hydroxy-carbamazepine in health subjects and in epileptic patients treated with phenobarbitone or valproic acid. *Br. J. Clin. Pharmacol.* 36, 366–368.
- Wagner, J., Schmid, K., 1987. Induction of microsomal enzymes in rat liver by oxcarbazepine, 10,11-dihydro-10-hydroxy-carbamazepine and carbamazepine. *Xenobiotica* 17, 951–956.

PHARMACEUTICA ACTA HELVETIAE

Notes to Contributors

Submission procedure: Submission of a paper to *Pharmaceutica Acta Helvetiae* is understood to imply that the data contained therein have not previously been published (except in abstract form) nor is being considered for publication elsewhere, and that it has been tacitly or explicitly approved by the responsible authorities in the laboratory where the work was carried out.

Manuscripts submitted under multiple authorship are reviewed under the assumption that all listed authors concur with the submission and have approved the final manuscript. If accepted, the papers shall not be published either whole or in part elsewhere in the same or in any other language without the consent of the Publisher.

Pharmaceutica Acta Helvetiae has no page charges.

Only submissions in English will be considered. One original plus three copies (copy sets of photographs must be original prints) should be submitted to: Prof. Dr. G. Folkers, Departement Pharmazie, ETH Zürich-Irchel, Winterthurerstr. 190, CH-8057 Zürich, Switzerland. Copies of submitted manuscripts will not be returned except in the case of rejected articles, where only the original manuscript and figures are returned to the author. Papers that do not conform to the general criteria for publication in *Pharmaceutica Acta Helvetiae* will be returned immediately to authors to avoid unnecessary delay in submission elsewhere. The decision of the Editor is final: the Editor cannot enter into correspondence about a paper that is considered unsuitable for publication.

Electronic manuscripts: Electronic manuscripts have the advantage that there is no need for the rekeying of text, thereby avoiding the possibility of introducing errors and resulting in reliable and fast delivery of proofs.

For the initial submission of manuscripts for consideration, hardcopies are sufficient. For the processing of **accepted papers**, electronic versions are preferred. After **final acceptance**, your disk plus three, final and exactly matching printed versions should be submitted together. Double density (DD) or high density (HD) diskettes (3 $\frac{1}{2}$ or 5 $\frac{1}{4}$ inch) are acceptable. It is important that the file saved is in the native format of the wordprocessor program used. Label the disk with the name of the computer and wordprocessing package used, your name, and the name of the file on the disk. Further information may be obtained from the Publisher.

Authors in Japan please note: Upon request, Elsevier Science K.K. will provide authors with a list of people who can check and improve the English of their paper (*before submission*). Please contact our Tokyo office: Elsevier Science K.K., 1-9-15 Higashi-Azabu, Minato-ku, 106-0044 Tokyo; Tel. (81) (3) 5561-5032; fax (81) (3) 5561-5045.

A complete **Guide for Authors** appears in the journal each year and can be found in volume 73, no. 5 on pages 261-262.

The instructions can also be found on the World Wide Web: access under <http://www.elsevier.nl> or <http://www.elsevier.com>

Enquiries concerning manuscripts and proofs: questions arising after acceptance of a manuscript by the editors, especially those relating to proofs, publication and reprints, should be directed to Elsevier Science Ireland Ltd., Elsevier House, Brookvale Plaza, East Park, Shannon, Co. Clare, Ireland. Tel.: +353 61 709652; fax: +353 61 709110.

Advertising information. Advertising orders and enquiries can be sent to: **USA, Canada and South America:** Mr Tino DeCarlo, The Advertising Department, Elsevier Science Inc., 655 Avenue of the Americas, New York, NY 10010-5107, USA; Tel.: (+1) (212) 633 3815; fax: (+1) (212) 633 3820; e-mail: t.decarlo@elsevier.com. **Japan:** The Advertising Department, Elsevier Science K.K., 9-15 Higashi-Azabu 1-chome, Minato-ku, Tokyo 106-0044, Japan; Tel.: (+81) (3) 5561 5033; fax: (+81) (3) 5561 5047. **Europe and ROW:** Rachel Gresle-Farthing, The Advertising Department, Elsevier Science Ltd., The Boulevard, Langford Lane, Kidlington, Oxford OX5 1GB, UK; Tel.: (+44) (1865) 843565; fax: (+44) (1865) 843976; e-mail: r.gresle-farthing@elsevier.co.uk.

Orders, claims, and product enquiries: please contact the Customer Support Department at the Regional Sales Office nearest you:

New York: Elsevier Science, PO Box 945, New York, NY 10159-0945, USA; Tel.: (+1) (212) 633 3730 [toll free number for North American customers: 1-888-4ES-INFO (437-4636)]; fax: (+1) (212) 633 3680; e-mail: usinfo-f@elsevier.com

Amsterdam: Elsevier Science, PO Box 211, 1000 AE Amsterdam, The Netherlands; Tel.: (+31) 20 485 3757; fax: (+31) 20 485 3432; e-mail: nlinfo-f@elsevier.nl

Tokyo: Elsevier Science, 9-15 Higashi-Azabu 1-chome, Minato-ku, Tokyo 106-0044, Japan; Tel.: (+81) (3) 5561 5033; fax: (+81) (3) 5561 5047; e-mail: info@elsevier.co.jp

Singapore: Elsevier Science, No. 1 Temasek Avenue, #17-01 Millenia Tower, Singapore 039192; Tel.: (+65) 434 3727; fax: (+65) 337 2230; e-mail: asiainfo@elsevier.com.sg

Rio de Janeiro: Elsevier Science, Rua Sete de Setembro 111/16 Andar, 20050-002 Centro, Rio de Janeiro - RJ, Brazil; Tel.: (+55) (21) 509 5340; fax: (+55) (21) 507 1991; e-mail: elsevier@campus.com.br [Note (Latin America): for orders, claims and help desk information, please contact the Regional Sales Office in New York as listed above]